This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

+ *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
+ *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
+ *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
+ *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at `http://books.google.com/`

3 2044 106 400 856

H831s
cop.2

W. G. FARLOW

H831s c.2

Harvard University

VE RI TAS

FARLOW
REFERENCE LIBRARY
OF
CRYPTOGAMIC BOTANY

- A 115 -

N.° 23.

NICOLAI THOMÆ HOST
MEDIC. DOCT.

SYNOPSIS PLANTARUM

IN

AUSTRIA

PROVINCIISQUE ADIACENTIBUS

SPONTE CRESCENTIUM.



VINDOBONÆ.

SUMPTIBUS CHRIST. FRIEDER. WAPPLER.

1797.

PRÆFATIO.

Enumerat Synopſis hæc ſtirpes, quas mihi in excurſionibus per Auſtriam inferiorem finitimamque Pannoniam; in itineribus per Styriam, Carinthiam, Tyrolim, Forojulium, Iſtriam, Carnioliam, Litorale finitimaque Croatiæ loca videre contigit. Exhibet plantas, quas, juſſu auſpiciisque FRANCISCI II. a me lectas, in hortum, AUGUSTISSIMI munificentia pro Floræ Auſtriacæ ſede extructum, repoſitas, jam per triennium omni anni tempore ſtudioſius examinare mihi licuit. Accedunt hisce permultæ, a me non viſæ, quas fide digniſſimi viri invenerunt.

In ordinandis ſtirpibus Linnæi veſtigiis inhæſi; ſtatutumque ab eo Syſtema, etiamſi undequaque minime perfectum, conſectari volui.

Genera, ut in Editione Generum Plantarum octava curante Cl. Schreber ordinata proſtant, paucis mutatis, ordinavi.

Præſtantiſſimorum virorum deſcriptiones in auxilium vocavi; ſynonyma ſelectiora adduxi.

Cum Fungi multum ſibi temporis vindicent; nec non longam operam, atque acriorem ſolertiam expoſcant, hos obiter pertractavi.

Arbori Arb., Frutici Fr., Perenni plantæ Per. appoſui, ſi id vel nomen triviale vel differentia ſpecifica non indicavit.

Annuis, biennibus durationis tempus; Muſcis vero, Hepaticis, Algis, Fungis tam durationis quam florendi tempus apponere ſuperfluum duxi.

— ❁ —

Me pauca hic præſtitiſſe; multa hic deſiderari, ingenue fateor. At palam quoque profiteri liceat, quod nec hæc præſtare valuiſſem, quodque plura deſiderarentur, niſi Cl. Viri, Iacquin Pater, Profeſſor meus in æternum venerandus, & Filius pro ſumma erga me measque res benevolentia in dubiis explicandis maxime benignos, in obſervationibus, plantis libriſque communicandis ſumme urbanos ſeſe præbuiſſent. Nec ſilentio præterire licet Ill. L. B. Carolum a Zois, Reverendum Dominum, jam ab incunabulis plus quam fratrem, Ioſephum Hoſt, quibus ſimilia officiorum genera accepta referre non piget. Perpetuo hærent hærebuntque animo impreſſa optimæ voluntatis indicia, quibus ille me excepit; quibus, alioquin ſumme arduum ſummeque aſperum, wochinenſium aliarumque carniolicarum alpium aſcenſum faciliorem reddere conatus eſt; quibus præſentem oretenus, per litteras abſentem docere ſtuduit. Communicavit hic rariſſimas, a me nondum inventas, ſtirpes, quas in arduis itineribus, ob Floræ amorem per *Li*torale, Iſtriam, Inſulas venetas, Forojulium finitimamque Carniam ſuſceptis, collegit; cumque ſtir-

pibus communicavit mecum, a ſe factas, obſervationes, Sit æternæ gratitudinis monumentum!

Tu autem Benevole Lector hos meos labores æqui bonique conſulito. Ego quod hic tum ob temporis anguſtias, tum aliarum occupationum copiam præſtare non valui, in poſterum præſtare ſatagam, ſi placere ſenſero officium meum.

AUCTORES CITATI.

Ait. kew. Hortus kewenſis; or a Catalogue of the Plants cultivated in the Royal Garden at Kew. By Wm. Aiton. 3 vol. London. 1789. 8vo.

Allion. nic. Caroli Allionii Stirpium Nicenſium Enumeratio. Pariſiis. 8vo.

Allion. pedem. Ejusdem Flora Pedemontana. Tomi 3. Auguſtæ Taurinorum. 1785. fol.

Ambros. Hyacinthi Ambroſini Phytologiæ partis primæ Tomus. 1. Bononiæ. 1666. fol.

Arduin. animadv. P. Arduini animadverſionum botanicarum ſpecimen 1. Patavii. 1759. Specimen 2. Venetiis. 1764. 4to.

Barr. ic. I. Barrelieri Plantæ per Galliam, Hiſpaniam, & Italiam obſervatæ. Pariſiis. 1714. fol.

Batſch. fung. Elenchus fungorum, conſcripſit A. I. G. C. Batſch Halæ Magdeburgicæ. 1783. &c. 4to.

Batt. fung. Fungorum agri ariminenſis hiſtoria. a I. A. Battarra. Faventiæ. 1759. 4to.

Bauh. hiſt. Hiſtoria plantarum univerſalis. Ioh. Bauhini & I. H. Cherleri. Tomi 3. Ebroduni. 1650. 1651. fol.

Bauh. pin. Caſp. Bauhini pinax Theatri botanici. Baſileæ. 1671. 4to.

Besl. eyſt. B. Besleri Hortus Eyſtettenſis. 1613. fol.

Blackw. herb. A. curioſus Herbal. By Eliz. Blackwell. 2 vol. London. 1759. fol.

Bocc. muſ. Muſeo di piante rare di Don Paulo Boccone. Venezia. 1697. 4to.

Bocc. ſic. Ejusdem Icones, & Deſcriptiones rariorum Plantarum Siciliæ, Melitæ, Galliæ, Italiæ. Oxonii. 1674. 4to.

Bolt. fil. Filices britanicæ. By Iames Bolton. With an Appendix. Huddersfield. 1790. 4to.

Bolt. fung. An Hiſtory of Funguſes growing about Halifax. By I. Bolton. 4 vol. Halifax. 1788. &c. fol.

Breyn. cent. I. Breynii Centuria 1 exoticarum, & minus cognitarum plantarum. Gedani. 1768. fol.

Burgsdorf abb. Vide Reitter und Abel.

Bux. cent. I. C. Buxbaumii Plantarum minus cognitarum Centuriæ 5. Petropoli. 1728. &c. 4to.

Cam. epit. De Plantis Epitome P. A. Matthioli, aucta a I. Camerario. Francofurti ad Mœnum. 1586. 4to.

Cam. hort. I. Camerarii hortus medicus & philoſophicus. Francofurti. 1588. 4to.

Cavan. diſſ. A. I. Cavanilles Diſſertationes botanicæ. Pariſiis. 1785. &c. 4to.

Clus. hiſt. C. Cluſii rariorum Plantarum Hiſtoria. Antverpiæ. 1601. fol.

Col. ecphr. F. Columnæ minus cognitarum Stirpium Ecphraſis. Romæ. 1616. 4to.

Col. phyt. Ejusdem Phytobaſanoc. Neapoli. 1561. fol.

Comm. gott. Commentationes Societatis Regiæ Scientiarum Gottingenſis. Gottingæ. 1779. &c. 4to.

Crantz faſc. N. Crantz Stirpium auſtriacarum faſciculi. 1769. vol. 2. 4to.

Curtis lond. Flora londinenſis. By. Wm. Curtis. London. fol.

Curtis magaz. Ejusdem botanical Magazine. London. 1787. &c. 8vo.

Dill. elth. — Ioh. Iac. Dillenii Hortus elthamenſis. vol. 2. Londini. 1732. fol.

Dill. giſſ. — Catalogus Plantarum ſponte circa Giſſam naſcentium cum Appendice. Francofurti. 1709. 8vo.

Dill. muſc. — Hiſtoria muſcorum. Oxonii. 1741. 4to.

Dod. pempt. Remberti Dodonæi Stirpium Hiſtoriæ Pemptades ſex. Antverpiæ. 1616. fol.

Dod. purg. Ejusdem purgantium & deleteriarum herbarum Hiſtoria. Antverpiæ. 1574. 4to.

Ehrh. beytr. Fr. Ehrhart Beyträge zur Naturkunde &c. 7 Bände. Hannover. 1787. &c. 8vo.

Fl. dan. Icones Plantarum ſponte nascentium in regnis Daniæ, Norvegiæ. Hafniæ. 1761. &c. fol.

Fl. lapp. C. Linnæi Flora Lapponica. Amſtellodami. 1737. 8vo.

Fl. N. I. E. Gunneri Flora Norvegica. Pars prior Nedroſiæ. 1766. Pars poſterior Hafniæ. 1772. fol.

Garid. aix. Garidel Hiſtoire des plantes, qui naiſſent aux environs d'Aix en Provence. Aix. 1715. fol.

Ger. prov. Lud. Gerarard Flora provincialis. Pariſiis. 1761. 8vo.

Gouan illuſtr. Ant. Gouan. Illuſtrationes & Obſervationes botanicæ. Tiguri. 1773. fol.

Gouan fl. — Flora monſpeliaca. Lugduni. 1765. 8vo.

Gouan hort. — Hortus regius monſpelienſis. Lugduni. 1762. 8vo.

Gmel. fuc. Gmelin Hiſtoria Fucorum Petropoli. 1768. 4to.

Gmel. ſib. Gmelin Flora Sibirica. Tomi 4. Petropoli. 1747. &c. 4to.

Guett. obs. Guettard Obſervations ſur les plantes. vol. 2. Paris. 1747. 8vo.

Hacquet pl. alp. Plantæ alpinæ carniolicæ. Collegit & deſcripſit Hacquet. 1782. 4to.

Hag. lich. C. G. Hagen Tentamen Hiſtoriæ Lichenum. Regiomonti. 1782. 8vo.

Hall. hiſt. Alberti v. Haller Hiſtoria ſtirpium indigenarum Helvetiæ inchoata. Tomi 3. Bernæ. 1768. fol.

Hall. nom. — Nomenclator ex Hiſtoria Plantarum excerptus. Bernæ. 1769. 8vo.

Han. magaz. Hannoveriſches Magazin. 1781. 4to.

Hedw. musc. Deſcriptio & Adumbratio muscorum frondoſorum. Auctore I. Hedwig. Tomi 4. Lipſiæ. 1787. &c. fol.

Herm. par. P. Hermanni Paradiſus Batavus. Lugduni Batavorum. 1705. 4to.

Hoffm. fl. Deutſchlands Flora von G. F. Hoffmann. vol. 2. Erlangen.

Hoffm. lich. — Deſcriptio & Adumbratio Plantarum, quæ lichenes dicuntur. vol. 2. Lipſiæ. fol.

Hoffm. ſal. — Hiſtoria ſalicum. Lipſiæ. 1787. fol.

Huds. angl. ed. 1. Gul. Hudſoni Flora anglica. Londini. 1762. 8vo.

Huds. angl. ed. 2. — Flora anglica. Editio altera. vol. 2. Londini. 1778. 8vo.

Iacqu. auſtr. N. I. Iacquin Flora Auſtriaca, vol. 5. Viennæ. 1773. &c. fol.

Iacqu. collect. — Collectanea ad Botanicam, Chemiam & Hiſtoriam naturalem ſpectantia. vol. 5. Vindobonæ. 1786. 4to.

Iacqu. en. — Enumeratio Stirpium, quæ ſponte creſcunt in agro vindobonenſi. Viennæ. 1762. 8vo.

Iacqu. hort. — Hortus botanicus vindobonenſis. Tomi 3. Viennæ. 1770. &c. fol.

Iacqu. ic. — Icones Plantarum rariorum. Tomi 3. Vindobonæ. 1781. &c. fol.

Iacqu. miscell. — Miscellanea auſtriaca. Vol. 2. Vindobonæ. 1778. &c. 4to.

Iacqu. obs. — Obſervationum botanicarum. Partes 4. Viennæ. 1764. &c. fol.

Iacqu. oxal. — Oxalis. Viennæ. 1794. 4to.

Koel. ſpic. Spicilegium obſervationum de Aconito. Auctore I. L. C. Koelle. Erlangæ. 1788. 8vo.

Kram. auſtr. Gul. Henr. Kramer Elenchus Vegetabilium & Animalium per Auſtriam inferiorem obſervatorum. Viennæ. 1756. 8vo.

Lam. fl. franc. Lamarck Flore françaiſe. Vol. 3. Paris. 1778. 8vo.

Leers herb. I. D. Leers Flora Herbornenſis. Editio altera. Coloniæ Alobrog. 1789. 8vo.

Leys. hal. F. W. a Leyſer Flora Halenſis. Halæ. 1783. 8vo.

L' Herit. corn. Car. Ludw. L' Heritier Cornus. Pariſiis. 1788. fol.

L' Herit. ger. — Geranologia. Pariſiis. 1788. fol.

L' Herit. ſtirp. nov. — Stirpes novæ deſcriptionibus & iconibus illuſtratæ. Pariſiis. 1784. &c.

Lindern hort. Hortus Alſaticus conſcriptus a F. B. von Lindern, Argentorati. 1747. 8vo.

Linn. cliff. C. Linnæi Hortus Cliffortianus. Amſtelodami. 1737. fol.

Linn. mant. — Mantiſſa Plantarum Generum editionis 6, & Specierum ed. 2. Holmiæ. 1767. 8vo.

Mantiſſa Plantarum altera. Holmiæ. 1771. 8vo.

Linn. ſuppl. Supplementum Plantarum Syſtematis Vegetabilium ed. 13. Generum Pl. ed. 6, & Specierum Pl. ed. 2. editum a C. a Linné filio. Brunsvigæ. 1781. 8vo.

Linn. trans. Transactions of the Linnean Society. 2 vol. London. 1791. &c. 4to.

Lob. adv. Stirpium Adverſaria nova, auctoribus Petro Pena, & Matthia de Lobel. Londini. 1571. fol.

Matthiæ de Lobel Adverſariorum altera pars. Londini. 1605. fol.

Lob. ic. M. Lobelii Adverſaria Stirpium. Antverpiæ. 1676. fol.

Loes. pruſſ. I. Loeſelii Flora Pruſſica. Regiomonti. 1703. 4to.

Magn. bot. Petri Magnol Botanicum Monſpelienſe. Monſpelii. 1697. 8vo.

Mentz. pug. Mentzelius in Pugillo rariorum Plantarum ad calcem Indicis. Berolini. 1682. fol.

Mich. gen. P. A. Michelii Nova Plantarum Genera. Florentiæ. 1729. fol.

Miller ic. Figures of Plants deſcribed in the Gardeners Dictionary, by Phil. Miller. 2 Vol. London. 1760. fol.

Miller illuſtr. ic. Ioh. Miller Illuſtrario Syſtematis ſexualis Linnæi. Londini. 1777. fol.

Miſc. taur. Miſcellanea Philoſophico - mathematica Taurinenſia. Vol. 5. Taurini. 1759. &c. 4to.

Mœnch. meth. Methodus plantas horti botanici & agri marburgenſis a ſtaminum ſitu deſcribendi, auctore Conrado Mœnch. Marburgi Cattorum. 1794. 8vo.

Moris. hiſt. Rob. Moriſon Hiſtoria Plantarum univerſalis Oxonienſis. Oxonii 1680. Pars 3. abſoluta a Iacobo Bobartio. Oxonii. 1699. fol.

Müll. fridr. Ott. Fridr. Müller Flora Fridrichsdaliana. Argentorati. 1767. 8vo.

Neck. meth. N. I. de Necker Methodus muscorum. Manheimii. 1771. 8vo.

Pallas it. P. S. Pallas Reiſe durch verſchiedene Provinzen des ruſſiſchen Reichs. 3 Theile. Petersburg. 1771. &c. 4to.

Pallas roſſic. — Flora Roſſica ſeu Stirpium Imperii roſſici per Europam & Aſiam indigenarum Deſcriptiones & Icones. Partes 2. Petropoli. 1784. fol.

Parkinſon theatr. Theatrum Botanicum: The Theatre of Plants, collected by Iohn Parkinſon. London. 1640. fol.

Plenk ic. Ioſ. Iac. Plenk Icones Plantarum medicinalium. Viennæ. 1788. &c. fol.

Pluk. alm. Leon. Plukenetii Almageſtum Botanicum. Londini. 1696. 4to.

Pluk phyt. — Phytographia. Londini. 1691. 1692. 4to.

Pollich palat. I. A. Polich Hiſtoria Plantarum in Palatinatu Electorali ſponte naſcentium. Tomi 3. Mannheimii. 1776. 8vo.

Regn. bot. La Botanique miſe à la portée de tout le Monde par P. & D. Regnault. Paris. 1774. fol.

Reitter und Abel Abbild. Abbildung der 100 deutſchen wilden Holzarten. Stuttgart. 1790. 4to.

Retzii ſcand. Floræ Scandinaviæ Prodomus, auctore A. I. Retzio. Editio altera. Lipſiæ. 1795. 8vo.

Retzii obs. — Obſervationum botanicarum faſciculi 4. Lipſiæ. 1779. &c. fol.

Riv. mon. A. Q. Rivini Ordo Plantarum, quæ ſunt flore irregulari monopetalo. Lipſiæ. 1691. fol.

Riv. pent. — Ordo Plantarum, quæ ſunt flore irregulari pentapetalo. Lipſiæ. 1699. fol.

Roth. germ. A. G. Rothii Tentamen Floræ germanicæ. Tomi 3. Lipſiæ. 1788. &c. 8vo.

Sauv. monſp. F. B. de Sauvages Flora Monſpelienſis. Hagæ. 1751. 8vo.

Schæff. fung. I. C. Schæffer Fungorum qui in Bavaria & Palatinatu circa Ratisbonam naſcuntur Icones. Tomi 4. 1762. &c. 4to.

Scheuch. agr. I. Scheuchzeri Agroſtographia. Tiguri. 1775. 4to.

Scheuch. it. — Itinera alpina. Vol. 2. Lugduni Batavorum. 1723. 4to.

Schrader ſpic. Spicilegium Floræ Germanicæ. Auctore A Schrader. Pars prior. Hannoveræ. 1794. 8vo.

Schreb. gr. I. C. D. Schrebers Beſchreibung der Gräſer, nebſt ihren Abbildungen. Leipzig. 1769. fol.

Schreb. ſpic. — Spicilegium Floræ Lipſicæ. Lipſiæ. 1771. 8vo.

Scop. ann. I. A. Scopoli Annus 1—5. hiſtorico-naturalis. Lipſiæ. 1769. &c. 8vo.

Scop. carn. ed. 1. — Flora Carniolica. Editio prima. Viennæ. 1760. 8vo.

Scop. carn. ed. 2. — Flora Carniolica. Editio altera. Tomi 2. Viennæ. 1772. 8vo.

Scop. inſubr. — Deliciæ Floræ & Faunæ Inſubricæ. Partes 3. Ticini. 1786. &c. fol.

Segu. ver. Plantæ Veronenſes. Auctore I. F. Seguierio. Vol. 2. Veronæ. 1745. Tomus tertius ſeu Supplementum. 1754. 8vo.

Sibb. ſcot. R. Sibbald Scotia illuſtrata. Edimburgi. 1684.

Sibthorp oxon. Flora oxonienſis. Auctore I. Sibthorp. Oxonii. 1794. 8vo.

Sp. pl. C. Linnæi Species Plantarum. Tomi 2. Editio ſecunda. Holmiæ. 1763. 8vo.

Syſt. veget. — Syſtema Vegetabilium, Editio 14. Curante I. A. Murray. Gottingæ. 1784. 8vo.

Syſt. veget. ed. 13. — Syſtema Vegetabilium. Editio 13. Gottingæ & Gothæ. 1774. 8vo.

Thunb. japon. C. P. Thunberg Flora Iaponica. Lipſiæ. 1784. 8vo.

Till. piſ. M. A. Tilli. Catalogus Plantarum Horti Piſani. Florentiæ. 1723. fol.

Timm meg. Floræ Megapolitanæ Prodromus. Auctore I. C. Timm. Lipſiæ 1788. 8vo.

Tourn. inſt. I. P. Tournefort Inſtitutiones Rei Herbariæ. Tomi 3. Lugduni. 1719. 4to.

Vaill. paris. Botanicum pariſienſe par S. Vaillant. Leide & Amſterdam. 1727. fol.

Vill. proſp. Villars Proſpectus de l' Hiſtoire des Plantes du Dauphiné. Grenoble. 1779. 8vo.

Web. dec. G. H. Weber Præſide Diſſ. Plantarum minus cognitarum Decuria. Kiloniæ. 1784. 4to.

Web. ſpic. D. G. H. Weberi Spicilegium Floræ Gottingenſis. Gothæ. 1778. 8vo.

Weis. crypt. Plantæ cryptogamicæ Floræ Gottingenſis. Collegit D. F. G. Weis. Gottingæ. 1770. 8vo.

Wigg. prim. F. H. Wiggers Primitiæ Floræ Holſatiæ. 1780. 8vo.

Willd. baumz. Berliniſche Baumzucht von K. L. Willdenow. Berlin. 1796. 8vo.

Willd. ber. — Floræ Berolinenſis Prodromis. Berolini 1787. 8vo.

Zann. iſt. Iſtoria delle piante, che naſcono ne' lidi intorno a Venezia, opera poſtuma di Gian-Girolamo Zannichelli accresciuta da Gian Iacopo figliuolo dello ſteſſo. In Venezia. 1735. fol.

MONANDRIA.

MONOGYNIA.

SALICORNIA. Calyx monophyllus, ventricoſus, integer. Corolla nulla. Semen unicum.

herbacea. 1. S. herbacea patula, articulis apice compreſſis emarginato-bifidis. *Sp. pl. p.* 5. *Fl. dan. t.* 303. *Scop. carn. tom.* 1. *p.* 4.

Kali geniculatum ſimplicius, et brevius, rectum, et annuum. *Bauh. pin.* 289. *Zann. iſt. p.* 232. *t.* 33. *n.* 2.

In Litorali auſtriaco in paluſtribus Duini, Aquilejæ. In Pannonia ad lacum Peiſonis, vulgo *Neuſidelſee.* Floret Iulio, Auguſto.

fruticoſa. 2. S. caule erecto fruticoſo. *Sp. pl. p.* 5.

Salicornia articulis ſubæqualibus, obtuſis, internodiis approximatis. *Ger. prov. p.* 328.

Kali geniculatum majus, ſempervirens. *Bauh. pin.* 289. *Zann. iſt. p.* 231. *t.* 32.

In arenoſis ad litus maris adriatici: in Litorali infrequens; frequentiſſima in inſulis venetis. Fl. cum præcedente.

Utrique caules, ramique aphylli, articulati. Flores in articulis.

HIPPURIS. Calyx minimus, bilobus, germen coronans. Corolla nulla. Stamen intra lobum anteriorem. Stylus e poſteriori lobo calycis.

vulgaris. 1. H. foliis octonis ſubulatis. *Syſt. veget. p.* 53. *Curtis lond. Fl. dan. t.* 87. *Scop. carn. tom.* 1. *p.* 5.

Limnopeuce. *Hall. hiſt. n.* 1572.

Equiſetum paluſtre, brevioribus foliis, polyſpermun. *Bauh. pin.* 15.

In aquis vel ſtagnantibus, vel lente fluentibus. Fl. Majo, Iunio. Perennis.

DIGYNIA.

CALLITRICHE. Calyx nullus. Petala duo. Capſula bilocularis, tetraſperma.

aquatica. 1. Callitriche. *Sibthorp oxon.* p. 2.

Callitriche verna, & C. autumnalis. *Sp. pl. p.* 6.

Callitriche fontana, & C. ſtagnalis. *Scop. carn. tom.* 2. *p.* 251.

In rivulis, foſſis, inundatis. Fl. primo vere, quandoque & autumno. Per.

Variat foliis omnibus ovalibus, vel linearibus; ſuperioribus ovalibus, reliquis linearibus; apice integris, vel bifidis; &, aſſerente Cl. Sibthorp, floribus monoicis, ac hermaphroditis.

BLITUM. Calyx trifidus. Corolla nulla. Capſula tenuiſſima, calyce baccato tecta. Semen unicum.

capitatum. 1. B. capitulis ſpicatis terminalibus. *Sp. pl. p.* 6.

Blitum foliis ſemitrilobis, rariter dentatis. *Hall. hiſt. n.* 1571.

Morocarpus capitatus. *Scop. carn. tom.* 1. *p.* 6.

Atriplex ſylveſtris lappulas habens. *Bauh. pin.* 119.

Atriplex ſylveſtris baccifera. *Clus. hiſt. p.* CXXXV.

In Litorali, Forojulio, Tyrolique calidiore. Fl. æſtate.

DIANDRIA.

MONOGYNIA.

LIGUSTRUM. Corolla quadrifida. Bacca bilocularis: loculamentis membrana tenui veſtitis. Semina bina.

vulgare. 1. L. foliis ovatis obtuſis, panicula ſimplici trichotoma. *Thunb. japon. p.* 17. *Sp. pl. p.* 10. *Scop. carn. tom.* 1. *p.* 7. *Hall. hiſt. n.* 530.

Le Troêne. Regn. bot.

In ſylvis, ad ſepes, alibique. Fl. Iunio. Frutex.

PHILLYREA. Corolla quadrifida. Bacca bilocularis. Semina ſolitaria; unum ſæpe abortiens.

media. 1. P. foliis oblongo-lanceolatis integerrimis ſerratisque. *Ait. Kew. tom.* 1. *p.* 11. *Sp. pl. p.* 10.

Phillyrea tertia. *Clus. hiſt. p.* 52.

Phillyrea folio Liguſtri. *Bauh. pin.* 476.

In Litorali in rupeſtribus ad mare. In agro fluminenſi in *Rezzine* quoque ad torrentem. Fl. Martio; Aprili. Fr.

latifolia. 2. P. foliis ovato-oblongis ſubcordatis ſerratis. *Ait. Kew. tom.* 1. *p.* 12. *Sp. pl. p.* 10.

Phillyrea prima; & P. ſecunda. *Clus. hiſt. p.* 51. 52.

Habitat & floret cum præcedente. Fr.

OLEA. Corolla quadrifida: laciniis ſubovatis. Nux ſubbilocularis.

europaea. 1. O. foliis lanceolatis. *Sp. pl. p.* 11. *Scop. carn. tom.* 1. *p.* 8.

Olea ſativa. *Clus. hiſt. p.* 26.

L'Olivier. Regn. bot.

In Litorali in olivetis, vineiſque neglectis fruticis forma creſcit, habet ramos ſpineſcentes, nunquam floret. In cultis arboris magnitudinem attingit, eſt inermis, & Fl. Aprili.

SYRINGA. Corolla quadrifida. Capſula bilocularis.

vulgaris. 1. S. foliis ovato-cordatis integris. *Syſt. veget. p.* 57. *Curtis magaz.* 183. *Haller hiſt. n.* 531.

Syringa cœruleo flore. *Cluſ. hiſt. p.* 56.

Occurrit per Auſtriam reliquaſque provincias ad ſepes. Fl. primo vere. Fr.

CIRCÆA. Calyx diphyllus. Corolla dipetala. Germen inferum. Capſula uncinato-hiſpida, bivalvis. Semina ſolitaria.

lutetiana. 1. C. caule erecto piloſo, foliis ovatis ſubſerratis. *Retzii ſcand. p.* 5. *obs. faſc.* 1. *p.* 9. *Sp. pl. p.* 12. *Fl. dan. t.* 256. *Curtis lond.*

La Circée Regn. bot.

In ſylvis umbroſis, ad ſepes humidas. Fl. Iunio, Iulio. Per.

alpina. 2. C. caule proſtrato glabro, foliis cordatis dentatis. *Retzii ſcand. p.* 5. *obs. faſc.* 1. *p.* 9. *Sp. pl. p.* 12. *Fl. dan. t.* 210.

In ſubalpinis. Fl. Iulio, Auguſto. Per.

Circæa lutetiana habet petala baſi rotundata, ſtylum baſi incraſſatum. Alpinæ contra petala ſunt baſi attenuata, ſtylus ubique æqualis craſſitiei.

VERONICA. Calyx quadri-vel quinque partitus. Corolla limbo quadripartito, lacinia infima anguſtiore. Capſula bilocularis.

* *Spicatæ.*

longifolia. 1. V. ſpicis terminalibus, foliis oppoſitis lanceolatis ſerratis acuminatis. *Sp. pl. p.* 13.

Lyſimachia cœruleo flore. *Cluſ. hiſt. p.* LIII.

In paluſtribus: in Carniolia circa Labacum; in Litorali in agro aquilejenſi. Copioſior occurrit in Auſtria, Pannoniaque in ſimilibus locis, & in ſylvis demiſſis. Fl. Iunio, Iulio. Per.

Radix promit caules plures, alios foliis oppoſitis, alios ternis, quosdam ſimul ternis & oppoſitis veſtitos. Laciniæ calycis in aliis floribus æquales ſunt, in aliis inæquales.

ſpicata. 2. V. ſpica terminali, foliis oppoſitis crenatis obtuſis, caule adſcendente ſimpliciſſimo. *Sp. pl. p.* 14. *Fl. dan. t.* 52.

Veronica foliis ellipticis, ſerratis, conjugatis, floribus ſpicatis. *Hall. hiſt. n.* 522.

In collibus herbidis, montoſis apricis. Fl. Iulio, Auguſto, Septembri. Per.

Pro diverſa ſoli natura, atque expoſitione variat caule ſimplici, ramoſo; ſoliis latioribus, anguſtioribus, acutis, & obtuſis.

hybrida. 3. V. ſpicis terminalibus, foliis oppoſitis obtuſe ſerratis ſcabris, caule erecto. *Sp. pl. p.* 14.

Veronica ſpicata cambro-britanica, bugulæ ſubhirſuto folio. *Raj. angl.* 3. p. 278 *t.* 11. *f.* 1.

In aſperis Carnioliæ calidioris, copioſiſſime in *Lippiza.* Fl. Iulio, Auguſto. Per.

officinalis. 4. V. ſpicis lateralibus pedunculatis, foliis oppoſitis, caule procumbente. *Sp. pl. p.* 14. *Fl. dan. t.* 248.

Veronica caule procumbente foliis ſcabris, petiolatis, ovatis, ex alis racemoſa. *Hall. hiſt. n.* 540.

La Veronique mâle, ou le Thé d' Europe. Regn. bot.

In ſylvis montanis, & ſubalpinis. Fl. Majo, Iunio. Per.

** *Corymboſo - racemoſæ.*

aphylla. 5. V. corymbo terminali, caule nudo. *Sp. pl. p.* 14. *Scop. carn. t.* 1, *p.* 10.

Veronica caule nudo, foliis ovatis ſerratis, racemo paucifloro. *Hall. hiſt. n.* 541.

Veronica parva ſaxatilis, cauliculis nudis. *Segu. ver. tom.* 1. *p.* 241. *t.* 3. *f.* 2.

Teucrium minimum. *Clus. hiſt. p.* 350.

In alpibus. Fl. Iulio, Auguſto. Per.

Tota planta, corolla excepta, obſita eſt pilis articulatis.

bellidioides. 6. V. corymbo terminali, caule adſcendente diphyllo, foliis obtuſis crenatis, calycibus hirſutis. *Syſt. veget. p.* 59.

Veronica caule ſimplici, paucifloro, foliis ſcabris, ſerratis, ovatis, ſpica pauciflora. *Hall. hiſt. n.* 543. *t.* 15.

Veronica alpina, Bellidis folio hirſuto. *Bauh. pin.* 247. *prodr.* 116.

In alpibus Styriæ ſuperioris, Carinthiæ, Tyrolis. Fl. Iulio, Auguſto. Per.

In planta pumila unicum par foliorum oppoſitorum, in majori duo et tria paria numeravi.

fruticuloſa. 7. V. corymbo terminali, foliis lanceolatis integerrimis crenatisque, caulibus fruticuloſis erectis.

Veronica caulibus fruticoſis, baſi procumbentibus; racemo terminali multifloro; foliis elliptico-lanceolatis, ſubdenticulatis; calycibus acutis. *Wulfen* in *Iacqu. collect. tom.* 4. *p.* 229. *t.* 5.

Veronica foliis ovatis, crenatis, fructu ovali; floribus in ſummo caule purpuraſcentibus. *Hall. hiſt. n.* 545.

Veronica frutescens. *Scop. carn. tom.* 1. *p.* 19.

In Carniolia in rupeſtribus ad radices montium in *der Wochein* & *Gariutſche.* Fl. Iunio, Iulio.

ſaxatilis. 8. V. corymbo terminali, foliis obovatis integerrimis crenatisque, caulibus fruticuloſis diffuſis.

Veronica racemo terminali; foliis ovatis crenatis, glabris, pedunculo brevioribus. *Scop. carn. ed.* 1. *p.* 302. *n.* 2.

Veronica corymbo terminali, foliis oppoſitis glabriusculis, pedunculis folia floralia ſuperantibus. *Linn. ſuppl. p.* 83.

Veronica fruticuloſa. *Fl. dan. t.* 342.

Veronica tertia fruticans. *Clus. hiſt. p.* 347.

In alpium rupeſtribus, ſaxoſisque. Fl. æſtate.

V. fruticuloſa habet caules erectos, ſimplices. Saxatilis ramoſos, diffuſosque. Saxatili ſunt folia obovata, plana. Fruticuloſæ oblonga, lanceolata, & ob margines parum conniventes ſubcanaliculata, acuta; utrique vero vel integerrima, vel plus vel minus crenulata. Saxatilis habitat tantum in alpibus, geritque corollas cœruleas. Fruticuloſa in demiſſis, habetque corollas carneas lineis purpureis variegatas.

alpina. 9. V. corymbo terminali, foliis oppoſitis, calycibus hispidis. *Sp. pl. p.* 15. *Fl. dan. t.* 16.

Veronica caule ſimplici, foliis ovatis glabris, ſubſerratis, ſpica pauciflora. *Hall. hiſt. n.* 544. *t.* 15. *f.* 2.

Teucrium ſextum pumilum. *Clus. hiſt. p.* 350.

In alpium herbidis. Fl. æſtate. Per.

ſerpyllifolia. 10. V. racemo terminali ſubſpicato, foliis ovatis glabris crenatis, *Sp. pl. p.* 15. *Curtis lond. Fl. dan. t.* 492. *Scop. carn. tom.* 1. *p.* 12.

Veronica caule recto, foliis ovatis, crenatis, glabris, petiolis ex alis unifloris, breviſſimis. *Hall. hiſt. n.* 546.

Ad margines ſylvarum, ad ſepes, in pratis, & alibi. Fl. Majo. Per.

Folia caulina oppoſita, ovata. Floralia alterna, lanceolata. Pedunculi axillares uniflori.

Becca-bunga. 11. V. racemis lateralibus, foliis ovatis planis, caule repente. *Sp. pl. p.* 16. *Curtis lond. Fl. dan. t.* 511. *Scop. carn. tom.* 1. *p.* 13.

Veronica foliis ovatis, ſerratis, glabris, ex alis racemoſa. *Hall. hiſt. n.* 534.

Le Beccabunga Regn. bot.

In rivulis, ad foſſas, fluvios. Fl. æſtate. Per.

Anagal-lis. 12. V. racemis lateralibus, foliis lanceolatis ſerratis, caule erecto. *Sp. pl. p.* 16. *Curtis lond. Fl. dan. t.* 903. *Scop. carn. tom.* 1. *p.* 13.

Veronica foliis lanceolatis, ſerratis, glabris, ex alis racemoſa. *Hall. hiſt. n.* 533.

Habitat & floret cum præcedente.

ſcutel-lata. 13. V. racemis lateralibus alternis: pedicellis pendulis, foliis linearibus integerrimis. *Sp. pl. p.* 16. *Fl. dan. t.* 209. *Scop. carn. tom.* 1. *p.* 20.

Veronica foliis lineari-lanceolatis, racemis ex alis paucifloris. *Hall. hiſt. n.* 532.

Anagallis aquatica anguſtifolia, ſcutellata. *Bauh. pin.* 252. *prodr.* 119.

In paluſtribus, & inundatis. Fl. æſtate. Per.

Teucri-um. 14. V. racemis lateralibus longiſſimis, foliis ovato-lanceolatis dentatis obtuſiuſculis, caulibus erectis. *Sp. pl. p.* 16.

Teucrii quarti tertia ſpecies. *Clus. hiſt. p.* 349.

In Auſtriæ, Pannoniæ collibus herbidis, pratisque ſiccioribus. Fl. Majo, Iunio. Per.

* *Caules erecti, plerumque pedales. Folia ovato-lanceolata, minime rugoſa, dentata, dentatisque pauca integerrima mixta. Racemi laterales, longiſſimi. Calyces quadripartiti.*

proſtrata. 15. V. racemis lateralibus, foliis oblongo-ovatis ſerratis, caulibus proſtratis. *Sp. pl. p.* 17.

Veronica foliis imis ellipticis, dentatis, ſupremis lanceolatis, ex alis racemoſa. *Hall. hiſt. n.* 538.

Teucrium quintum. *Cluſ. hiſt. p.* 350.

In collibus, pratis, ad vias, alibique. Fl. Majo, Iunio. Per.

Caules ſemper proſtrati, in feliciori ſolo ex geniculis radices agentes. Calyx quadripartitus; lacinia unica quandoque bifida.

montana. 16. V. racemis lateralibus paucifloris, calycibus hirſutis, foliis ovatis rugoſis crenatis petiolatis, caule debili. *Sp. pl. p.* 17. *Iacqu. auſtr.* t. 109.

Veronica caule procumbente, foliis hirſutis, cordatis, retuſis, racemis paucifloris. *Hall. hiſt. n.* 539.

Veronica ſubbiscutata. *Crantz faſc.* 343.

In Auſtriæ, Styriæ ſylvis umbroſis montanis, & ſubalpinis. Fl. Iunio, Iulio. Per.

Chamædrys. 17. V. racemis lateralibus, foliis ovatis ſeſſilibus rugoſis dentatis, caule bifariam piloſo. *Syſt. veget. p.* 60. *Fl. dan. t.* 448. *Curtis lond.*

Veronica foliis cordatis, ſubrotundis, hirſutis, nervoſis, ex alis racemoſa. *Hall. hiſt. n.* 536.

Chamædrys quorundam ſylveſtris. *Cluſ. hiſt.* p. 352.

In ſylvis, pratis, ad ſepes. Fl. Majo, Iunio. Per.

Planta in umbroſis humidis habet folia ſubglabra; in apricis ſiccis tota pilis caneſcit.

auſtriaca. 18. V. racemis lateralibus, foliis lineari-lanceolatis pinnato-dentatis. *Sp.* pl. *p.* 17. *Iacqu. auſtr. t.* 329.

In Carniolia calidiore, Litorali, Croatiaque finitima in collibus herbidis, pratisque montanis. Fl. Majo, Iunio. Per.

In aprico asperoque ſolo radix emittit caules paucos, ſemipedales, ſæpe breviores, foliis veſtitos anguſtioribus, vix pinnatifidis; pinnis raro dentatis. In tali ſtatu eſt V. Teucrio valde ſimilis. In pratis contra inter elatas ſtirpes ex unica radice exeunt caules plures, pedales, al-

tioreſque, foliis prædíti amplioribus, pinnatifidis; pinnis dentatis. Calyx eſt ſemper hiſpidus. Differt a V. multifida, quæ eſt planta ſibirica: nam huic, docente Cl. Smith Linnæani herbarii poſſeſſore, ſunt folia multipartita: laciniis pinnatifidis; lobis decurrentibus. Calyces glabri. Vid. Linn. Trans. tom. I. on Veronica.

urticæfolia. 19. V. racemis lateralibus, foliis ovato-lanceolatis hirtis argute ſerratis, caule erecto. *Syſt. veget.* p. 60. *Iacqu. auſtr. t.* 59.

In Auſtria occurrit prope *Schottwien* in monte *Sömmering*, extenditur exinde per Styriam, reliquasque provincias, ubi in montanis & ſubalpinis creſcit. Fl. æſtate. Per.

latifolia. 20. V. racemis lateralibus, foliis cordatis rugoſis dentatis, caule ſtricto. *Sp. pl.* p. 18.

Veronica foliis cordatis obtuſis, nervoſis, ſerratis, racemis denſiſſimis. *Hall. hiſt. n.* 537.

Veronica Pseudo-Chamædrys. *Iacqu. auſtr. t.* 60.

Teucrium quartum. *Clus. hiſt. p.* 349.

In Auſtria, Pannonia in fruticoſis, ad ſepes, ſylvarum margines, in pratis ſiccioribus. Fl. Iunio, Iulio. Per.

*** *Pedunculis unifloris.*

agreſtis. 21. V, floribus ſolitariis, foliis cordatis inciſis pedunculo brevioribus. *Sp. pl. p.* 18. *Curtis lond. Fl. dan. t.* 449. *Scop. carn. tom.* I. *p.* 21.

Veronica caule procumbente, foliis petiolatis, ovatis, crenatis. *Hall. hiſt. n.* 549.

In cultis, ruderatis. Fl. a fine hyemis ad ſerum autumnum.

arvenſis. 22. V. floribus ſolitariis, foliis cordatis inciſis pedunculo longioribus. *Sp. pl. p.* 18. *Curtis lond.*

Veronica caule erecto, foliis ovatis, ſubhirſutis, dentatis, petiolis breviſſimis. *Hall. hiſt. n.* 542.

In arvis, alibique. Fl. primo vere.

hederæ-folia. 23. V. caule procumbente, foliis lobatis, petiolatis, pedunculis unifloris. *Hall. hiſt. n.* 550. *Sp. pl. p.* 19. *Curtis lond.*

In cultis, ad vias, in ruderatis. Fl. primo vere.

Folia pro diverſa ſoli natura, atque expoſitione ſunt plus vel minus hirſuta. Loborum numerus varius.

triphyllos. 24. V. floribus ſolitariis, foliis digitato-partitis, pedunculis calyce longioribus. *Syſt. veget. p.* 60. *Fl. dan. t.* 627. *Scop. carn. tom.* 1. *p.* 21.

Veronica foliis ovatis, tripartitis, & quinque partitis, petiolis unifloris. *Hall. hiſt. n.* 551.

In arvis, hortis, alibique. Fl. primo vere.

verna. 25. V. floribus ſolitariis, foliis digitato-partitis, pedunculis calyce brevioribus. *Syſt. veget. p.* 60. *Fl. dan. t.* 252.

Veronica foliis ovatis, tripartitis & quinquepartitis, petiolis unifloris, breviſſimis. *Hall. hiſt. n.* 552.

Habitat & floret cum præcedente.

Differt a V. triphylla pedunculis breviſſimis, minimo calyce, & corolla. Haller l. c.

acinifolia. 26. V. floribus ſolitariis pedunculatis, foliis ovatis glabris crenatis, caule erecto ſubpiloſo. *Sp. pl. p.* 19.

Veronica minima, clinopodii minoris folio glabro, romana. *Bocc. mus. p.* 29. *t.* 102. *Vaill. paris. p.* 201. *t.* 23. *f.* 3.

Veronica romana. *Allion. pedem. tom.* 1. *p.* 79. *t.* 85. *f.* 2. *Scop. carn. tom.* 1. *p.* 18.

In apricis herbidis, ruderatis, & cultis. Fl. Majo.

Veronica romana exulare debet. Omnia ejus ſynonima in Speciebus plantarum ed. 1. perti-

nent ad V. acinifoliam. Specimen in herbario Linnaeano proſtans, ad quod differentia ſpecifica, & deſcriptio mant. p. 317, facta eſt, non eſt aliud quam V. peregrina. Smith in Linn. Trans. tom. 1. on Veronica.

PÆDEROTA. Calyx quinquepartitus. Corolla bilabiata, ecalcarata. Capſula bilocularis.

cœrulea. 1. P. foliis oppoſitis ſerratis. *Scop. ann.* 2. *p.* 41. *Syſt. veget. p.* 61.

Pæderota Bonarota. *Iacqu. auſtr. app. t.* 39. *Syſt. veget. p.* 61.

Veronica petræa ſempervirens. *Pon. in Cluſ. hiſt. p.* CCCXXXVI.

Bonarota montana, italica, Chamædryos folio, flore cœruleo. *Mich. gen. p.* 19. *t.* 15. *f.* 1.

Bonarota montana, italica, Chamædryos folio rotundiore, & quaſi circinato, ſpica cœrulea habitiore. *Mich. l. c. t.* 19. *f.* 2.

In Tyroli circa Leontium. In Carinthia in ſummitatibus apricis prope Pontebam *Trog*, aut Trigel dictis, nec non Malborghetti, ubique in rupeſtribus. *Wulfen.* Fl. Iunio, Iulio. Per.

Folia ſunt omnia oppoſita, ſeſſilia, ſupra glabra, ſubtus villoſa: inferiora breviora; ſuperiora oblonga, ſumma lineari-lanceolata. Flores ſpicati. Bracteæ lineares. Calyx quinquepartitus: laciniis linearibus, rufis, villoſis. Corolla violacea, raro alba. Labium ſuperius tridentatum, inferius plerumque integrum. Iacqu. l. c.

lutea. 2. P. foliis oppoſitis alternis terniſque ſerratis. *Scop. ann.* 2. *p.* 41. *Syſt. veget. p.* 61.

Pæderota Bonarota. *Iacqu. hort. tom.* 2. *p.* 55. *tab.* 121.

Pæderota Ageria. *Syſt. veget. p.* 61:

In rupeſtribus Carnioliæ in alpibus wochinenſibus, tolmienſibus, in agro idrienſi. In Carinthia in monte Loibl. In Forojulio circa

Goritiam ad latera Liſoncii; alibique. Fl. Iunio, Iulio. Per.

Caulis ut in præcedente vel ſimplex, vel baſi ramoſus. Folia, in aliis caulibus inferiora oppoſita, quæ medium caulem occupant alterna, ſumma terna; dum in aliis ſumma ſunt alterna, quæ medium caulem veſtiunt, oppoſita. infima terna. Flores ſpicati. Corollæ tubus baſi ſubgloboſo-gibbus. Limbus bilabiatus: labium ſuperius in quibusdam trifidum, in aliquibus bifidum, & modo emarginatum, modo integrum in aliis. Faux barbata.

WULFENIA. Corolla ringens, ecalcarata, ſemihians, ad faucem barbata.

carinthiaca. 1. Wulfenia. *Syſt. veget. p. 65. Iacqu. miscell. tom. 2. p. 60. t. 8. f. 1. ic. t. 2.*

In Carinthiæ alpe Kibeggenſi (*Kühewegeralpe*) vallis Gilienſis (*im Gailthale*) prope Fanum ſancti Hermagoræ (vulgo *Ermachor*, aut *Michor*) paulo ſupra ſuprema paſtorum mapalia. *Wulfen.* Fl. Iulio. Per.

GRATIOLA. Calyx heptaphyllus: foliolis duobus exterioribus patulis. Corolla irregularis, reſupinata. Stamina quatuor, quorum duo ſterilia. Capſula bilocularis.

officinalis. 1. G. foliis lanceolatis ſerratis, floribus pedunculatis. *Sp. pl. p. 24. Fl. dan. t. 363. Scop. carn. tom. 1. p. 24. Hall. hiſt. n. 329.*

La Gratiole. Regn. bot.

In humidis. Fl. æſtate. Per.

Folia ſeſſilia, ſemiamplexicaulia, ſerrata, inferne integerrima. Pedunculi axillares, ſolitarii, uniflori. Bracteæ duæ ad baſin cujuslibet calycis. Gouan. hort. p. 13.

PINGUICULA. Calyx bilabiatus. Labium ſuperius trifidum; inferius bifidum. Corolla ringens, calcarata. Capſula unilocularis.

vulgaris. 1. P. nectario cylindraceo longitudine petali. *Sp. pl. p.* 25. *Fl. dan. t.* 93.

Pinguicula calcare floris longitudine. *Hall. hiſt. n.* 292.

In Auſtriæ pratis paluſtribus, circa *Erla, Lachſenburg, Himberg, Schwadorf an der Fiſcha, Gallbrunn.* Fl. Majo, Iunio. Per.

alpina. 2. P. nectario conico petalo breviore. *Sp. pl. p.* 25. *Fl. lapp. n.* 12. *t.* 12. *f.* 3. *Fl. dan. t.* 453. *Scop. carn. tom.* 1. *p.* 25.

Pinguicula calcare breviſſimo. *Hall. hiſt. n.* 293.

In alpinis & ſubalpinis. Fl. a Majo ad Iulium. Per.

UTRICULARIA. Calyx diphyllus, æqualis. Corolla ringens, calcarata. Capſula unilocularis.

vulgaris. 1. U. nectario conico, ſcapo paucifloro. *Sp. pl. p.* 26. *Fl. dan. t.* 138. *Hall. hiſt. n.* 290.

In Auſtria in aquis ſtagnantibus circa *Himberg, Bruck an der Leitha, Rohrau,* & in inſulis danubialibus. Fl. æſtate Per.

Folia omnia ſunt ſubmerſa, multifido - capillaria, veſiculas aqua repletas geſtantia.

LYCOPUS. Corolla quadrifida: lacinia unica emarginata. Stamina diſtantia. Semina quatuor, retuſa.

europæus. 1. L. foliis ſinuato - ſerratis. *Sp. pl. p.* 30. *Curtis lond.*

Lycopus foliis acute ſerratis, appendiculatis. *Hall. hiſt. n.* 220.

Pſeudo - marrubiaſtrum paluſtre. *Riv. mon. t.* 21.

Ad rivulos, fluvios, foſſas; & in aliis humidis locis. Fl. æſtate. Per.

ROSMARINUS. **Corolla inæqualis, labio ſuperiore bipartito, erecto; labio inferiore reflexo, trifido. Filamenta ſimplicia cum dente.**

officinalis. 1. Rosmarinus. *Sp. pl. p.* 33. *Scop. carn. tom.* 1. *p.* 26. *Hall. hiſt. n.* 250.

Le Romarin. Regn. bot.

In Litorali locis rupeſtribus ad mare inter Flumen & Portum Regium. Fl. Aprili. Fr.

SALVIA. **Corolla inæqualis. Filamenta transverſe pedicello affixa.**

officinalis. 1. S. foliis lanceolato-ovatis integris crenulatis, floribus ſpicatis, calycibus acutis. *Sp. pl. p.* 34. *Scop. carn. tom.* 1. *p.* 27.

La petite Sauge. Regn. bot.

In ſaxoſis Forijulii, Litoralis: copioſiſſime in agro fluminenſi. Fl. Majo, Iunio. Per.

Folia lato-lanceolata, crenulata, aliqua tantum cordata, omnia denſo villo, uti & caulis, obtecta. Flores verticillati. Calyx bilabiatus: labio ſuperiore bi- vel tridentato; labio inferiore bifido. Corollæ galea emarginata. Labium inferius reflexum, trifidum, lacinia intermedia emarginata.

Flumine præcipue primo vere folia cedunt in pabulum ovium; hinc fit, quod lac ovinum imbutum ſit gratiſſimo ſalviæ odore. Apes amant flores, nec non pueri, qui liquorem melleum in corollæ tubo latitantem ſugunt. Vetulas novi, quae atrophicis, pro phtiſicis habitis, in quibus ob ventriculi languorem digeſtio non rite perficiebatur, jejuno adhuc ſtomacho, foliorum juniorum maſticationem, atque maſticatione expreſſi ſucci, deglutitionem cum ſuperpoto vasculo limpidiſſimæ aquae optimo cum effectu ſuadebant.

ſylveſtris. 2. S. foliis cordato-lanceolatis ſerratis, bracteis coloratis flore brevioribus. *Sp. pl. p.* 34. *Iacqu. auſtr. t.* 212. *Crantz faſc.* 242.

Hormini ſylveſtris quinti ſpecies altera. *Cluſ. hiſt. p.* XXXI.

In Auſtria ad agrorum margines, vias, ſepes, & alibi. Fl. æſtate. Per.

Caules ramoſi. Folia ſubſeſſilia, ſubcordata, obtuſa, inæqualiter et obtuſe ſerrata. Flores ſpicati. Bracteæ ex rotundo acuminatæ, longitudine calycum, plus vel minus purpurascentes. Calyx bilabiatus. Labium ſuperius obſolete tridentatum; inferius bifidum. Laciniæ labii inferioris corollæ ſunt concavæ; laterales oblongæ, intermedia ſubrotunda, emarginata. Iacqu. l. c.

Variat foliis planis, & rugoſiusculis; floribus cæruleis, purpureis & albis; corollæ labio inferiore patente, reflexoque. Salvia nemoroſa, quam Linnæus in Auſtria crescere dicit, eſt S. ſylveſtris foliis planis, corollæ labio infimo reflexo.

auſtriaca. 3. S. foliis ovatis cordatisque eroſo-ſinuatis; radicalibus petiolatis, caule ſubaphyllo, ſtaminibus corolla duplo longioribus. *Ait.* Kew. *tom.* 1. *p.* 41. *Iacqu. auſtr. t.* 112. *Syſt. veget. p.* 69. *Crantz faſc.* 236.

Hormini ſylveſtris quarti altera ſpecies. *Cluſ. hiſt. p.* XXX.

In pratis Auſtriæ: infrequens circa Vindobonam; copioſior circa *Swächat*, *Bruck an der Leitha*, *Rohrau*, & in finitima Pannonia. Fl. Majo, Iunio. Per.

Folia radicalia petiolata, terræ appreſſa. Caulina ſeſſilia. Bracteæ cordatæ, integræ. Calycis labium ſuperius emarginatum, inferius bifidum. Corollæ pallide ochroleucæ galea compreſſa, emarginata. Labii inferioris lacinia intermedia ſubrotunda, atque emarginata porrigitur; laterales oblongæ, obtuſæ, duplo breviores eriguntur. Iacqu. l. c.

pratenſis. 4. S. foliis cordato - oblongis : ſummis amplexicaulibus, verticillis ſubnudis, corollis galea glutinoſis. *Sp. pl. p.* 35. *Scop. carn. tom.* 1. *p.* 30.

Salvia foliis imis cordatis, ſuperioribus longe lanceolatis, verticillis nudis, in ſpicam continuatis. *Hall. hiſt. n.* 253.

Hormini ſylveſtris quarti prima ſpecies. *Clus. hiſt. p.* XXX.

In pratis, & aliis herbidis locis. Fl. Iunio, Iulio. Per.

Folia ima longe, ſuperiora breviter petiolata, ſumma ſeſſilia. Bracteæ acuminatæ. Calycis labium ſuperius tridenticulatum, inferius bifidum. Corollæ galea compreſſa, emarginata. Labium inferius trifidum; laciniis duabus lateralibus reflexis, intermedia majore emarginata, concava.

verticillata. 5. S. foliis cordatis crenato - dentatis, verticillis ſubnudis, ſtylo corollæ labio inferiori incumbente. *Syſt. veget. p.* 70. *Scop. carn. tom.* 1. *p.* 28.

Salvia foliis cordatis, ſerratis, verticillis denſiſſimis, in ſpicam continuatis. *Hall. hiſt. n.* 251.

Horminum ſylveſtre tertium. *Clus. hiſt. p.* XXIX.

Ad vias, ſepes, alibique in locis cultis & ſylvaticis. Fl. æſtate. Per.

Radix ſtolonibus repens. Folia petiolata, quam plurima cordata, inferiora plerumque alata: omnia crenata; crenis apice tuberculo brevi notatis. Flores ſecundi. Bracteæ jam ante florum expanſionem marcescentes. Calycis labium utrumque tridentatum; dentibus labii inferioris longioribus. Corollæ galea compreſſa, pubescens. Labium inferius trifidum; laciniis lateralibus minimis, truncatis, reflexis, intermedia maxima obcordata, concava.

glutinoſa. 6. S. foliis cordato - ſagittatis ſerratis acutis. *Sp. pl. p.* 37. *Scop. carn. tom.* 1. *p.* 28. *Hall. hiſt. n.* 252.

Horminum ſylveſtre ſecundum, Colus Iovis. *Clus. hiſt. p.* XXVIII.

In ſylvis umbroſis, ad collium montiumque radices. Fl. æſtate. Per.

Folia ſagittata. Bracteæ ovato-lanceolata Flores, ut caulis & rami, lento viſcidoque humore manant. Calycis labium ſuperius integrum, inferius bifidum. Lacinia intermedia labii inferioris corollæ eſt maxima, denticulata.

Sclarea. 7. S. foliis rugoſis cordatis oblongis villoſis ſerratis, bracteis floralibus calyce longioribus concavis acuminatis. *Sp. pl. p.* 38.

Horminum ſylveſtre primum. *Clus. hiſt. p.* XXVIII.

Horminum Sclarea dictum. *Bauh. pin.* 238.

L'Orvale. Regn. bot.

In Litorali circa *Moschienizze*, præſertim ad latera viæ, quæ ab Eccleſia ad litus maris ducit. Fl. Iunio, Iulio.

Caulis in apricis ſimplex, in ceſpitoſis, & ad ſepes ramoſus. Folia ima petiolata, ſuperiora ſeſſilia. Calycis labium ſuperius emarginatum, inferius bidentatum; dentibus mucronatis. Corolla guture inflata. Galea compreſſa, obcordata. Labium inferius trifidum; laciniis duabus lateralibus reflexis, intermedia emarginata concava.

Aethiopis. 8. S. foliis oblongis eroſis verticillisque lanatis, corollæ labio crenato, bracteis recurvatis ſubſpinoſis. *Syſt. veget. p.* 71. *Iacqu. auſtr. t.* 211. *Crantz faſc.* 238.

In Auſtria ad ſcrobes, margines agrorum, in agro vindobonenſi, & neoſtadienſi, viciniſque locis. Fl. Iunio, Iulio.

Tota planta lanata. Folia ovata, margine ſinuato-crenulata: radicalia petiolata; reliqua ſeſſilia. Bracteæ ſubrotundæ, concavæ, mucronatæ. Calycis labium ſuperius eſt tri-, inferius bidentatum; dentibus mucronatis. Corollæ galea carinata. Laciniæ labii inferioris laterales oblongæ; media cucullata, ac poſtice fiſſa. Iacqu. l. c.

DIGYNIA.

ANTHOXANTHUM. Calyx gluma bivalvis uniflora Corolla gluma bivalvis, longitudine valvulæ majoris calycinæ: valvula utraque ariſtam e dorſo emittente.

odoratum. 1. A. ſpica oblonga ovata, floſculis ſubpedunculatis ariſta longioribus. *Sp. pl. p.* 40. *Curtis lond. Fl. dan. t.* 666. *Leers herb. p.* 7. *t.* 2. *f.* 1. *Schreb. gr. t.* 5.

Avena diantha, foliculo villoſo, calycis glumis inæqualibus, altera de imo dorſo, altera de ſummo ariſtata. *Hall. hiſt. n.* 1491.

Gramen pratenſe ſpica flaveſcente. *Bauh. pin.* 3. *Scheuch. agr.* 88.

In pratis. Fl. primo vere. Per.

CRYPSIS. Calyx gluma bivalvis, uniflora. Corolla gluma bivalvis, mutica.

aculeata. 1. Crypſis. *Ait. Kew. tom.* 3. *p.* 505.

Schœnus aculeatus. *Sp. pl. p.* 63.

Anthoxanthum aculeatum. *Syſt. veget. p.* 73.

Phleum ſchœnoides. *Iacqu. auſtr. app. t.* 7.

Agroſtis aculeata. *Scop. carn. tom.* 1. *p.* 62.

Gramen ſupinum aculeatum. *Bauh. hiſt.* 2. *p.* 461. *Zann. iſt. p.* 121. *t.* 69.

In Litorali in paluſtribus aquilejenſibus non procul a flumine *Anfora*. In Pannonia ad litus lacus Peiſonis. Fl. Iulio, Auguſto.

TRIANDRIA.

MONOGYNIA.

VALERIANA. Calyx nullus. Corolla monopetala, baſi hinc gibba, ſupera. Semina ſolitaria.

rubra. 1. V. floribus monandris caudatis, foliis lanceolatis integerrimis. *Sp. pl. p.* 44.

Valeriana foliis glaberrimis, floribus calcaratis. *Hall. hiſt. n.* 213.

Valeriana marina latifolia major rubra. *Moris. hiſt.* 3. *p.* 102. *ſect.* 7. *t.* 14. *f.* 15.

In Croatia prope *Fucine. I. Hoſt.* Fl. æſtate. Per.

dioica. 2. V. floribus triandris dioicis, foliis radicalibus ſubintegris, caulinis pinnatis. *Wild. ber. p.* 18. *Sp. pl. p.* 44. *Scop. carn. tom.* 1. *p.* 33. *Curtis lond. Fl. dan. t.* 687.

Valeriana foliis radicalibus petiolatis, ovatis, caulinis pinnatis, ſexubus diſtinctis. *Hall. hiſt. n.* 208.

Valeriana ſylveſtris ſecunda minor. *Clus. hiſt. p. LV.*

In pratis udis. Fl. Majo, Iunio. Per.

officinalis. 3. V. floribus triandris, foliis omnibus pinnatis. *Sp. pl. p.* 45. *Fl. dan. t.* 570. *Scop. carn. tom.* 1. *p.* 32.

Valeriana foliis pinnatis, pinnis dentatis. *Hall. hiſt. n.* 210.

Valeriana ſylveſtris prima. *Clus. hiſt. p. LV.*

In ſylvis, ad ſepes, fluvios, in montium declivibus. Fl. Iunio, Iulio. Per.

tripteris. 4. V. floribus triandris, foliis dentatis; caulinis ternatis ovato-oblongis. *Sp. pl. p.* 45. *Iacqu. auſtr. t.* 268. *Scop. carn. tom.* 1. *p.* 34.

Valeriana foliis radicalibus petiolatis, cordatis, ſerratis, caulinis tripteris. *Hall. hiſt. n.* 211.

In ſubalpinis, & alpinis. Fl. Iunio, Iulio. Per.

Folia radicalia ſimplicia, cordata. Caulina ternata; foliolis ſæpe baſi confluentibus, modo ſerratis, modo integerrimis, rariſſime pinnata. Summa ſimpliciſſima. Flores paniculati. Iacqu. l. c.

montana. 5. V. floribus triandris, foliis ovato-oblongis ſubdentatis, caule ſimplici. *Sp. pl. p.* 45. *Iacqu. auſtr. t.* 269.

Valeriana foliis ovato-lanceolatis, obiter dentatis. *Hall. hiſt. n.* 212.

Valeriana ſylveſtris alpina prima latifolia. *Cluſ. hiſt. p. LV.*

Cum præcedente. Fl. a Iunio ad Auguſtum. Per.

celtica. 6. V. floribus triandris, foliis ovato-oblongis obtuſis integerrimis. *Sp. pl. p.* 46. *Iacqu. collect. tom.* 1. *p.* 24. *t.* 1. *Scop. carn. tom.* 1. *p.* 36.

Valeriana foliis integerrimis, radicalibus ovatis, caulinis linearibus obtuſis. *Hall. hiſt. n.* 209.

Nardus alpina. *Cluſ. hiſt. p. LVII.*

In Styria ſuperiore prope *Eiſenärzt* in monte *Reicheſtein* dicto; in alpibus ſeckavienſibus, waſſerbergenſibus, judenburgenſibus. Invenit quoque *Cl. Wulfen* in Carinthia in alpibus reichenavienſibus ad S. Laurentium, in *Winterthal, Eiſenhut, Hadneralpen, Leitenſteig, Stangelalpen, Wolkart;* in leontinis tyrolenſibus. Fl. æſtate. Per.

ſaxatilis. 7. V. floribus triandris, foliis ſubdentatis: radicalibus ovatis: caulinis lineari-lanceolatis. *Sp. pl. p.* 46. *Iacqu. auſtr. t.* 267. *Scop. carn. tom.* 1. *p.* 35.

Valeriana ſylveſtris alpina ſecunda ſaxatilis. *Cluſ. hiſt. p. LVI.*

In ſiſſuris rupium alpinarum. Fl. Iulio, Auguſto. Per.

Folia radicalia ſunt ſubovata aut lanceolata, aut obverſe ovata, modo integerrima, modo parum inciſa. Caulina oppoſita, lineari lanceolata, integerrima, quandoque ſubdentata. Flores dioici. Maribus eſt corolla multo major cum tubo infundibuliformi. Stamina tria. Germen ſtylo carens. Feminis corolla rotata, tubus breviſſimus. Germen donatur ſtigmate tripartito. Iacqu. l. c.

elongata. 8. V. floribus triandris, foliis radicalibus ovatis; caulinis cordatis ſeſſilibus inciſo ſubhaſtatis. *Sp. pl. app. p.* 1664. *Iacqu. auſtr. t.* 219. *en. p.* 205. *t.* 1. *Scop. carn. tom.* 1. *p.* 35.

In ſummis alpium jugis, locis rupeſtribus, humidiuſculis. Fl. Iulio, Auguſto. Per.

olitoria. 9. V. foliis oblongis rariter inciſis, feminis mucrone ſimplici. *Hall. hiſt. n.* 214.

Valeriana caule dichotomo, foliis lanceolatis integris, fructu ſimplici. *Linn. cliff. p.* 16.

Valeriana Locuſta olitoria. *Sp. pl. p.* 47. *Curtis lond.*

La Mache, ou Blanchette. Regn. bot.

In agris, arvis, & alibi. Fl. Aprili, Majo.

coronata. 10. V. caule dichotomo, foliis lanceolatis dentatis, fructu ſexdentato. *Linn. cliff. p.* 16.

Valeriana floribus triandris, caule dichotomo, pedunculis uniformibus, ſeminibus ſtellatis. *Ger. prov. p.* 219.

Valeriana Locuſta coronata. *Sp. pl. p.* 48.

Valerianella ſemine ſtellato. *Tourn. inſt.* 133.

Valerianella ſemine ſcabioſæ ſtellato hirſuto. *Col. ecphr. p.* 207. *t.* 209.

In Litorali. Fl. Aprili, Majo.

dentata. 11. V. foliis oblongis ſerratis, ſemine tridentato. *Hall. hiſt. n.* 215.

Valeriana Locuſta dentata. *Sp. pl. p.* 48.

Valeriana Locuſta. *Scop. carn. tom.* 1. *p.* 36.

In Carnioliæ, Litoralis, Auſtriæ cultis & ſylvaticis. Fl. Majo.

echinata. 12. V. floribus triandris regularibus, foliis dentatis, fructu lineari tridentato: extimo majore recurvato. *Sp. pl. p.* 47.

Valeriana floribus triandris, caule dichotomo, pedunculis conicis. *Ger. prov. p.* 218.

Valerianella cornucopioides echinata. *Col. ecphr. p.* 204. *t.* 206.

In Carniolia in monte *Nanas*; in Carniæ locis ſylvaticis. *I. Hoſt.* Fl. Iunio.

ſupina. 13. V. floribus tetrandris, involucellis hexaphyllis trifloris, foliis integris. *Syſt. veget. p.* 81. *Wulfen in Iacqu. miſcell. tom.* 2. *p.* 114. *t.* 17. *f.* 2.

Arduin. animadv. 2. *p.* 13. *t.* 3.

In humidis ſummarum Tyrolim inter & gilienſem Carinthiæ vallem alpium locis, *in dem innerſten Laſerz gegen die Lugau zu;* tum ubi ex ſumma alpe Kirſchbaumenſi ultra adſcenditur, per horam parumper verſus orientem, ad ipſas nives; denique *in der öden Au bei Lientz. Wulfen.* Ego quoque in Tyroli in humidis celſiſſimi montis *Donnersberg* prope *Sterzing.* Fl. Iulio, Auguſto. Per.

POLYCNEMUM. Calyx triphyllus. Petala quinque, calyciformia. Stamina duo, ſeu tria. Semen unicum, ſubnudum.

arvenſe. 1. Polycnemum. *Sp. pl. p.* 50. *Iacqu. auſtr. t.* 365. *Hall. hiſt. n.* 1570.

In arvis, ſabuloſisque Auſtriæ, Pannoniæ locis. Fl. æſtate.

CROCUS. Spatha duplicata, exterior vaginans, interior unilateralis. Corolla tubuloſa: limbo ſexpartito æquali. Stamina tria æqualia. Stylus unicus ſemitrifidus. Stigmata tria convoluta criſtata. Capſula trivalvis. *Mœnch - meth. p.* 498.

vernus. 1. C. foliis linearibus carinatis: margine patulo. *Mœnch. l. c. Iacqu. auſtr. app. t.* 36.

Crocus ſativus vernus. *Sp. pl. p.* 50.

In herbidis montanis, & ſubalpinis Litoralis, Forijulii, Carnioliæ, Carinthiæ. Fl. primo vere. Per.

IXIA. Spatha bivalvis. Corolla monopetala, ſupera, ſexpartita. Tubus infundibuliformis, brevis. Limbi laciniæ lanceolatæ, planæ; tribus exterioribus anguſtioribus. Germen ſubrotundum, triquetrum. Stylus filiformis. Stigma trifidum; laciniis bifidis, patentiſſimis, ſupra villoſulis. Capſula trilocularis, trivalvis. *Iacqu. collect. tom.* 3. *p.* 265. *ic. t.* 271.

Bulbocodium. 1. I. ſcapo unifloro breviſſimo, foliis angulatis caulinis, ſtigmatibus trifidis, laciniis bifidis. *Syſt. veget. ed.* 13. *p.* 76. *Ait. Kew. tom.* 1. *p.* 56.

Crocus vernus anguſtifolius quartus. *Clus. hiſt. p.* 208.

Siſyrinchium asprenſinm, anguſto folio, alterum. *Col. ecphr.* 2. *p.* 5. *t.* 7. *f.* 1.

In pratis ſiccis, aliisque herbidis locis agri fluminenſis, *in Belvedere, Braida, San Nicolò, Bergudi*, inter Terſactum & Eccleſiam S. Laurentii, alibique. Fl. Februario, Martio. Per.

Bulbus ſubrotundus, bulbillo emarcido ſuperimpoſitus. Folia linearia, ſic antice margini-

bus ſuis per totam longitudinem conniventia, ut folia mentiantur compreſſula & multiſulcata. Caulis uniflorus. Spatha terminalis, bivalvis: foliolis anguſte lanceolatis, acutis, concavis, oppoſitis; altero erecto, viridi, & fere longitudine corollæ, altero paulo breviori, dorſo virente, cœterum membranaceo, patulo, unde flos evadit veluti lateralis. Iacqu. l. c.

GLADIOLUS. Corolla ſexpartita, irregularis, inæqualis. Stigmata tria.

communis. 1. G. foliis enſiformibus, floribus diſtantibus. *Sp. pl. p.* 52. *Curtis magaz. p.* 86. *Scop. carn. tom.* 1. *p.* 38. *Hall. hiſt. n.* 1261.

Gladiolus floribus uno verſu diſpoſitis. *Bauh. pin.* 43. *Zann. iſt. p.* 113. *t.* 53.

In Carniolia circa lacum zirchnicenſem, & Vipacnm. *Scopoli.* Ego in agro fluminenſi in herbidis, vineisque *a San Nicoló, ſotto i Piopi*, & in toto illo collium tractu, qui inter *Bergudi* ac Flanonam (*Fianona*) ieterjacent. Fl. Majo, Iunio. Per.

IRIS. Corolla ſexpartita: laciniis tribus exterioribus reflexis; interioribus erectis. Stigmata petaliformia, cucullato-bilabiata.

pumila. 1. I. corollis barbatis, caule unifloro foliis breviore. *Sp. pl. p.* 56. *Iacqu. auſtr. t.* 1.

Chamæiris latifolia minor. *Clus. hiſt. p.* 225.

In collibus apricis herbidis: in Auſtria prope *Bertholdsdorf, Medling, Baaden, Haimburg*; in Pannonia inter *Bruck an der Leitha* & *Bahrendorf*, alibique. Fl. Aprili, Majo. Per.

variegata. 2. I. corollis barbatis, caule ſubfolioſo multifloro longitudine foliorum. *Sp. pl. p.* 56. *Iacqu. auſtr. t.* 5.

Iris major latifolia decima quinta. *Clus. hiſt. p.* 221.

Iris latifolia pannonica, colore multiplici. *Bauh.* pin. 31.

In Auſtria in *Leithaberg* ſupra *Männersdorf;* in collibus herbidis non procul ab *Enzersdorf im Langenthale.* In Pannonia inter *Bruck an der Leitha* & *Bahrendorf*, locis herbidis ſylvaticis, alibique. Fl. Majo, Iunio. Per.

germanica. 3. I. corollis barbatis, caule foliis longiore multifloro, floribus inferioribus pedunculatis. *Sp. pl. p.* 55. *Scop. carn. tom.* 1. *p.* 39.

Iris foliis enſiformibus diſtichis, petalis reflexis, barbatis, tubalium arcu obtuſo. *Hall. hiſt. n.* 1258.

Iris latifolia major vulgaris obſoleta. *Clus. hiſt. p.* 224.

Iris, ou la Flambe. Regn. bot.

In udis Auſtriæ inferioris, Pannoniæ. In muris hortorum vetuſtis Carnioliæ, Forijulii, Litoralis. Fl. Majo. Per.

ſambucina. 4. I. corollis barbatis, caule foliis altiore multifloro, laciniis deflexis planis: erectis emarginatis. *Sp. pl. p.* 55. *Iacqu. hort. tom.* 1. *p.* 1. *t.* 2.

Iris major latifolia octava. *Clus. hiſt. p.* 219.

Iris latifolia germanica, ſambuci odore. *Bauh. pin.* 31.

In Croatiæ montibus, qui a fluminenſibus *mala Planina* vocantur. Fl. Iunio. Per.

Pſeud-Acorus. 5. I. corollis imberbibus: laciniis interioribus ſtigmate minoribus, foliis enſiformibus. *Sp. pl. p.* 56. *Curtis lond. Fl. dan. t.* 494. *Scop. carn. tom.* 1. *p.* 38.

Iris caule inflexo, foliis enſiformibus, petalis erectis minimis, reflexis imberbibus. *Hall. hiſt. n.* 1260.

Acorus adulterinus. *Bauh. pin.* 34. *Blackw. herb. t.* 261.

In paluſtribus, foſſis, ad fluviorum ripas, lacus. Fl. Majo, Iunio. Per.

graminea. 6. I. corollis imberbibus, germinibus ſexangularibus, caule ancipiti, foliis linearibus. *Sp. pl. p.* 58. *Iacqu. auſtr. t.* 2. *Scop. carn. tom.* 1. *p.* 39.

Iris anguſtifolia ſexta. *Clus. hiſt. p.* 230.

In Auſtria in herbidis collibus inter *Gumpoldskirchen* & *Baaden.* In Carniolia in montibus tolmienſibus, & in monte *Nanas.* In Croatia in pratis montium *mala Planina* dictorum. In pratis agri fluminenſis in *Rezzine, Grohovo.* Fl. Majo, Iunio. Per.

ſpuria. 7. I. corollis imberbibus, germinibus ſexangularibus, caule tereti, foliis ſublinearibus. *Sp. pl. p.* 58. *Iacqu. auſtr. t.* 4.

Iris anguſtifolia prima. *Clus. hiſt. p.* 228.

In pratis circa *Miſtelbach, Waltersdorf,* & *Männersdorf,* alibique. Fl. Iunio. Per.

ſibirica. 8. I. corollis imberbibus, germinibus trigonis, caule tereti, foliis linearibus. *Sp. pl. p.* 57. *Iacqu. auſtr. t.* 3.

Iris foliis linearibus, caule ſubnudo, petalis reflexis, imberbibus, venoſis, tubalium arcu acuminato. *Hall. hiſt. n.* 1259.

In pratis udis planitiei, in Auſtria, Pannonia, Croatia. Fl. Majo, Iunio. Per.

SCHOENUS. Glumæ paleaceæ, univalves, congeſtæ. Corolla nulla. Semen unicum, ſubrotundum inter glumas, vel nudum, vel ſetis minimis, e receptaculo proprio enatis, cinctum.

Mariscus. 1. S. culmo tereti, foliis margine dorſoque aculeatis. *Sp. pl. p.* 62.

Mariscus panicula ramoſa, foliorum oris dorſoque ſerratis. *Hall. hiſt. n.* 1343.

Pſeudocyperus paluſtris, foliis & carina ſerratis. *Scheuch. agr.* 375. *t.* 8. *f.* 7, 8, 9, 10, 11.

In aquis ſtagnantibus, pratis paluſtribus, in Auſtria prope *Himberg*; in Pannonia ad lacum Peiſonis. Fl. Iulio. Per.

Eſt planta diandra, ferens nucem, includentem nucleum quadrialatum. Ehrh. beytr. tom. 4. p. 48.

nigricans. 2. S. culmo tereti nudo, capitulo ovato, involucri diphylli valvula altera ſubulata longa. *Sp. pl. p.* 64.

Cyperus ſpicis conglomeratis, calyce multifido. *Hall. hiſt. n.* 1347.

Iuncus lævis minor, panicula glomerata nigricante. *Moris. hiſt.* 3. *p.* 233. *ſect.* 8. *t.* 10. *f.* 28.

Iunco affinis, capitulo glomerato, nigricante. *Scheuch. agr.* 349. *t.* 7. *f.* 12, 13, 14.

In paluſtribus Auſtriæ, Pannoniæ. Fl. Iunio, Iulio. Per.

Huic ſpeciei eſt receptaculum nudum, ſemina ſunt nitida. Ehrh. l. c.

compreſſus. 3. S. culmo ſubtriquetro nudo, ſpica diſticha. *Sp. pl. p.* 65. *Leers herb. p.* 9. *t.* 1. *f.* 1. *Pollich palat. tom.* 1. p. 35. *f.* 2.

Scirpus *caricis*, culmo nudo ſubtriquetro, ſpica diſticha, involucro monophyllo. *Retzii ſcand. p.* 16. *obs. faſc.* 1. *p.* 10.

Scirpus planifolius, ſpica terminante diſticha. *Hall. hiſt. n.* 1342.

Carex uliginoſa. *Sp. pl. p.* 1381.

Gramen cyperoides, ſpica ſimplici compreſſa, diſticha. *Scheuch. agr.* 490. *t.* 11. *f.* 6.

In pratis paluſtribus, aliisque herbidis humidisque locis. Fl. Majo, Iunio. Per.

albus. 4. S. culmo ſubtriquetro folioſo, floribus faſciculatis, foliis ſetaceis. *Sp. pl. p.* 65. *Fl. dan. t.* 320. *Scop. carn. tom.* 1. *p.* 40.

Scirpus foliis ſetaceis, panicula plana, locuſtis teretibus. *Hall. hiſt. n.* 1341.

Gramen cyperoides paluſtre leucanthemum. *Scheuch. agr.* 503. *t.* 11. *f.* 11.

In paluſtribus. Fl. Iunio. Per.

Eſt planta diandra; ſtylum habet gluma paulo longiorem; ſetas receptaculi decem, longitudine ſeminis. Differt ergo a Schœno fuſco, cui ſunt ſtamina tria; ſtylus gluma duplo longior; ſetæ receptaculi tres, ſemine multo longiores. Ehrh. beytr. tom. 4. p. 154.

CYPERUS. Glumæ paleaceæ, diſtiche imbricatæ. Corolla nulla. Semen unicum, nudum.

longus. 1. C. culmo triquetro folioſo, umbella folioſa ſupradecompoſita, pedunculis nudis, ſpicis alternis. *Sp. pl. p.* 67. *Iacqu. collect. tom.* 3. *p.* 191. *ic. t.* 297. *Scop. carn. tom.* 1. *p.* 42.

Cyperus odoratus, radice longa, ſeu Cyperus officinarum. *Bauh. pin.* 14. *Scheuch. agr.* 378. *t.* 8. *f.* 12.

Le Souchet. Regn. bot.

In Auſtria ad Thermas Badenſes ad rivulos balnei S. Peregrini. In Litorali in foſſis Duini, Aquilejæ, nec non in Forojulio locis paluſtribus, & inundatis. Fl. Iulio, Auguſto. Per.

Semen eſt triquetrum.

flaveſcens. 2. C. culmo triquetro nudo, umbella triphylla, pedunculis ſimplicibus inæqualibus, ſpicis confertis lanceolatis. *Sp. pl. p.* 68. *Scop. carn. tom.* 1. *p.* 41.

Cyperus umbella trifolia, ſpicis ſeſſilibus, umbellatis, glumis obtuſis. *Hall. hiſt. n.* 1348.

Cyperus minor pulcher, panicula lata, compreſſa, ſubflaveſcente. *Moris. hiſt.* 3. *p.* 239. *ſect.* 8. *t.* 11. *f.* 37.

Cyperus minimus panicula ſparſa ſubflaveſcente. *Tourn. inſt.* 527. *Scheuch. agr.* 385.

Ad foſſas, lacus, fluvios, alibique. Fl. Iulio, Auguſto.

fuſcus. 3. C. culmo triquetro nudo, umbella trifida, pedunculis ſimplicibus inæqualibus, ſpicis confertis linearibus. *Sp. pl. p.* 69. *Fl. dan. t.* 179. *Scop. carn. tom.* 1. *p.* 41. *Leers herb. p.* 9. *t.* 1. *f.* 2.

Cyperus umbella trifolia, ſpicis petiolatis congeſtis, glumis ovato-lanceolatis. *Hall. hiſt. n.* 1349.

Cyperus minimus panicula ſparſa nigricante. *Tourn. inſt.* 527. *Scheuch. agr.* 384.

Habitat & floret cum præcedente.

Cyperus flaveſcens habet ſemen lentiforme; fuſcus vero triquetrum. Ehrh. beytr. tom. 4. p. 155.

pannonicus. 4. C. culmo triquetro, ſpicis ſeſſilibus, glumis lateralibus ſphacelatis. *Syſt. veget. p.* 98. *Iacqu. auſtr. app. t.* 6.

In Pannonia in litore lacus Peiſonis. Fl. Auguſto, Septembri.

Huic ſemina late ovata & compreſſa. Iacqu. l. c.

SCIRPUS. Glumæ paleaceæ, undique imbricatæ. Corolla nulla. Semen unicum, vel nudum, vel villis cinctum.

paluſtris. 1. S. culmo tereti nudo, ſpica ſubovata terminali. *Sp. pl. p.* 70. *Fl. dan. t.* 273. *Leers herb. p.* 10. *t.* 1. *f.* 3. *Scop. carn. tom.* 1. *p.* 43.

Scirpus caule terete, ſpica terete pauciflora. *Hall. hiſt. n.* 1336.

Scirpus equiſeti capitulo majore. *Tourn. inſt.* 528. *Scheuch. agr.* 360.

Ad rivulos, fluvios, lacus, foſſas. Fl. æſtate. Per.

Semen cingitur ſetis plerumque quatuor calyce brevioribus.

acicularis. 2. S. culmo tereti nudo ſetiformi, ſpica ovata bivalvi, ſeminibus nudis. *Sp. pl. p.* 71. *Fl. dan. t.* 287.

Mariscus foliis ſetaceis, mollibus, ſpica nuda pauciflora. *Hall. hiſt. n.* 1346.

In Auſtriæ arenoſis, inundatis in inſulis danubialibus, & ad fluvium Leitham. Fl. æſtate.

lacuſtris. 3. S. culmo tereti nudo, ſpicis ovatis pluribus pedunculatis terminalibus. *Sp. pl. p.* 72. *Scop. carn. tom.* I. *p.* 45.

Scirpus caule terete, panicula laterali, ramoſa, locuſtis ovatis. *Hall. hiſt. n.* 1337.

Scirpus paluſtris altiſſimus. *Tourn. inſt.* 528. *Scheuch. agr.* 354.

Iuncus aquaticus maximus. *Lob. ic. p.* 85.

In aquis ſtagnantibus, vel lente fluentibus. Fl. æſtate. Per.

Baſim germinis cingunt ſetæ quinque, vel ſex.

romanus. 4. S. culmo tereti nudo, capitulo laterali unico vel pluribus, ſeſſilibus pedunculatisque, bractea reflexa.

Scirpus culmo tereti nudo, capitulo laterali conglobato, bractea reflexa. *Sp. pl. p.* 72. *Iacqu. auſtr.* t. 448.

In paluſtribus Auſtriæ, Pannoniæ, Litoralis. Fl. æſtate. Per.

Variat, dum aut in omnibus ejusdem cespitis culmis capitulum ſeſſile ſolitarium conſpicitur, aut in alio ceſpite ſeſſile illud plura alia pedunculata ubique comitantur; aut dum in ejusdem plantæ ceſpite, alii culmi ſolitaria, alii plura capitula gerunt. At vero quandoquidem non minor apud nos numerus eſt culmorum multicapitatorum quam illorum, qui unico capitulo gaudent. Iacqu. l. c.

ſetaceus. 5. S. culmo nudo ſetaceo, ſpica terminali ſeſſili. *Syſt. veget. p.* 99. *Fl. dan. t.* 311. *Scop. carn. tom.* I. *p.* 46. *Leers herb. p.* 10. *t.* I. *f.* 6.

Mariſcus ſetaceus, capitulis lateralibus perpaucis. *Hall. hiſt. n.* 1345.

Scirpus omnium minimus, capitulo breviore. *Scheuch. agr.* 358.

In paluſtribus Duini, Aquilejæ Fl. Iunio.

mucronatus. 6. S. culmo triangulo nudo acuminato, ſpicis conglomeratis ſeſſilibus lateralibus. *Syſt. veget. p.* 100. *Scop. carn. tom.* 1. *p.* 45.

Scirpo-Cyperus panicula glomerata, e ſpicis imbricatis compoſita. *Scheuch. agr.* 404. *t.* 9. *f.* 14.

In Litorali in paluſtribus Duini, Aquilejæ; in ſimilibus locis in Auſtria, Pannonia, & ad Danubii ripas. Fl. Iulio, Auguſto. Per.

Ad baſin culmi folium unicum, breve, vaginans. Germen circumdant ſetæ tres.

maritimus. 7. S. culmo triquetro, panicula conglobata foliacea, ſpicularum ſquamis trifidis: intermedia ſubulata. *Sp. pl. p.* 74. *Curtis lond. Fl. dan. t.* 937. *Scop. carn. tom.* 1. *p.* 43.

Scirpus caule triquetro, panicula folioſa, ramoſa, locuſtis ovatis, nutantibus, glumis dentatis ariſtatis. *Hall. hiſt. n.* 1339.

Gramen panicula ſparſa ex ſpicis longioribus tenuioribus teretibus compoſita. *Scheuch. agr.* 398. *t.* 9. *f.* 7, 8.

In foſſis, inundatis. Fl. Iunio. Per.

Setæ quatuor aut quinque ad baſin germinis.

ſylvaticus. 8. S. culmo triquetro folioſo, umbella foliacea, pedunculis nudis ſupradecompoſitis, ſpicis confertis. *Sp. pl. p.* 75. *Fl. dan. t.* 307. *Leers herb. p.* 11. *t.* 1. *f.* 4.

Scirpus planifolius, panicula ramoſiſſima, locuſtis ovatis. *Hall. hiſt. n.* 1340.

Cyperus gramineus, *Bauh. hiſt.* 2. *p.* 504. *Scheuch. agr.* 393.

In udis, ad rivulos, præcipue in locis montanis. Fl. Iunio. Per.

Setis ſex circiter cingitur ſemen.

micheli-anus. 9. S. culmo triquetro, capitulo globoſo, involucro polyphyllo longo. *Sp. pl. p.* 76.

Cyperus italicus omnium minimus, locuſtis in capitulo ſubrotundo collectis. *Till. pis.* 51. *t.* 20. *f.* 5.

Tergeſti *alle Saule. I. Hoſt.*

ERIOPHORUM. Glumæ paleaceæ, undique imbricatæ. Corolla nulla. Semen unicum, lana longiſſima cinctum.

vagina-tum. 1. E. culmis vaginatis teretibus, ſpica ſcarioſa. *Sp. pl. p.* 76. *Curtis lond. Fl. dan. t.* 236.

Eriphorum caule terete, ſpica erecta ovata. *Hall. hiſt. n.* 1332.

Linagroſtis vaginata. *Scop. carn. tom.* 1. *p.* 47.

Iuncus alpinus, capitulo lanuginoſo, ſeu Schœnolagurus. *Bauh. pin.* 12. *prodr.* 23. *Scheuch. agr.* 302. *t.* 7. *f.* 1, 2, 3. *prodr. t.* 7.

In alpium udis. Fl. Iunio, Iulio. Per.

polyſta-chyon. 2. E. culmis teretibus, foliis planis, ſpicis pedunculatis. *Sp. pl. p.* 76. *Curtis lond. Leers herb. p.* 11. *t.* 1. *f.* 5.

Eriophorum foliis planis, ſpicis pendulis. *Hall. hiſt. n.* 1331.

Linagroſtis polyſtachya. *Scop. carn. tom.* 1. *p.* 48.

Linagroſtis panicula ampliore. *Tourn. inſt.* 664. *Scheuch. agr.* 306.

In pratis udis, montanis & demiſſis. Fl. Aprili, Majo. Per.

alpinum. 3. E. culmis nudis triquetris, ſpica pappo breviore. *Sp. pl. p.* 77. *Fl. dan. t.* 620.

Eriophorum caule foliisque triquetris, ſpica paucifiora. *Hall. hiſt. n.* 1333.

Linagroſtis alpina. *Scop. carn. tom.* 1. *p.* 48.

Linagroſtis juncea alpina, capitulo parvo, tomento rariore. *Scheuch. agr.* 305. *t.* 7. *f.* 4.

Iuncus alpinus bombycinus. *Bauh. pin.* 12. *prodr.* 6. *Scheuch. prodr. t.* 8.

In Carniolia in paludibus labacenſibus, & in udis alpium wochinenſium. Fl. Majo, Iunio. Per.

NARDUS. Calyx nullus. Corolla bivalvis.

ſtricta. 1. N. ſpica ſetacea recta ſecunda. *Syſt. veget. p.* 102. *Schreb. gr. t.* 7. *Scop. carn. tom.* 1. *p.* 49. *Leers herb. p.* 11. *t.* 1. *f.* 7. *Hall. hiſt. n.* 1410.

Gramen ſparteum junciſolium. *Bauh. pin.* 5. *Scheuch. agr.* 90. *t.* 2. *f.* 10.

In montanis & alpinis. Fl. Iunio, Iulio. Per.

ariſtata. 2. N. calycibus ariſtatis. *Sp. pl. p.* 78.

Nardus ſpica ſubulata, articulata, incurva, tereti. *Ger. prov. p.* 73.

Nardus ſpica incurva æquali, gluma calycina exteriore ariſtato-acuminata longiſſima, floribus diſtantibus alternis. *Gouan hort. p.* 33. *Fl. p.* 114.

Gramen junceum, nodoſum, minimum, capillare. *Barr. rar.* 106. *t.* 117. *f.* 1.

In Iſtria circa Petinam & Piſinum. *I. Hoſt.* Fl. Iunio.

DIGYNIA.

ALOPECURUS. Calyx bivalvis. Corolla univalvis.

pratenſis. 1. A. culmo ſpicato erecto, glumis villoſis, corollis muticis. *Sp. pl. p.* 88. *Curtis lond. Schreb. gr. t.* 19. *f.* 1. *Leers herb. p.* 15. *t.* 2. *f.* 4.

Gramen myurum ſpica molli candicante villoſa. *Scheuch. agr.* 70.

In pratis, præſertim humidis. Fl. æſtate. Per.

genicu-latus. 2. A. culmo ſpicato infracto, corollis muticis. *Syſt. veget. p.* 108. *Curtis lond. Leers herb. p* 16. *t.* 2. *f.* 7. *Scop. carn. tom.* 1. *p.* 58. *Hall. hiſt. n.* 1541.

Gramen aquaticum geniculatum ſpicatum. *Bauh. pin.* 3. *Scheuch. agr.* 72.

In uliginoſis, ad foſſas, in inundatis. Fl. æſtate. Per.

monſpeli-enſis. 3. A. panicula ſubſpicata, glumis ſcabris, corollis ariſtatis. *Sp. pl. p.* 89.

Alopecurus *ariſtatus*, panicula erecta ovata, glumis calycinis ovatis ariſtatis. *Gouan hort. p.* 37.

Phleum *crinitum*, calycibus linearibus baſi ſubventricoſis hiſpidis ariſtatis: ariſtis capillaceis longiſſimis. *Schreb. gr. p.* 151. *t.* 20. *f.* 3.

Agroſtis panicula oblonga, calycibus æqualibus; terminatis ariſtis longis rectis. *Ger. prov. p.* 80.

In agro tergeſtino circa *Dolline. I. Hoſt.*

ASPRELLA. Calyx nullus. Corolla bivalvis clauſa. Stamina corolla breviora.

oryzoides. 1. Asprella. *Retzii ſcand. p.* 17.

Phalaris *oryzoides*, panicula effuſa, glumarum carinis ciliatis. *Sp. pl. p.* 81. *Iacqu. en. p.* 12. *Schreb. gr. t.* 22.

Homalocenchrus. *Hall. hiſt. n.* 1411.

Gramen miliaceum paluſtre ſerotinum paniculis ex locuſtis compreſſis compoſitis. *Mont. prodr. p.* 51. *ic.* 58.

Gramen aquaticum aſperum ſerotinum & elatius, radice repente, orycæ panicula villoſa flaveſcente, ſemine griſeo milii aut lini. *Till. pis. p.* 73.

In Auſtriæ aquis ſtagnantibus in inſulis danubialibus, ad fluvium Leitham, & Fiſcham. Fl. Iulio, Auguſto. Per.

PHALARIS. Calyx bivalvis, carinatus, longitudine æqualis, corollam includens.

canarienſis. 1. P. panicula ſubovata ſpiciformi, glumis carinatis. *Sp. pl. p.* 79. *Schreb. gr. t.* 10. *f.* 2.

Phalaris panicula ſpiciformi ovata, calycibus carinatis piloſis, nectariis binis. *Retzii obs. fasc.* 3. *p.* 8.

Flumine, Lauranæ, vicinisque locis copioſe ad litus maris. Fl. Iunio.

utriculoſa. 2. P. panicula ſpicata, petalis ariſta articulata, vagina ſupremi folii ſpathiformi. *Sp. pl. p.* 80.

Gramen pratenſe, ſpica purpurea ex utriculo prodeunte, ſeu Gramen folio ſpicam amplexante. *Bauh. pin.* 3. *theatr.* 44.

Tergeſti *alle Saule. I. Hoſt.* Fl. Iunio.

phleoides. 3. P. panicula cylindrica ſpiciformi glabra, paſſim vivipara. *Syſt. veget. p.* 104. *Fl. dan. t.* 531.

In collibus apricis, ad vias, & alibi. Fl. Iunio. Per.

arundinacea. 4. P. panicula oblonga ventricoſa ampla. *Syſt. veget. p.* 104. *Leers herb. p.* 17. *t.* 7. *f.* 3.

Arundo locuſtis unifloris, faſciculatis, imbricatis, glumis floralibus nitentibus ſubhirſutis. *Hall. hiſt. n.* 1524.

Gramen miliaceum aquaticum, phalaridis ſemine. *Scheuch. agr.* 126. *t.* 3. *f.* 4.

In udis, ad foſſas, fluvios, lacus. Fl. Iunio, Iulio. Per.

Nectarii loco ſub quavis gluma corollina penicillum pilorum gerit. Retzii obs. faſc. 3. p. 8.

PANICUM. Calyx trivalvis: valvula tertia minima, uniflora.

verticillatum. 1. P. ſpica verticillata, racemis quaternis, involucellis unifloris biſetis, culmis diffuſis. *Sp. pl. p.* 82. *Curtis lond.*

Panicum ſpica unica, paniculata, ſetis perpaucis. *Hall. hiſt. n.* 1543.

Gramen paniceum, ſpica aſpera. *Bauh. pin.* 8. *Scheuch. agr.* 47.

In cultis, ruderatis, ad vias. Fl. Iulio, Auguſto.

glaucum. 2. P. ſpica tereti, involucellis bifloris faſciculato-ſetoſis, ſeminibus undulato-rugoſis. *Sp. pl. p.* 83. *Schreb. gr. t.* 25.

Panicum ſpica unica, floſculis ſeſſilibus ſolitariis, ſetis numeroſis. *Hall. hiſt. n.* 1542.

Cum præcedente.

viride. 3. P. ſpica tereti, involucellis bifloris faſciculato-piloſis, ſeminibus nervoſis. *Sp. pl. p.* 83. *Leers herb. p.* 13. *t.* 2. *f.* 2. *Curtis lond.*

Cum præc.

Crus Galli. 4. P. ſpicis alternis, conjugatisque, ſpiculis ſubdiviſis, glumis ariſtis hiſpidis, rachi quinquangulari. *Sp. pl. p.* 83. *Curtis lond. Leers herb. p.* 13. *t.* 2. *f.* 3. *Scop. carn. tom.* 1. *p.* 51.

Panicum ſpica ramoſa, ſetis nullis. *Hall. hiſt. n.* 1544.

In cultis, inundatis, ad foſſas, rivulos. Fl. Iulio, Auguſto.

Variat culmis erectis, & proſtratis; floſculis muticis, & ariſtatis.

hirtellum. 5. P. ſpica compoſita: ſpiculis adpreſſis alternis, calycibus geminis, valvulis omnibus ariſtatis; ariſta extima longiſſima. *Sp. pl. p.* 83. *Wulfen in Iacqu. collect. tom.* 1. *p.* 265.

Panicum foliis lanceolatis undulatis, caule repente, geniculis radicatis. *Arduin. animadv.* 2. *p.* 14. *t.* 4.

In Forojulio. Per.

ſanguinale. 6. P. ſpicis digitatis baſi interiore nodoſis, floſculis geminis muticis, vaginis foliorum punctatis. *Sp. pl. p.* 84. *Leers herb. p.* 14. *t.* 2. *f.* 6. *Schreb. gr. t.* 16.

Digitaria foliis ſubhirſutis, caule debili, ſpicis verticillatis, ſcapo ancipite. *Hall. hiſt. n.* 1526.

Digitaria ſanguinalis. *Scop. carn. tom.* 1. *p.* 52.

Iſchæmon. *Clus hiſt. p.* CCXVII.

Gramen dactylon, folio latiore. *Bauh. pin.* 8. *Scheuch. agr.* 101. *t.* 2. *f.* 2.

In cultis, & ſylvaticis. Fl. Iulio, Auguſto.

Dactylon. 7. P. ſpicis digitatis patentibus baſi interiore villoſis, floribus ſolitaris, ſarmentis repentibus. *Sp. pl. p.* 85.

Digitaria ſtolonibus teretibus, foliis patulis glabris, ſpicis digitatis. *Hall. hiſt. n.* 1527.

Digitaria Dactylon. *Scop. carn. tom.* 1 *p.* 52.

Panicum Dactylon, radice repente, ſeu officinarum. *Scheuch. agr.* 104.

Gramen legitimum. *Clus. hiſt. p. CCXVII.*

Le Chien dent, ou le Pied de Poule. Regn. bot.

In cultis, pratis, ruderatis, ad vias. Fl. Auguſto, Septembri. Per.

PHLEUM. Calyx bivalvis, ſeſſilis, linearis, truncatus, apice bicuſpidato, uniflorus. Corolla incluſa.

pratenſe. 1. P. ſpica cylindrica longiſſima ciliata, culmo erecto. *Syſt. veget. p.* 107. *Leers herb. p.* 16. *t.* 3. *f.* 1. *Scop. carn. tom.* 1. *p.* 54.

Phleum calycibus linearibus ciliatis truncatis ariſtatisque: ariſtis brevibus. *Schreb. gr. p.* 102. *t.* 14.

Phleum caule erecto, ſpica cylindrica longiſſima, glumis calycinis oblique truncatis. *Hall. hiſt. n.* 1528.

In pratis. Fl. Iunio, Iulio. Per.

nodoſum. 2. P. ſpica cylindrica, culmo adſcendente, foliis obliquis, radice bulboſa. *Syſt. veget. p.* 108. *Fl. dan. t.* 380. *Leers herb. p.* 17. *t.* 3. *f.* 2.

Phleum caule imo bulboſo, declinante, glumis calycinis obtuſe truncatis. *Hall. hiſt. n.* 1530.

In ſylvaticis, pratisque paluſtribus. Fl. cum præc. Per.

alpinum. 3. P. ſpica ovato-cylindracea. *Sp. pl. p.* 88. *Fl. dan. t.* 213. *Scop. carn. tom.* 1. *p.* 54.

Phleum ſpicis ovatis hirſutis, locuſtis longe bicornibus. *Hall. hiſt. n.* 1529.

Gramen typhoides alpinum; ſpica brevi, denſa & veluti villoſa. *Scheuch. agr.* 64. *prodr. t.* 3.

In pratis alpinis. Fl. æſtate Per.

Schœnoides. 4. P. ſpicis ovatis obvolutis, foliis breviſſimis mucronatis amplexicaulibus. *Sp. pl. p.* 88. *Iacqu. ic. t.* 15.

Gramen maritimum typhinum, brevi & craſſiori ſpica ad ſingula genicula prodeunte. *Scheuch. agr.* 86.

In Auſtriæ, Pannoniæ, Litoralis uliginoſis, paluſtribus. Fl. Iulio, Auguſto.

MILIUM. Calyx bivalvis, uniflorus: valvulis ſubæqualibus. Corolla breviſſima ſtigmata penicilliformia.

effuſum. 1. M. floribus paniculatis diſperſis muticis. *Sp. pl. p.* 90. *Curtis lond. Leers herb. p.* 18. *t.* 8. *f.* 7.

Milium paniculis raris, petiolis longiſſimis. *Hall. hiſt. n.* 1525.

Gramen ſylvaticum, panicula miliacea ſparſa. *Bauh. pin.* 8. *Scheuch. agr.* 133.

In ſylvis montanis, & ſubalpinis. Fl. Iunio, Iulio. Per.

parado-xum. 2. M. floribus paniculatis, ariſtatis. *Sp. pl. p.* 90. *Scop. carn. tom.* 1. *p.* 58. *t.* 1.

Agroſtis racemis ternatis pendulis; ſpiculis ariſtatis; ariſta undulata longa. *Scop. carn. ed.* 1. *p.* 186.

In Carniolia in ſylvis circa Idriam; prope viam regiam inter *Loitſch* & *Oberlaibach*. *Scop.* Ego in ſylvis agri fluminenſis in *Rezzine*, *Grohovo*, *Lopazza*. Fl. Iulio. Per.

AGROSTIS. Calyx bivalvis, uniflorus, corolla paulo minor. Stigmata longitudinaliter hiſpida.

Spica venti. 1. A. petalo exteriore ariſta recta ſtricta logiſſima, panicula patula. *Syſt. veget. p.* 110. *Fl. dan. t.* 853. *Leers herb. p.* 18. *t.* 4. *f.* 1.

Avena monantha, panicula adſcendente, multiflora, calyce lævi, florali ariſta longiſſima. *Hall. hiſt. n.* 1480.

Inter ſegetes, ad vias, & alibi. Fl. Iunio, Iulio.

Calamagroſtis. 2. A. panicula incraſſata, petalo exteriore toto lanato apice ariſtato, caule ramoſo. *Sp. pl. p.* 92.

Arundo locuſtis unifloris, ex ſumma gluma ariſtatis. *Hall. hiſt. n.* 1521.

Gramen arundinaceum panicula ariſtata, denſa, viridi-argentea, ſplendente. *Scheuch. agr.* 146. *t.* 3. *f.* 11. *Segu. ſuppl. p.* 163.

In rupeſtribus alpium tyrolenſium, carniolicarum. Fl. Iulio. Per.

arundinacea. 3. A. panicula oblonga, petalo exteriore baſi villoſo ariſtaque torta calyce longiore *Sp. pl. p.* 91.

Arundo locuſtis unifloris, ex imo dorſo ariſtatis, *Hall. hiſt. n.* 1522.

Gramen avenaceum montanum panicula anguſta, e dilutiſſimo fuſco albicante & papposa, *Scheuch. agr. app.* 507.

In ſylvis montanis, & ſubalpinis. Fl. Iulio, Auguſto, Per.

ſerotina. 4. A. panicula floſculis oblongis mucronatis, culmo obtecto foliis breviſſimis. *Syſt. veget. p.* 110.

Feſtuca ſerotina. *Sp. pl. p.* 111.

Gramen loliaceum ſerotinum, panicula diſpanſa. *Segu. ver.* 3. *p.* 146. *t.* 3. *f.* 2.

In Iſtria circa Petinam, & Piſinum. *I. Hoſt.* Per.

canina. 5. A. calycibus elongatis, petalorum ariſta dorſali recurva, culmis proſtratis ſubramoſis. *Sp. pl. p.* 92. *Leers herb. p.* 19. *t.* 4. *f.* 2.

Avena monantha, panicula diffuſa, multiflora, adſcendente, gluma calycina exaſperata. *Hall. hiſt. n.* 1479.

Gramen paniculatum ſupinum ad infima culmorum genicula foliorum capillarium faſciculis donatum. *Scheuch. agr.* 141. *t.* 3. *f.* 9.

Agroſtis alpina. *Scop. carn. tom.* 1. *p.* 60. *Fl. dan. t.* 161.

In pratis, collibus, & in alpibus. Fl. Iulio, Auguſto. Per.

Ariſta, quæ primo recta glumisque brevior, dein recurvatur ſitque glumis longior.

ſtolonifera. 6. A. panicula ramulis patentibus muticis, culmo repente, calycibus æqualibus. *Sp. pl. p.* 93. *Fl. dan. t.* 564.

Poa monantha, ſtolonifera, calycibus ſubaſperis. *Hall. hiſt. n.* 1473.

In ruderatis, ad vias, alibique. Fl. Iunio, Iulio. Per.

alba. 7. A. panicula laxa, calycibus muticis æqualibus, culmo repente. *Syſt. veget. p.* 111. *Leers herb. p.* 21. *t.* 4. *f.* 5.

In paluſtribus, ad foſſas, in inundatis. Fl. Iulio. Per.

capillaris. 8. A. panicula capillari patente, calycibus ſubulatis æqualibus hiſpidiuſculis, coloratis, floſculis muticis monopetalis. *Retzii ſcand. p.* 20. *Sp. pl. p.* 93. *Fl. dan. t.* 163. *Leers herb. p.* 20. *t.* 4. *f.* 3.

Poa monantha, caule erecto, panicula diffuſa, calycibus dorſo exaſperatis. *Hall. hiſt. n.* 1475.

Gramen montanum, panicula ſpadicea delicatiori. *Bauh. pin.* 3. *Scheuch. agr.* 129. *t.* 3. *f.* 5.

In pratis, collibus, aliiſque herbidis locis. Fl. Iulio. Per.

AIRA. Calyx bivalvis, biflorus. Floſculi abſque interjecto rudimento.

aquatica. 1. A. panicula patente, floribus muticis lævibus calyce longioribus, foliis planis. *Syſt. veget. p.* 112. *Curtis lond. Fl. dan. t.* 381. *Scop. carn. tom.* 1. *p.* 66.

Poa locuſtis bifloris, glabris, florali gluma majori, plicata, ſerrata. *Hall. hiſt. n.* 1471.

Gramen miliaceum, aquaticum. *Scheuch. agr.* 176.

In rivulis, inundatis. Fl. æſtate. Per.

ſubſpicata. 2. A. foliis planis, panicula ſpicata, floſculis medio ariſtatis: ariſta reflexa latiore. *Syſt. veget. p.* 112. *Sp. pl. p.* 95. *Fl. dan. t.* 228. *Wulfen in Iacqu. collect. tom.* 3. *p.* 32. *t.* 19. *f.* 3.

Avena diantha, panicula denſa, ſpicata. *Hall. hiſt. n.* 1490.

Gramen avenaceum paniculatum, alpinum, humile, locuſtis in ſpicam collectis, varicoloribus, ariſtatis. *Scheuch. agr.* 221. *prodr. t.* 6.

In alpibus. Fl. Iulio, Auguſto. Per.

ceſpitoſa. 3. A. foliis planis, panicula patente, petalis baſi villoſis ariſtatiſque: ariſta recta brevi. *Sp. pl.*

p. 96. *Fl. dan. t.* 240. *Leers herb. p.* 23. *t.* 4. *f.* 8. *Scop. carn. tom.* 1. *p.* 65.

Avena diantha, panicula ampliſſima, patula, bicolore. *Hall. hiſt. n.* 1487.

In pratis. Fl. Iunio, Iulio. Per.

flexuoſa. 4. A. foliis ſetaceis, culmis ſubnudis, panicula divaricata, pedunculis flexuoſis. *Sp. pl. p.* 96. *Fl. dan. t.* 157. *Leers herb. p.* 24. *t.* 5. *f.* 1.

Avena diantha, pediculis flexuoſis, floribus baſi villoſis, locuſtis ovatis. *Hall. hiſt. n.* 1486.

In collibus herbidis, montanis ſylvaticis. Fl. Iunio, Iulio. Per.

caryophyllea. 5. A. foliis ſetaceis, panicula divaricata, floribus ariſtatis diſtantibus. *Sp. pl. p.* 97. *Fl. dan. t.* 382. *Leers herb. p.* 24. *t.* 5. *f.* 7.

Avena diantha, foliis ſetaceis, panicula divaricata. *Hall. hiſt. n.* 1482.

Gramen phalaroides, ſparſa panicula, minimum anguſtifolium. *Barr. rar.* 1218. *t.* 44. *f.* 1.

In agro vindobonenſi in pratis ſiccioribus montis *Gallitzinberg.* Fl. Majo, Iunio.

MELICA. Calyx bivalvis, biflorus. Rudimentum floris inter flosculos.

ciliata. 1. M. flosculi inferioris petalo exteriore ciliato. *Sp. pl. p.* 97.

Arundo locuſtis bifloris, ſpicatis, gluma florali exteriore ciliata. *Hall. hiſt. n.* 1517.

Gramen avenaceum, ſpica ſimplici, locuſtis denſiſſimis, candicantibus & lanuginoſis. *Tourn. inſt.* 524. *Scheuch. agr.* 174.

Gramen montanum Avenæ ſemine. *Clus. hiſt. p.* CCXIX.

In ſaxoſis, aſperis, ac ad margines ſylvarum. Fl. Iunio, Iulio. Per.

nutans. 2. M. petalis imberbibus, panicula nutante ſimplici, calycibus trifloris. *Retzii ſcand.* *p.* 21. *Sp. pl. p.* 98. *Schreb. gr. t.* 6. *Leers herb. p.* 25. *t.* 3. *f.* 4.

Gramen montanum ſpicatum. *Clus. hiſt. p.* CCXIX.

In ſylvaticis. Fl. Majo. Per.

uniflora. 3. M. panicula rara, calycibus bifloris: floſculo altero hermaphrodito: altero ſterili. *Retzii ſcand. p.* 21. *obs. faſc.* 1. *p.* 10. *Curtis lond.*

In ſylvis umbroſis montanis Auſtriæ, Pannoniæ. Fl. Majo, Iunio. Per.

cœrulea. 4. M. panicula coarctata, floribus cylindricis. *Syſt. veget. p.* 113. *Curtis lond.*

Aira *cœrulea*, panicula coarctata, floribus pedunculatis muticis convoluto-ſubulatis, foliis planis. *Sp. pl. p.* 95. *Fl. dan. t.* 239. *Leers herb. p.* 22. *t.* 4. *f.* 7. *Gouan. illuſtr. p.* 3.

Arundo locuſtis bifloris, conicis, petiolis ſubvilloſis. *Hall. hiſt. n.* 1518.

Gramen paniculatum autumnale, panicula ampliore ex viridi nigricante. *Scheuch. agr.* 207. *t.* 4. *f.* 11, 12.

In Auſtria, Pannonia in paluſtribus, ad foſſas, atque in ſylvis montanis umbroſis. Fl. Auguſto, Septembri. Per.

Variat panicula parva, ramis coarctatis brevibus; & panicula ampliori, pedali, longioreque, ramis diffuſis, laxis.

POA. Calyx bivalvis, multiflorus. Spiculæ ovatæ, valvulis acutiuſculis, margine ſcarioſis.

aquatica. 1. P. panicula diffuſa, ſpiculis ſexfloris linearibus. *Sp. pl. p.* 98. *Fl. dan. t.* 920. *Leers herb. p.* 26. *t.* 5. *f.* 5. *Curtis lond. Scop. carn. tom.* 1. *p.* 75.

Poa altiſſima, foliis latiſſimis, panicula ampliſſima, locuſtis diſtichis multifloris. *Hall. hiſt. n.* 1454.

Gramen paluſtre paniculatum altiſſimum. *Bauh. pin.* 3. *Scheuch. agr.* 191.

Ad lacus, fluvios, in foſſis, rivulis. Fl. Iunio. Per.

Spiculæ variant floſculis a quinque ad octo.

alpina. 2. P. panicula diffuſa ramoſiſſima, ſpiculis ſexfloris cordatis. *Sp. pl. p.* 99. *Fl. dan. t.* 807.

Poa latifolia, culmo debili, locuſtis quadrifloris, calycis glumis æqualibus, floralium dorſo ſericeo. *Hall. hiſt. n.* 1456.

Gramen alpinum majus, panicula ſpecioſa, variegata. *Scheuch. agr.* 186. *prodr. t.* 3.

Gramen montanum, ſpica ſpecioſa graminea. *ejusd. it. alp.* 38. *t.* 4. *f.* 2.

In alpinis & ſubalpinis, varians ſpiculis viviparis. Creſcit quoque in Auſtriæ collibus herbidis prope *Bertholdsdorf*, *Medling*, *Baaden*, alibique. Fl. æſtate. Per.

trivialis. 3. P. panicula diffuſa, ſpiculis ſubtrifloris, culmo erecto ſcabro, ligula foliorum acuminata. *Curtis lond. Sp. pl. p.* 99.

In paſcuis præcipue humidiusculis. Fl. Majo, Iunio. Per.

Radix fibroſa. Culmus erectus, baſi repens. Folia ſcabriuscula. Ligula ad baſin foliorum longa, acuminata. Panicula erecta, diffuſa. Spiculæ parvæ, biſloræ, aut triſloræ, quandoque quadriſloræ, ovato-acuminatæ, ſubcompreſſæ. Glumæ calycinæ bivalves; valvulis inæqualibus, acuminatis, carina ſcabris. Lanugo ad baſin petali. Valvulæ corollæ ſubæquales, acutæ. Semen oblongo-acuminatum, anguloſum, baſi lanugine cinctum. Curtis l. c.

pratenſis. 4. P. panicula diffuſa, ſpiculis ſubquinquefloris, culmo erecto lævi, ligula foliorum obtuſa. *Curtis lond. Sp. pl. p.* 99.

Ubique in pratis. Fl. cum præc. Per.

Radix repens. Culmi erecti. Folia lævia, ligula brevi, obtuſa inſtructa. Panicula erecta, diffuſa. Spiculæ ovatæ, acutæ, utrinque compreſſæ, plerumque quinquefloræ. Calycis valvulæ acuminatæ. Corollæ valvulæ ſubæquales: altera concava, carina ſcabriuſcula; altera plana. Lanugo ad baſin petali. Semen anguloſum, acuminatum, baſi lana circumdatum. Curtis l. c.

anguſtifolia. 5. P. panicula diffuſa, ſpiculis quadrifloris pubeſcentibus, culmo erecto tereti. *Sp. pl. p.* 99. *Leers herb. p.* 27. *t.* 6. *f.* 3.

Poa foliis duris, ſe convolventibus, locuſtis quadrifloris, villoſis. *Hall. hiſt. n.* 1460.

In pratis, ad margines agrorum, & alibi. Fl. Majo. Per.

annua. 6. P. paniculis diffuſis angulis rectis, ſpiculis obtuſis, culmo obliquo compreſſo. *Sp. pl. p.* 99. *Curtis lond. Leers herb. p.* 29. *t.* 6. *f.* 1. *Scop. carn. tom.* 1. *p.* 71.

Poa culmo infracto, panicula triangulari, locuſtis trifloris, glabris. *Hall. hiſt. n.* 1466.

In cultis, ruderatis, ad vias. Fl. a primo vere ad ſerum autumnum.

rigida. 7. P. panicula lanceolata ſubramoſa, floribus alternis ſecundis. *Sp. pl. p.* 101. *Curtis lond.*

Gramen loliaceum, aſperum locuſtis unam partem ſpectantibus. *Cat. bon. prodr.* 42. *Zann. iſt. p.* 123. *t.* 171.

Flumine in herbidis. Fl. Iunio.

compreſſa. 8. P. panicula coarctata ſecunda, culmo obliquo compreſſo. *Sp. pl. p.* 101. *Leers herb. p.* 29. *t.* 5. *f.* 4. *Fl. dan. t.* 742. *Scop. carn. tom.* 1. *p.* 72.

Poa culmo compreſſo, declinato, locuſtis diſtichis, ſexfloris. *Hall. hiſt. n.* 1455.

Gramen paniculatum, radice repente, culmo compreſſo, locuſtis diſtichis, pulchellis. *Scheuch. agr.* 198.

In muris hortorum veſtutis, in uliginoſis, ruderatis, ad vias. Fl. Iulio, Auguſto. Per.

nemoralis. 9. P. panicula attenuata, ſpiculis ſubbifloris mucronatis ſcabris, culmo incurvo. *Sp. pl. p.* 102. *Fl. dan. t.* 749. *Leers herb. p.* 29. *t.* 5. *f.* 3.

Poa culmo debili, panicula nutante, locuſtis bifloris, folliculis ſubvilloſis. *Hall. hiſt. n.* 1469.

Gramen paniculatum anguſtifolium alpinum, locuſtis rarioribus & anguſtioribus non ariſtatis. *Scheuch. agr.* 164. *prodr. t.* 2.

In ſylvis montanis, ſubalpinis; & pratis alpinis. Fl. Iunio, Iulio. Per.

bulboſa. 10. P. panicula ſecunda patentiuſcula, ſpiculis quadrifloris. *Sp. pl. p.* 102. *Scop. carn. tom.* 1. *p.* 71.

Poa foliis bulboſis, panicula diffuſa, locuſtis quadrifloris, ſubvilloſis. *Hall. hiſt. n.* 1461.

Gramen vernum, radice Aſcalonitis. *Vaill. paris. t.* 17. *f.* 1.

Gramen arvenſe anguſtifolium, panicula denſa foliacea, foliolis in panicula anguſtiſſimis. *Scheuch. agr.* 212.

In cultis, ruderatis, ad vias, in apricis herbidis, occurrit ſpiculis plerumque viviparis. Fl. primo vere.

diſtans. 11. P. panicula ramis ſubdiviſis, floribus quinquefloris: floſculis diſtantibus obtuſis. *Syſt. veget. p.* 115.

Poa *retroflexa*, panicula diffuſa, ramis poſt floreſcentiam retroflexis, ſpiculis ſublinearibus ſubquinquefloris. *Curtis lond.*

Vulgaris in pratis ſiccioribus, locis arenoſis, ruderatis, ad vias Auſtriæ, Pannoniæ. Per.

criſtata. 12. P. panicula ſpicata, calycibus ſubpiloſis ſubquadrifloris pedunculo longioribus, petalis ariſtatis. *Syſt. veget. p.* 115. *Leers herb. p.* 30. *t.* 5. *f.* 6.

Feſtuca locuſtis bifloris, mucronatis, confertim imbricatis, panicula ſpicata. *Hall. hiſt. n.* 1444.

Aira criſtata. *Sp. pl. p.* 94. *Scop. carn. tom.* 1. *p.* 65.

Gramen ſpica criſtata, ſubhirſutum. *Bauh. pin.* 13. *prodr.* 8. *Scheuch. agr.* 166.

In apricis ſiccis herbidis; nec non in pratis humidis, ſylvisque umbroſis. Fl. Iunio, Iulio. Per.

In apricis paniculæ rami ſunt breves, culmoque adpreſſi; in humidis, atque umbroſis elonganatur, & patent.

diſticha. 13. P. ſpica diſticha terminali, hinc convexula, illinc plana. *Syſt. veget. p.* 115. *Wulfen in Iacqu. miſcell. tom.* 2. *p.* 74. *ic. t.* 19.

Seſleria locuſtis quinquefloris; diſtichis, calyce bidentato. *Hall. hiſt. n.* 1447.

Poa ſeſleroides. *Allion. pedem. tom.* 2. *p.* 246. *t.* 91. *f.* 1.

In ſummis Tyrolis, Carinthiæ, Styriæ ſuperioris alpium jugis. Fl. Iulio, Auguſto. Per.

BRIZA. Calyx bivalvis, multiflorus, flores in ſpiculam diſticham colligens. Valvulæ obtuſæ, ventricoſæ: interiore minuta.

media. 1. B. ſpiculis ovatis, calyce floſculis ſeptem breviore. *Sp. pl. p.* 103. *Fl. dan. t.* 258. *Leers herb. p.* 26. *t.* 7. *f.* 2.

Poa petiolis tenuiſſimis, locuſtis ovatis. *Hall. hiſt. n.* 1448.

Gramen tremulum majus. *Bauh. pin.* 2. *theatr.* 22. *Scheuch. agr.* 204. *t.* 4. *f.* 8.

Amourettes tremblantes. Clus. hiſt. p. CCXVIII.

In pratis. Fl. Iunio. Per.

maxima. 2. B. ſpiculis cordatis, floſculis ſeptemdecim. *Sp. pl. p.* 103.

Gramen paniculatum locuſtis maximis candicantibus, & phœniceis, tremulis. *Tourn. inſt.* 523.

Gramen tremulum maximum. *Bauh. pin.* 2. *prodr.* 5. *Scheuch. agr.* 202. *t.* 4. *f.* 7.

Tergeſti *alle Saule. I. Hoſt.* Fl. Iunio.

Eragroſtis. 3. B. ſpiculis lanceolatis, floſculis viginti. *Sp! pl. p.* 103.

Gramen paniculis elegantiſſimis. *Bauh. pin.* 2. *Scheuch. agr.* 194. *t.* 4. *f.* 4.

Gramen Eranthemum, ſeu Eragroſtis, Phalaroides paniculatum multiplici, tenui Brizæ ſpica. *Barr. rar.* 43. *t.* 4. *f.* 4.

Gramen *Amourettes. Clus. hiſt. p.* CCXVIII.

In cultis, ruderatis, ad vias Auſtriæ, Litoralis, Forijulii. Fl. a fine Iunii ad autumnum.

DACTYLIS. Calyx bivalvis, compreſſus; altera valvula majore carinata.

glomerata. 1. D. panicula ſecunda glomerata. *Sp. pl. p.* 105. *Fl. dan. t.* 743. *Leers herb. p.* 22. *t.* 3. *f.* 3. *Schreb. gr. t.* 8.

Bromus locuſtis quadrifloris, faſciculatis, imbricatis. *Hall. hiſt. n.* 1512.

Bromus glomeratus. *Scop. carn. tom.* I. *p.* 76.

Gramen ſpicatum folio aſpero. *Bauh. pin.* 3. *prodr.* 9. *Scheuch. agr.* 299. *t.* 6. *f.* 15.

In cultis, pratis, ruderatis, ad vias. Fl. æſtate. Per.

CYNOSURUS. Calyx bivalvis, multiflorus. Receptaculum proprium unilaterale, foliaceum.

criſtatus. 1. C. bracteis pinnatifidis. *Sp. pl. p.* 105. *Schreb. gr. t.* 8. *f.* 2. *Fl. dan. t.* 238. *Leers herb. p.* 47. *t.* 7. *f.* 4.

Cynoſurus bracteis pinnatis, retuſis. *Hall. hiſt. n.* 1545.

Phleum criſtatum. *Scop. carn. tom.* 1. *p.* 57.

Gramen pratenſe, criſtatum, ſeu ſpica criſtata lævi. *Bauh. pin.* 3, *Scheuch. agr.* 79.

In pratis. Fl. Iunio. Per.

echinatus. 2. C. bracteis pinnato-paleaceis, ariſtatis. *Sp. pl. p.* 105.

Cynoſurus bractearum dentibus lanceolato-linearibus. *Hall. hiſt. n.* 1546.

Gramen alopecuroides, ſpica aſpera. *Bauh. pin.* 4. *Scheuch. agr.* 80.

Gramen alopecurum ſpica aſpera. *Barr. rar.* 123.

Tergeſti *alle Saule. I. Hoſt.* Fl. Iunio.

durus. 3. C. ſpiculis alternis ſecundis ſeſſilibus rigidis obtuſis appreſſis. *Sp. pl. p.* 105. *Pollich. palat. tom.* 1. *p.* 98. *f.* 1.

Gramen arvenſe, polypodii panicula craſſiore. *Barr. rar.* 1213. *t.* 50.

In cultis, ruderatis, ad vias Auſtriæ, Litoralis. Fl. Majo, Iunio.

ſphærocephalus. 4. C. bracteis integris, ſpicis globoſis. *Syſt. veget. p.* 117. *Wulfen in Iacqu. miscell. tom.* 2. *p.* 71. *ic. t.* 20.

Sesleria ſphærocephala; caule ſimplici, nudo; ſpica ſubrotunda, involucrata. *Arduin. animadv. p.* 20. *t.* 2.

In ſummis jugis alpium Kibeggenſium vallis gilienſis; in Fladnizenſibus alpibus *Winter-*

thal, *Eiſenhut* in Carinthia; & in leontinis in Tyroli. *Wulfen.* Ego in Carniolia in alpibus wochinenſibus. Fl. Iulio. Per.

cæruleus. 5. C. bracteis integris, ſpicis ſubovatis. *Syſt. veget. p.* 117. *Wulfen in Iacqu. miscell. tom.* 2. *p.* 66. *ic. t.* 21.

Sesleria locuſtis trifloris, imbricatis, flore tricorni. *Hall. hiſt. n.* 1446.

Sesleria cœrulea. *Scop. carn. tom.* 1. *p.* 63.

Aira varia. *Iacqu. en. p.* 15.

Gramen glumis variis. *Bauh. pin.* 10. *Scheuch. agr.* 83. *t.* 2. *f.* 9.

In alpibus, collibus apricis, montibus, pratis planitiei. Pro diverſa ſoli elevatione Fl. ab Aprili ad Iulium. Per.

Variat ſpicis cæruleis, & albis.

FESTUCA. Calyx bivalvis floſculos in ſpiculam tenuem, oblongam, teretiuſculam colligens. Corollæ valvula inferior valvula interiore, & calyce major, teretiuſcula, acuminata, in mucronem terminata.

ovina. 1. F. panicula ſecunda coarctata ariſtata, culmo tetragono nudiuſculo, foliis ſetaceis. *Sp. pl. p.* 108. *Leers herb. p.* 31. *t.* 8. *f.* 3.

Bromus ovinus. *Scop. carn. tom.* 1. *p.* 77.

Gramen foliis junceis brevibus, majus, radice nigra. *Bauh. pin.* 5. *prodr.* 34. *Scheuch. agr.* 279.

In collibus apricis, aridis, & in alpibus. Fl. Majo, Iunio. Per.

rubra. 2. F. panicula ſecunda ſcabra, ſpiculis ſexfloris ariſtatis: floſculo ultimo mutico, culmo ſemitereti. *Sp. pl. p.* 109. *Leers herb. p.* 32. *t.* 8. *f.* 1.

Gramen alpinum pratenſe, panicula duriore laxa ſpadicea: locuſtis majoribus. *Scheuch. agr.* 287. *t.* 6. *f.* 9.

In montanis, & alpinis. Fl. Iunio, Iulio. Per

amethyſtina. 3. F. panicula flexuoſa, ſpiculis ſecundis inclinatis ſubmuticis, foliis ſetaceis. *Sp. pl. p.* 109.

Gramen montanum, foliis capillaribus longioribus, panicula heteromala ſpadicea & velut amethyſtina. *Scheuch. agr.* 276. *t.* 6. *f.* 7.

In collibus apricis, in pratis ſiccioribus Auſtriæ, Pannoniæ, Styriæ. Fl. Iunio. Per.

duriuſcula. 4. F. panicula ſecunda oblonga, ſpiculis ſexfloris oblongis lævibus, foliis ſetaceis. *Sp. pl. p.* 108. *Fl. dan. t.* 848.

Gramen tenue duriuſculum & pene junceum. *Bauh. hiſt.* 2. *p.* 463.

Gramen pratenſe, panicula duriore, laxa, unam partem ſpectante. *Raj. hiſt.* 1286. *Scheuch. agr.* 285.

In pratis, locisque montoſis ſylvaticis. Fl. Majo, Iunio. Per.

dumetorum. 5. F. panicula ſpiciformi pubescente, foliis filiformibus. *Sp. pl. p.* 109. *Fl. dan. t.* 700.

In pratis, ad margines ſylvarum. Fl. Iunio. Per.

elatior. 6. F. panicula ſecunda erecta, ſpiculis ſubariſtatis: exterioribus teretibus. *Sp. pl. p.* 111. *Schreb. gr. t.* 2. *Leers. herb. p.* 34. *t.* 8. *f.* 6.

Gramen paniculatum elatius, paniculis, ſeu ſpicis muticis, ſquamoſis. *Raj. cat. p.* 153. *Scheuch. agr.* 202.

In pratis, ad vias. Fl. æſtate. Per.

ſpadicea. 7. F. panicula erecta, ſpiculis ovatis quadrifloris, glumis acuminatis muticis, foliis ſetaceis glabris. *Smith linn. trans. on the Feſt. Syſt. veget. p.* 118. *Gouan. illuſtr. p.* 4.

Poa panicula erecta, ſpiculis trifloris glabris, corollis acuminatis, calyce duplo longioribus. *Ger. prov. p.* 91. *t.* 2. *f.* 1.

Poa culmo recto, locuſtis trifloris glaberrimis, calycibus ariſtatis. *Hall. hiſt. n.* 1463.

Feſtuca varia. *Hænke. in Iacqu. collect. tom.* 2. *p.* 94.

Poa Gerardi. *Allion. pedem. tom.* 2. *p.* 245.

Anthoxanthum paniculatum. *Sp. pl. p.* 40.

Gramen ſparteum, panicula flaveſcente. *Rudbeck elys. tom.* 1. *p.* 40. *f.* 14.

Gramen paniculatum, alpinum, radice craſſiſſima, foliis rigidis, ſtriatis, & aſperis, panicula fuſca non ariſtata. *Mich. hort. pis.* 75.

Gramen montanum panicula ſpadicea craſſiore. *Tourn. inſt.* 524.

Nardus ſpuria narbonenſis. *Bauh. pin.* 13.

Nardus Gangitis ſpuria Narbonæ. *Lobel. adv.* 43.

In Auſtria in ſaxoſis, collibusque apricis circa *Medling*, *Briel.* In Styria in monte *Schakl* prope Græcium; copioſiſſime in alpibus judenburgenſibus, ſekavienſibus. In Tyroli in alpibus leontinis, & in montibus prope *Sterzing.* Fl. Iunio, Iulio, Auguſto. Per.

Tota planta glabra. Radix fibroſa emittit foliorum & culmorum denſum ceſpitem. Culmi ſunt erecti, ſtricti, teretes, ſtriati; geniculo uno alterove purpureo. Folia involuto-ſetacea, ſtricta, ſtriata, mucronato-pungentia, glauca; baſi dilatato-membranacea, vaginantia, albida. Ligulæ breviſſimæ, aut nullæ. Panicula erecta, ramoſa, multiflora, laxa, aureo-ſpadicea, ramis plerumque binis. Pedunculi angulati, ſubflexuoſi, erecti. Spiculæ ovatæ, compreſſæ, plerumque quadrifloræ. Calycis valvulæ ſubæquales, carinatæ, acuminatæ, nec ariſtatæ, membranaceo-pellucidæ, baſi fuſcæ. Corollæ bivalves, calyce longiores, acuminatæ; valvula majori carinata alteram amplectente. Stamina tria. Styli duo breviſſimi. Stigmata plumoſa. Semen unicum, oblongum,

utrinque acutum, ſuperne ſulco longitudinali notatum, fuſcum. Smith. l. c.

Culmi apud nos ſunt modo pedales, modo bi- aut tripedales. Panicula plus minusve ramoſa. Spicularum color, qui initio ex ſpadiceo, fuſco, viridi, aureo varius, in ſeneſcente panicula evadit aureo-ſpadiceus.

decumbens. 8. F. panicula erecta, ſpiculis ſubovatis muticis, calyce floſculis majore, culmo decumbente. *Sp. pl. p.* 110. *Leers herb. p.* 34. *t.* 7. *f.* 5. *Fl. dan. t.* 162.

Gramen montanum, avenaceum, locuſtis muticis, tumentibus, piloſum. *Scheuch. agr.* 170. *t.* 3. *f.* 16.

In pratis ſiccioribus, paſcuisque montanis. Fl. Iunio. Per.

fluitans. 9. F. panicula ramoſa erecta, ſpiculis ſubſeſſilibus teretibus muticis. *Sp. pl. p.* 111. *Fl. dan. t.* 237. *Schreb. gr. t.* 3. *Curtis lond. Leers herb. p.* 35. *t.* 8. *f.* 5.

Poa locuſtis teretibus multifloris, glumis floralibus exterioribus truncatis, interioribus bifidis. *Hall. hiſt. n.* 1453.

Poa fluitans. *Scop. carn. tom.* 1. *p.* 73.

Gramen aquaticum fluitans, multiplici ſpica. *Bauh. pin* 3. *Scheuch agr.* 199. *t.* 4. *f.* 5.

In foſſis, lacubus, rivulis. Fl. æſtate. Per.

BROMUS. Calyx bivalvis flores multos in ſpiculam oblongam, diſticham colligens. Corolla bivalvis: valvula inferior major, obtuſa, bifida, ariſtam infra apicem rectam emittens; valvula ſuperior parva, mutica.

ſecalinus. 1. B. panicula patente, ſpiculis ovatis, ariſtis rectis, ſeminibus diſtinctis. *Syſt. veget. p.* 119.

Feſtuca graminea, glumis hirſutis, *Bauh. pin.* 9. *Scheuch. agr.* 250. *t.* 5. *f.* 9.

Inter ſegetes, & alibi. Fl. æſtate.

mollis. 2. B. panicula erectiuscula, ſpiculis ovatis pubescentibus: ariſtis rectis, foliis molliſſime villoſis. *Syſt. veget. p.* 117. *Schreb. gr. t.* 6. *Leers herb. p.* 36. *t.* 11. *f.* 1.

Gramen avenaceum locuſtis villoſis anguſtis candicantibus & ariſtatis. *Tourn. inſt.* 526. *Scheuch. agr.* 254. *t.* 5. *f.* 12.

In agris, ruderatis, pratis, ad vias. Fl. Iunio, Iulio.

ſquarroſus. 3. B. panicula nutante, ſpicis ovatis: ariſtis divaricatis. *Sp. pl. p.* 112.

Feſtuca graminea, glumis vacuis. *Scheuch. agr.* 251. *t.* 5. *f.* 11.

In pratis ſiccis agri vindobonenſis circa *Pœtzelsdorf.* Fl. Majo, Iunio.

inermis. 4. B. panicula erecta, ſpiculis ſubteretibus ſubulatis nudis ſubmuticis. *Syſt. veget. p.* 119. *Schreb. gr. t.* 13.

Bromus panicula diffuſa, ſpiculis teretibus ſubulatis muticis ariſtatisve. *Leys. hal.* 90.

In pratis, montoſis ſylvaticis, ad vias. Fl. Iunio. Per.

aſper. 5. B. panicula nutante ſcabra, ſpiculis villoſis ariſtatis, foliis ſcabris. *Syſt. veget. p.* 119. *Retzii ſcand. ed.* 2. *p.* 25.

Bromus *hirſutus*, panicula nutante ſcabra, ſpiculis teretibus ſublinearibus decemfloris, ariſtis rectis, vaginis foliorum hirſutis. *Curtis lond. Sibthorp. oxon. p.* 48.

Bromus *montanus*, panicula nutante, ſpiculis ſubſexfloris, ariſtis longioribus. *Pollich. palat. p.* 116. *Scop. carn. tom.* 1. *p.* 82. *Retzii ſcand. ed.* 1. *p.* 19. *obs. faſc.* 2. *p.* 7.

Bromus ramoſus. *Huds. angl. ed.* 1. 40.

Bromus nemoralis. *ejusd. angl. ed.* 2. 51.

In montanis ſylvaticis. Fl. Iunio, Iulio. Per.

ſterilis. 6. B. panicula patula, ſpiculis oblongis diſtichis, glumis ſubulato-ariſtatis. *Sp. pl. p.* 113. *Curtis lond. Leers herb. p.* 37. *t.* 11. *f.* 4.

Bromus panicula nutante, locuſtis ſeptifloris, glumis argute lanceolatis, lineatis, ſubhirſutis. *Hall. hiſt. n.* 1505.

Gramen avenaceum, panicula ſparſa, locuſtis majoribus, & ariſtatis. *Scheuch. agr.* 258. *t.* 5. *f.* 14.

Ad pagos, agrorum margines, ſepes, vias. Fl. Iunio, Iulio.

In hoc pedunculi ſcabri, minime piloſi; in ſequenti piloſi, minime ſcabri. Retzii obs. fasc. 1. p. 11.

tectorum. 7. B. panicula nutante, ſpiculis linearibus. *Sp. pl. p.* 114. *Leers herb. p.* 38. *t.* 10. *f.* 2.

Bromus hirſutus panicula nutante, pauciflora, locuſtis quinquefloris, hirſutis, glumis argute lanceolatis. *Hall. hiſt. n.* 1508.

In ruderatis, muris, tectis, & alibi. Fl. Majo.

Paniculas digitis coarctatas repetitis vicibus naribus inſerunt extrahuntque ſanguinis ciendi cauſa fluminenſes pueri.

arvenſis. 8. B. panicula nutante, ſpiculis ovato-oblongis. *Sp. pl. p.* 113. *Leers herb. p.* 37. *t.* 11. *f.* 3. *Fl. dan. t.* 293.

Bromus glaber, locuſtis glabris, nutantibus, octifloris, glumis marginatis. *Hall. hiſt. n.* 1509.

Feſtuca graminea, effuſa juba. *Bauh. pin.* 9. *prodr.* 66. *Scheuch. agr.* 262. *t.* 5. *f.* 15.

In arvis, ad vias, agrorumque margines. Fl. Iulio.

giganteus. 9. B. panicula nutante, ſpiculis quadrifloris: ariſtis brevioribus. *Sp. pl. p.* 114. *Curtis lond. Leers herb. p.* 38. *t.* 10. *f.* 1. *Schreb. gr. t.* 11.

Gramen bromoides, aquaticum, latifolium, panicula ſparſa, tenuiſſime ariſtata. *Scheuch. agr.* 264. *t.* 5. *f.* 17.

In ſylvis, ad fluvios. Fl. Iulio, Auguſto, Septembri. Per.

rubens. 10. B. panicula fasciculata, ſpiculis ſubſeſſilibus villoſis, ariſtis erectis. *Sp. pl. p.* 114.

Tergeſti *alle Saule*, & in Iſtria. *I. Hoſt.* Fl. Iunio.

pinnatus. 11. B. ſpiculis ſeſſilibus, culmo indiviſo, ariſtis corollæ gluma brevioribus. *Pollich. palat. tom.* 1. *p.* 117. *Sp. pl. p.* 115. *Leers herb. p.* 39. *t.* 10. *f.* 3. *Fl. dan. t.* 164.

Triticum ſpicis diſtichis, locuſtis teretibus, hirſutis, ariſtatis, duodecimfloris. *Hall. hiſt. n.* 1431.

Feſtuca *pinnata*, ſpicata ſubdiſticha; ſpiculis alternis teretibus ſubariſtatis. *Huds. angl. ed.* 2. 48. *Sibthorp. oxon. p.* 46.

In pratis, collibus apricis, montoſis ſylvaticis, ad vias. Fl. Iunio, Iulio. Per.

ſylvaticus. 12. B. ſpiculis ſeſſilibus, culmo indiviſo, ariſtis corollæ gluma longioribus. *Pollich. palat. tom.* 1. *p.* 118. *Fl. dan. t.* 164.

Feſtuca *ſylvatica*, ſpicata, ſpiculis alternis ſubdiſtichis ſubſeſſilibus ariſtatis. *Huds. angl. ed.* 1. 41. *Sibthorp. oxon. p.* 46.

Triticum locuſtis remotis, hirſutis, teretibus, exiguis, culmis floriferis gracilibus. *Hall. hiſt. n.* 1432.

Gramen loliaceum, corniculatum, montanum, ſpica partiali ſubhirſuta, fragili. *Scheuch. agr.* 38.

In montanis ſylvaticis, nec non in ſylvis demiſſis ad fluvios. Fl. cum præc. Per.

Præcedenti radix repens, huic minime.

STIPA. Calyx bivalvis, uniflorus. Corolla bivalvis: valvula exterior apice terminata ariſta longiſſima, tortili, baſi articulata, recta; valvula interior linearis, mutica.

pennata. 1. S. ariſtis lanatis. *Sp. pl. p.* 115. *Scop. carn. tom.* 1. *p.* 85.

Stipa ariſtis plumoſis. *Hall. hiſt. n.* 1514.

Spartum auſtriacum pennatum. *Cluſ. hiſt. p.* CCXXI.

In collibus apricis, aſperis. Fl. Majo, Iunio.

juncea. 2. S. ariſtis nudis rectis, calycibus ſemine longioribus, foliis intus lævibus. *Syſt. veget. p.* 121. *Hall. hiſt. n.* 1513.

Feſtuca junceo folio. *Bauh. pin.* 9. *prodr.* 66.

Feſtuca longiſſimis ariſtis. *Bauh. pin.* 10. *Scheuch. agr.* 151. *t.* 3. *f.* 13.

In collibus, campis, Auſtriæ, Pannoniæ. Fl. Iulio. Per.

AVENA. Calyx bivalvis, ſæpius multiflorus. Corolla bivalvis: valvula inferior utrinque acuminata, e dorſo ariſtam ſpiraliter intortam, geniculo veluti reflexam, emittens.

elatior. 1. A. paniculata, calycibus bifloris, floſculo hermaphrodito ſubmutico, maſculo ariſtato. *Sp. pl. p.* 117. *Fl. dan. t.* 165. *Leers herb. p.* 40. *t.* 10. *f.* 4. *Curtis lond.*

Avena diantha, folliculis baſi villoſis, majoris ariſta geniculata. *Hall. hiſt. n.* 1492.

Avena calycibus biſloris, panicula laxe ſpicata: pedunculis prioribus faſciculatis. *Guettard. obs.* 1. *p.* 182.

Holcus *avenaceus*, glumis bifloris, ariſta floris maſculi dorſali refracta. *Wigg. prim. p.* 72. *Sibthorp. oxon. p.* 4. *Scop. carn. tom.* 2. *p.* 276.

Gramen avenaceum, elatius, juba longa ſplendente. *Raj. meth.* 179. *Scheuch. agr.* 239.

Gramen nodoſum, avenacea panicula. *Bauh. pin.* 2. *prodr.* 3. *Scheuch. agr.* 237. *t.* 4. *f.* 27, 28.

In pratis, pomariis. Fl. æſtate. Per.

fatua. 2. A. paniculata, calycibus trifloris, floſculis omnibus ariſtatis baſique piloſis. *Syſt. veget. p.* 122. *Schreb. gr. t.* 15. *Leers herb. p.* 42. *t.* 9. *f.* 4.

Avena triantha, locuſtis patulis villoſis. *Hall. hiſt. n.* 1495.

Gramen avenaceum locuſtis lanugine flaveſcentibus. *Tourn. inſt.* 525. *Scheuch. agr.* 239. *t.* 5. *f.* 1.

Inter ſegetes, ad margines agrorum, ad pagos. Fl. Majo, Iunio.

dubia. 3. A. paniculata, calycibus ſubtrifloris: floſculo infimo terminali ariſta recta; reliquis apice biariſtatis, dorſali ariſta reflexa. *Leers herb. p.* 41. *t.* 9. *f.* 3.

In agro vindobonenſi in ſterilibus montis *Galitzinberg.* Fl. Iunio.

flaveſcens. 4. A. panicula laxa, calybicus trifloris brevibus, floſculis omnibus ariſtatis. *Sp. pl. p.* 118. *Schreb. gr. t.* 9. *Curtis lond. Leers herb. p.* 43. *t.* 10. *f.* 5.

Avena triantha, locuſtis teretibus, calycina gluma altera minima, petiolo villoſo. *Hall. hiſt. n.* 1497.

In pratis, pomariis, & paſcuis alpinis. Fl. æſtate. Per.

ſeſqui-tertia. 5. A. paniculata, calycibus ſubtrifloris, floſculis omnibus ariſtatis, receptaculis barbatis. *Syſt. veget. p.* 122.

Gramen pratenſe, villoſum, panicula denſa, ex argenteo dilute ſpadicea, vel fuſca. *Scheuch. agr.* 220. *t.* 4. *f.* 17.

In pratis alpinis, & ſubalpinis. Fl. Iulio. Per.

Scheuchzeri. 6. A. paniculata, calycibus ſubquinquefloris, floſculis omnibus ariſtatis baſi pubescentibus.

Avena petiolis ramoſis, locuſtis teretibus, quinquefloris, baſi pubescentibus. *Hall. hiſt. n.* 1500. *Allion. pedem. tom.* 2. *p.* 255.

Gramen avenaceum, paniculatum, alpinum, locuſtis varicoloribus, majoribus, ariſtatis. *Scheuch. agr.* 231. *prodr. t.* 3.

In pascuis alpium Styriæ ſuperioris, Tyrolis. Fl. Iulio, Auguſto. Per.

pubescens. 7. A. ſubſpicata, calycibus ſubtrifloris, baſi piloſis, foliis planis pubescentibus. *Sp. pl. app. p.* 1665. *Leers herb. p.* 42. *t.* 9. *f.* 2.

Avena triantha, locuſtis teretibus erectis, petiolis ſericeis, gluma exteriori lacera. *Hall. hiſt. n.* 1498.

Gramen avenaceum, panicula purpureo-argentea, ſplendente. *Scheuch. agr.* 226.

In Auſtriæ, Pannoniæ pratis humidiusculis. Fl. Majo, Iunio. Per.

pratenſis. 8. A. ſubſpicata, calycibus quinquefloris. *Syſt. veget. p.* 122. *Leers herb. p.* 43. *t.* 9. *f.* 1.

Avena petiolis brevibus unifloris, locuſtis teretibus quinquefloris, baſi pubescentibus. *Hall. hiſt. n.* 1499.

In collibus apricis, pratis ſiccioribus. Fl. Iunio. Per.

fragilis. 9. A. ſpicata, calycibus quadrifloris floſculo longioribus. *Syſt. veget. p.* 122. *Schreb. gr. t.* 24. *f.* 3.

Gramen loliaceum lanuginoſum, ſpica fragili articulata, glumis piloſis, ariſtatum. *Scheuch. agr.* 32.

Gramen loliaceum lanuginoſum, ſpica fragili articulata, glumis glabris, ariſtatum. *ejusd.* 33.

In Iſtria. *I. Hoſt.* Fl. Iunio.

ARUNDO. Calyx bivalvis, uni- vel multiflorus. Corolla bivalvis; e valvularum baſi lanugo longitndine fere floris aſſurgit.

Donax. 1. A. calycibus trifloris, panicula diffuſa, culmo fruticoſo. *Syſt. veget. p.* 121. *Scop. carn. tom.* 1. *p.* 89.

Arundo caule lignoſo, geniculato, foliis latiſſimis, locuſtis trifloris. *Hall. hiſt. n.* 1516.

Arundo ſativa, quæ Donax Dioſcoridis. *Bauh. pin.* 17. *Scheuch. agr.* 159. *t.* 3. *f.* 14.

In Litorali in paluſtribus aquilejenſibus. In Iſtria in ſimilibus locis circa *Cepichi*, & inter Novacum & Piſinum, alibique.

phragmitis. 2. A. calycibus quinquefloris, panicula laxa. *Sp. pl. p.* 120. *Leers herb. p.* 44. *t.* 7. *f.* 1. *Scop. carn. tom.* 1. *p.* 89.

Arundo foliis ſecantibus, locuſtis trifloris, pappoſis, muticis. *Hall. hiſt. n.* 1515.

Arundo vulgaris, ſive Phragmitis Dioſcoridis. *Bauh. pin.* 17. *Scheuch. agr.* 161. *t.* 3. *f.* 14.

In paluſtribus, foſſis, lacubus, ad fluvios. Fl. Iulio, Auguſto. Per.

epigejos. 3. A. calycibus unifloris, panicula erecta, foliis ſubtus ſcabris, culmo ſimplici. *Retzii ſcand. p.* 27.

Gramen arundinaceum, paniculatum, montanum; panicula ſpadiceo-viridi, ſemine pappoſo. *Scheuch. agr.* 124.

In montanis, collibus, in arenoſis ad fluvios, alibique. Fl. Iulio, Auguſto. Per.

Medio valvulæ corollinæ dorſo adnaſcitur ariſta valvula ſua longior, & parum geniculata.

Calamagroſtis. 4. A. calycibus unifloris lævibus, corollis lanuginoſis, culmo ramoſo. *Syſt. veget. p.* 121.

Gramen arundinaceum, panicula molli, ſpadicea, majus. *Bauh. pin.* 7. *Scheuch. agr.* 122. *prodr. t.* 5.

In Styria in alpibus judenburgenſibus, præprimis locis humidiuſculis, & ad lacus. Fl. Auguſto. Per.

Ariſta recta valvulæ corollæ adnata valvula ſua vix longior.

Leerſii. 5. A. panicula incraſſata, calycibus unifloris ſcabris, petalo exteriore fiſſo ariſtato. *Hoffm. fl. p.* 40.

Agroſtis Calamagroſtis. *Leers herb. p.* 19.

In inſulis danubialibus. Fl. Iulio. Per.

LOLIUM. Receptaculum commune elongatum in ſpicam, flores diſtiche ſpicatos angulo culmi apprimens. Calycis gluma univalvis, rachi ita oppoſita, ut culmus valvulæ alterius vicem gerat.

perenne. 1. L. ſpica mutica: ſpiculis compreſſis multifloris. *Sp. pl. p.* 122. *Fl. dan. t.* 747. *Leers herb. p.* 46. *t.* 12. *f.* 1. *Scop. carn. tom.* 1. *p.* 90.

Lolium radice perenni, locuſtis octifloris, contiguis. *Hall. hiſt. n.* 1416.

Gramen loliaceum, anguſtiori folio & ſpica. *Bauh. pin.* 9. *Scheuch. agr.* 25.

In cultis, pratis, ad vias. Fl. æſtate. Per.

tenue. 2. L. ſpica mutica tereti, ſpiculis triflotis *Sp. pl. p.* 122.

Lolium perenne locuſtis diſſitis quadrifloris. *Hall. hiſt. n.* 1418.

Gramen loliaceum, foliis & ſpicis tenuiſſimis. *Vaill. paris.* 81.

In Iſtriæ herbidis. *I. Hoſt.* Fl. Iunio.

temulentum. 3. L. ſpica ariſtata, ſpiculis compreſſis multifloris. *Syſt. veget. p.* 124. *Fl. dan. t.* 160. *Leers herb. p.* 47. *t.* 12. *f.* 2. *Scop. carn. tom.* 1. *p.* 90.

Lolium annuum, locuſtis diſſitis. *Hall. hiſt. n.* 1420.

Gramen loliaceum ſpica longiore, ſeu Lolium Dioscoridis. *Bauh. pin.* 9. *Scheuch. agr.* 31.

Inter ſegetes. Fl. Iunio, Iulio.

Variat ſpica mutica.

ELYMUS. Receptaculum commune elongatum in ſpicam. Calycis gluma tetraphylla, diſtachya. Foliolis duobus ſingulæ ſpiculæ ſubjectis. Corolla bivalvis: valvula exterior acuminata, ariſtata.

europæus. 1. E. ſpica erecta: ſpiculis bifloris involucro æqualibus. *Syſt. veget. p.* 125.

Hordeum ſpica cylindrica, rigida, calycinis glumis ariſtatis. *Hall. hiſt. n.* 1537.

Gramen hordeaceum montanum ſpica ſtrigoſiori brevius ariſtata. *Scheuch. agr.* 16. *prodr. t.* 1.

In Auſtriæ, Styriæ ſylvis montanis & ſubalpinis. Fl. Iunio, Iulio. Per.

HORDEUM. Receptaculum commune elongatum in ſpicam. Calyx gluma hexaphylla, triflora: floribus ſeſſilibus; omnibus tribus hermaphroditis fertilibusque, vel duobus lateralibus maribus, intermedio hermaphrodito, et fertili.

murinum. 1. H. floſculis lateralibus maſculis ariſtatis, involucris intermediis ciliatis. *Sp. pl. p.* 126. *Fl. dan. t.* 629. *Curtis lond. Scop. carn. tom.* 1. *p.* 277.

Hordeum ſpica craſſa, longe ariſtata, calycinis glumis ariſtatis. *Hall. hiſt. n.* 1536.

Gramen hordeaceum minus & vulgare. *Bauh. pin.* 9. *Scheuch. agr.* 14.

In cultis, ruderatis, ad vias. Fl. æſtate.

TRITICUM. Receptaculum commune elongatum in ſpicam. Calyx gluma bivalvis, ſubtriflora. Corolla bivalvis: valvula exterior ventricoſa, obtuſa, in aliis ariſtata, in aliis mutica; valvula interior plana.

junceum. 1. T. calycibus truncatis quinquefloris, foliis involutis. *Sp. pl. p.* 128. *Fl. dan. t.* 916.

Triticum radice repente, culmo duro, foliis hirſutis, locuſtis quinquefloris. *Hall. hiſt. n.* 1428.

In collibus, ad vias agri vindobonenſis. Fl. Iunio, Iulio. Per.

repens. 2. T. calycibus quadrifloris ſubulatis acuminatis, foliis planis. *Syſt. veget. p.* 127. *Fl. dan. t.* 748. *Leers herb. p.* 44. *t.* 12. *f.* 3.

Triticum radicibus repentibus vaginoſis, foliis hirſutis, locuſtis quinquefloris. *Hall. hiſt. n.* 1426.

Gramen caninum arvenſe ſeu Gramen Dioſcoridis. *Bauh. pin.* 1. *Scheuch. agr.* 5.

In cultis, pratis, ruderatis, ad vias. Fl. Iulio. Per.

caninum. 3. T. calycibus ſubulatis ſubquadrifloris, floſculis ariſtatis. *Huds. angl. ed.* 2. 58. *Sibthorp. oxon. p.* 52.

Triticum radice fibroſa, foliis hirſutis, locuſtis quinquefloris ariſtatis. *Hall. hiſt. n.* 1429.

Elymus caninus. *Sp. pl. p.* 124. *Leers herb. p.* 45. *t.* 12. *f.* 4.

In ſylvis. Fl. Iulio, Auguſto. Per.

mariti-mum. 4. T. calycibus multifloris, floſculis mucronatis, ſpica ramoſa. *Sp. pl. p.* 128.

Gramen arvenſe, filicina, duriori panicula gracilius. *Barr. rar.* 1219. *Zann. iſt. p.* 133. *t.* 123, 126.

Gramen loliaceum, panicula ramoſa, maritimum. *Bauh. theatr.* 130. *Scheuch. agr.* 274. *t.* 6. *f.* 5.

In Litorali. Fl. Iunio. Per.

tenellum. 5. T. calycibus ſubquadrifloris, floſculis muticis acutis, foliis ſetaceis, *Sp. pl. p.* 127.

Triticum radice fibroſa, foliis anguſtiſſimis, locuſtis muticis, ovatis, obeſis. *Hall. hiſt. n.* 1430.

Feſtuca ſpica ſecunda recta, ſpiculis compreſſis ſubariſtatis. *Ger. prov. p.* 96.

Gramen loliaceum, foliis & ſpicis tenuiſſimis. *Moris. hiſt.* 3. *p.* 182. *ſect.* 8. *t.* 2. *f.* 3.

In Iſtriæ, Carniæ herbidis. *I. Hoſt.* Fl. Iunio, Iulio. Per.

Variat ſpiculis in eodem culmo muticis & ariſtatis.

TRIGYNIA.

MONTIA. Calyx diphyllus, triphyllusque. Corolla quinquepartita: laciniis tribus alternis minoribus tantum, quandoque omnibus quinque ſtaminiferis. Capſula unilocularis, bivalvis.

fontana. 1. Montia. *Sp. pl. p.* 129. *Curtis lond. Fl. dan. t.* 131. *Hall. hiſt. n.* 301.

In alpibus ſtyriacis, carinthiacis, tyrolenſibus ad rivulos, torrentes, & in aliis udis locis. Fl. æſtate.

HOLOSTEUM. Calyx pentaphyllus. Petala quinque. Capſula unilocularis, ſubcylindracea, apice dehiſcens.

umbellatum. 1. H. floribus umbellatis. *Sp. pl. p.* 130.

Alſine floribus umbellatis, petalis ſerratis. *Hall. hiſt. n.* 879.

In cultis, campis, ad vias, alibique. Fl. primo vere.

POLYCARPON. Calyx pentaphyllus. Petala quinque, minima, ovata. Capſula unilocularis, trivalvis.

tetraphyllum. 1. P. caule ramoſo proſtrato, foliis quaternis. *Syſt. veget. p.* 129. *Scop. carn. tom.* 1. *p.* 91.

Anthyllis maritima alſinefolia. *Bauh. pin.* 282. *Zann. iſt. p.* 138. *t.* 203.

In arenoſis, & in vineis Litoralis. Fl. æſtate.

TETRANDRIA.

MONOGYNIA.

GLOBULARIA. Calyx communis imbricatus; proprius tubulatus, inferus. Corolla propria labio ſuperiore bipartito: inferiore tripartito. Receptaculum paleaceum.

vulgaris. 1. G. caule herbaceo, foliis radicalibus tridentatis; caulinis lanceolatis. *Sp. pl. p.* 139. *Scop. carn. tom.* 1. *p.* 92.

Globularia foliis radicalibus emarginatis, nervo ariſtato, caulinis lanceolatis. *Hall. hiſt. n.* 218.

Globularia. *Clus. hiſt. p. VI.*

In collibus herbidis, in montoſis apricis. Fl. Majo, Iunio. Per.

cordifolia. 2. G. caule ſubnudo, foliis cuneiformibus tricuspidatis: intermedio minimo. *Sp. pl. p.* 139. *Iacqu. auſtr. t.* 245. *Scop. carn. tom.* 1. *p.* 93.

Globularia caule unifolio & unifloro, foliis emarginatis, nervo ariſtato. *Hall. hiſt. n.* 216.

Scabioſa decima ſive repens. *Clus. hiſt. p. V.*

In collibus apricis aſperis. Fl. Aprili, Majo. Per.

nudicaulis. 3. G. caule nudo, foliis integerrimis lanceolatis. *Sp. pl. p.* 140. *Iacqu. auſtr. t.* 230. *Scop. carn. tom.* 1. *p.* 93.

Globularia foliis ovatis oblongis, integerrimis, caule nudo unifloro. *Hall. hiſt. n.* 217.

In petroſis alpinis, & ſubalpinis. Fl. Iunio, Iulio. Per.

DIPSACUS. Calyx communis polyphyllus; proprius ſuperus. Receptaculum paleaceum.

ſylveſtris. 1. D. foliis integris, paleis rectis. *Syſt. veget. p.* 143. *Iacqu. auſtr. t.* 402. *Curtis lond. Fl. dan. t.* 965.

Dipſacus fullonum. *Scop. carn. tom.* 1. *p.* 97.

In ruderatis, ad ſepes, vias, & alibi. Fl. Iulio, Auguſto.

laciniatus. 2. D. foliis connatis ſinuatis. *Sp. pl. p.* 141. *Iacqu. auſtr. t.* 403.

In Auſtria ad foſſas, vias, ſepes humidas. Fl. a Iunio ad Auguſtum.

piloſus. 3. D. foliis petiolatis appendiculatis. *Sp. pl. p.* 141. *Iacqu. auſtr. t.* 248. *Curtis lond.*

Dipſacus foliis biauribus, capitulis hemiſphæricis. *Hall. hiſt. n.* 199.

In Auſtria, Pannonia, Carniolia in ſylvis, ad fluvios, ad ſepes, in apricis. Fl. Iunio, Iulio, Auguſto.

Folia ſunt ovata, vel lanceolata, utrinque piloſa, ſerrata, ſuperiora auriculata.

SCABIOSA. Calyx communis polyphyllus; proprius duplex, ſuperus. Receptaculum paleaceum ſeu nudum.

leucantha. 1. S. corollulis quadrifidis ſubæqualibus, ſquamis calycinis ovatis imbricatis, foliis pinnatifidis. *Sp. pl. p.* 142. *Scop. carn. tom.* 1. *p.* 95.

In aſperis Litoralis, Forijulii. Fl. æſtate. Per.

ſucciſa. 2. S. corollulis quadrifidis æqualibus, caule ſimplici, ramis approximatis, foliis lanceolato-ovatis. *Syſt. veget. p.* 144. *Curtis lond. Fl. dan. t.* 279. *Scop. carn. tom.* 1. *p.* 95.

Succiſa caule trifloro, floribus convexis, foliolis radicalibus ovatis, caulinis lanceolatis. *Hall. hiſt. n.* 201.

Succiſa ſeu Morſus Diaboli. *Matthiol. comm. p.* 623.

In pratis paluſtribus, paſcuis humidiusculis, alibique. Fl. Iulio, Auguſto. Per.

Variat foliis integerrimis, inciſis, & variter dentatis.

arvenſis. 3. S. corollis quadrifidis radiantibus, foliis pinnatifidis inciſis, caule hiſpido. *Syſt. veget. p.* 144. *Curtis lond. Fl. dan. t.* 447. *Scop. carn. tom.* I. *p.* 93.

Scabioſa foliis petiolatis, ovato-lanceolatis, dentatis, ſuperioribus ſemipinnatis. *Hall. hiſt. n.* 206.

La Scabieuſe des Prés. Regn. bot.

In arvis, pratis, ad vias. Fl. æſtate. Per.

Folia radicalia & caulina ima ſunt integra. Superiora pinnatifida: pinna impari maxima.

ſylvatica. 4. S. corollulis quadrifidis radiantibus, foliis omnibus indiviſis ovato-oblongis ſerratis, caule hispido. *Syſt. veget. p.* 144. *Iacqu. auſtr. t.* 362. *Scop. carn. tom.* I. *p.* 94.

Scabioſa caule hiſpido; foliis ovato-lanceolatis ſubhirſutis, inferioribus dentatis. *Hall. hiſt. n.* 204.

Scabioſa latifolia pannonica flore rubro ſeu tertia. *Clus. hiſt. p.* I.

Scabioſa latifolia pannonica purpuraſcente flore ſeu quarta. *Clus. hiſt. p.* II.

In ſylvis. Fl. æſtate. Per.

columbaria. 5. S. corollulis quinquefidis radiantibus, foliis radicalibus ovatis crenatis, caulinis pinnatis ſetaceis. *Sp. pl. p.* 143. *Scop. carn. tom.* I. *p.* 96. *Fl. dan. t.* 314.

Succiſa foliis imis ovatis, ſuperioribus pinnatis, pinnis ſemipinnatis, acutis, ciliis floſculorum longitudine. *Hall. hiſt. n.* 202.

Scabioſa quinta. *Clus. hiſt. p. II.*

In collibus herbidis, montoſis ſylvaticis. Fl. æſtate. Per.

gramini-folia. 6. S. corollulis quinquefidis, foliis lineari-lanceolatis integerrimis, caule herbaceo. *Sp. pl. p.* 145. *Scop. carn. tom.* 1. *p.* 96.

Succiſa foliis gramineis tomentoſis. *Hall. hiſt. n.* 203.

Scabioſa argentea anguſtifolia. *Bauh. pin.* 270. *prodr.* 127.

In Forojulio, Carnia, Tyroli calidiore. Fl. æſtate. Per.

ochro-leuca. 7. S. corollulis quinquefidis radiantibus, foliis bipinnatis linearibus. *Syſt. veget. p.* 146. *Iacqu. auſtr. t.* 439.

Scabioſa ſeptima. *Clus. hiſt. p. III.*

In collibus herbidis, ſiccis, aſperis, in montanis. Fl. æſtate. Per.

SHERARDIA. Corolla monopetala, infundibuliformis. Semina duo, tridentata.

arvenſis. 1. S. foliis omnibus verticillatis, floribus terminalibus. *Sp. pl. p.* 149. *Curtis lond. Fl. dan. t.* 439.

Sherardia foliis ſenis, lanceolatis, floribus ſeſſilibus umbellatis. *Hall. hiſt. n.* 734.

In arrvis. Fl. æſtate.

Tam caulis inferne, quam folia margine aculeis ſcabra. Pedunculi axillares, foliis duplo longiores, terminati bractea magna, multiflora.

ASPERULA. Corolla monopetala, infundibuliformis. Semina duo, globoſa.

odorata. 1. A. foliis octonis lanceolatis, florum faſciculis pedunculatis. *Sp. pl. p.* 150. *Curtis lond. Fl. dan. t.* 562.

Aſperula caule erecto, foliis octonis, petiolis ramoſis erectis, ſeminibus hirſutis. *Hall. hiſt. n.* 728.

Galium odoratum. *Scop. carn. tom.* 1. *p.* 105.

Aſperula odorato flore. *Clus. hiſt. p. CLXXV.*

Le Muguet des Bois, ou Hepatique etoilée. Regn. bot.

In ſylvis umbroſis. Fl. Aprili, Majo. Per.

Radix repens. Caulis erectus, tetragonus. Folia lanceolata, aculeato-ciliata. Flores albi, odori. Fructus, vertice excepto, aculeis hamatis ſcaber.

arvenſis. 2. A. foliis ſenis, floribus ſeſſilibus aggregatis terminalibus. *Sp. pl. p.* 150. *Scop. carn. tom.* 1. *p.* 106.

Aſperula foliis octonis, obtuſis, floribus ſeſſilibus congeſtis, bracteis ciliatis. *Hall. hiſt. n.* 733.

In arvis. Fl. Majo, Iunio.

Caulis erectus, ſcaber. Folia ſena, vel octona, lineari-lanceolata. Flores cærulei. Fructus glabri.

tinctoria. 3. A. foliis linearibus: inferioribus ſenis; intermediis quaternis, caule flaccido, floribus plerisque trifidis. *Sp. pl. p.* 150.

Aſperula caule flaccido, foliis linearibus quaternis, umbellulis petiolatis. *Hall. hiſt. n.* 729.

Galium tinctorium. *Scop. carn. tom.* 1. *p.* 101.

Galium album. *Tabern. hiſt.* 433. *t.* 733. *f.* 1.

In ſaxoſis montanis. Fl. Iunio, Iulio. Per.

cynan-chica. 4. A. foliis quaternis, linearibus: ſuperioribus oppoſitis, caule erecto, floribus quadrifidis. *Syſt. veget. p.* 149.

Aſperula caule firmo, ramoſo, foliis linearibus quaternis, ſupremis conjugatis. *Hall. hiſt. n.* 730.

Galium cynanchicum. *Scop. carn. tom.* 1. *p.* 101.

Galium album minus. *Tabern. hiſt.* 433. *t.* 733. *f.* 1.

In apricis. Fl. æſtate. Per.

GALIUM. Corolla monopetala, rotata; tubo nullo. Semina duo, ſubrotunda.

* *Fructu glabro.*

paluſtre. 1. G. foliis quaternis obovatis inæqualibus, caulibus diffuſis. *Sp. pl. p.* 153. *Fl. dan. t.* 423.

Galium caule radicato, diffuſo, foliis quaternis ovatis, obtuſis. *Hall. hiſt. n.* 719.

In paluſtribus, ad foſſas, rivulos, ſepes humidas. Fl. æſtate Per.

Caules diffuſi, tetragoni; angulis obtuſis, ſcabris. Folia quaterna lanceolato-ovata, margine ſcabra, inæqualia. Flores albi. Semen plerumque unicum, altero abortiente.

uliginoſum. 2. G. foliis ſenis lanceolatis retrorſum aculeato-ſerratis mucronatis rigidis, corollis fructu majoribus. *Sp. pl. p.* 153.

In uliginoſis. Fl. Majo, Iunio. Per.

Radix repens. Caulis erectus, tetragonus ſtriatus. Folia lineari-lanceolata, mucronata, margine aculeato-ſerrata, facie læte, dorſo pallide viridia.

ſcabrum. 3. G. foliis ſuboctonis ſcabris mucronatis, ramis floriferis ſubtrichotomis. *Syſt. veget. p.* 151. *Iacqu. auſtr. t.* 422.

Galium *aſperum*, foliis ſeptenis cauleque inferne pubeſcentibus, paniculæ ramis ſtrictis ſubtrichotomis. *Schreb. ſpicil. p.* 3.

In montoſis ſylvaticis, alpinis, & ſubalpinis. Fl. æſtate. Per.

Folia ſeptena vel octona, in ramis pauciora, linearia, aliqua ad apicem dilatata, utrinque ſulcata, mucronata, villis, ut caulis inferne, inſperſa, aſperaque. Flores valde odorati. Corolla alba, patentiſſima, quadrifida, rarius trifida; laciniis lanceolatis & acutis. Iacqu. l. c.

verum. 4. G. foliis octonis linearibus ſulcatis, ramis floriferis brevibus. *Sp. pl. p.* 155. *Scop. carn. tom.* I. *p.* 103.

Galium foliis octonis, linearibus, perangustis, racemis multifloris ſpicatis. *Hall. hiſt. n.* 710.

Le Caille-lait. Regn. bot.

In pratis, collibus, aggeribus. Fl. Iunio, Iulio. Per.

Caulis erectus, obſolete tetragonus. Folia caulina octona, denaque, ramea pauciora, linearia, facie ſulcata, margine revoluta, mucronata. Flores paniculati, flavi, valde odori.

Mollugo. 5. G. foliis octonis ovato-linearibus ſubſerratis patentiſſimis mucronatis, caule flaccido, ramis patentibus. *Sp. pl. p.* 155. *Fl. dan. t.* 455.

Galium foliis octonis, ellipticis, caule anguloſo. *Hall. hiſt. n.* 711.

Mollugo. *Clus. hiſt. p.* CLVI.

Ad ſepes, in ceſpitoſis. Fl. æſtate. Per.

Caulis ramoſiſſimus, debilis, tetragonus, glaber, ad genicula incraſſatus. Folia octona, glabra, margine aculeato-ciliata. Flores albi.

ſylvaticum. 6. G. foliis octonis lævibus, ſubtus ſcabris: floralibus binis, pedunculis capillaribus, caule lævi, *Sp. pl. p.* 155.

Galium foliis octonis, ellipticis, caule terete. *Hall. hiſt. n. 712.*

Galium glaucum. *Fl. dan. t.* 609.

In ſylvis. Fl. æſtate. Per.

Caulis erectus, ſubteres, geniculatus, glaber. Folia octona, quandoque plura, mucronata, margine & coſta aculeis ſcabra, ex glauco viridia. Flores albi.

auſtriacum. 7. G. foliis linearibus, glabris mucronatis, caulibus tetragonis diffuſis. *Iacqu. auſtr. t.* 80.

In ſaxoſis apricis Auſtriæ. Fl. Majo, Iunio. Per.

Caules tenues, nitidi & glaberrimi. Folia a quinis ad dena, linearia, glabra, plana, mucronata. Fructus obverſe ovati, parum incurvati, nigricantes. Iacqu. l. c.

glaucum. 8. G. foliis verticillatis linearibus, pedunculis dichotomis, caule lævi. *Syſt. veget. p.* 151. *Iacqu. auſtr. t.* 81.

In ſaxoſis apricis, ad ſepes, in cespitoſis. Fl. Iunio. Per.

Caules erecti, teretes. Folia ſublinearia, patentiſſima vel erecta, rarius deflexa, facie glaberrima & virentia, dorſo palentia, & aſperiuscula, mucronata, octona circiter, lateribus revoluta. Corollæ *albæ tubus campanulatus.* Iacqu. l. c.

rubrum. 9. G. foliis verticillatis linearibus patulis, pedunculis breviſſimis. *Sp. pl. p.* 156. *Scop. carn. tom.* 1. *p.* 103.

Galium rubro flore. *Clus. hiſt. p.* CLXXV.

In ſaxoſis Litoralis, Carnioliæ calidioris. Fl. Iulio, Auguſto. Per.

Caules & rami ſuperne tetragoni, pilis breviſſimis obſiti. Folia linearia, margine revoluta, ſcabra, mucronata. Pedunculi uniflori. Corollæ rubræ.

** *Fructu hiſpido.*

boreale. 10. G. foliis quaternis lanceolatis trinerviis glabris, caule erecto, ſeminibus hiſpidis. *Sp. pl. p.* 156. *Scop. carn. tom.* 1. *p.* 102. *Hall. hiſt. n.* 722.

In uliginoſis. Fl. Iunio, Iulio. Per.

Caulis tetragonus, pilis brevibus adpreſſis inſperſus. Folia quaterna, ovato-lanceolata, inæqualia, aculeato-ciliata, pleraque trinervia. Corollæ albæ.

rotundifolium. 11. G. foliis quaternis ovatis aculeato-ciliatis, ſeminibus hiſpidis. *Sp. pl. p.* 156. *Iacqu. auſtr. t.* 94. *Scop. carn. tom.* 1. *p.* 202.

Galium foliis quaternis, petiolis longiſſimis, brachiatis, ſeminibus hirſutis. *Hall. hiſt. n.* 727.

In ſylvis ſubalpinis. Fl. Iulio, Auguſto. Per.

Caules tetragoni, procumbentes. Folia quaterna, obverſe ovata, obtuſa cum exiguo mucrone, trinervia, ciliata. Corolla campanulato-rotata. Semina pilis uncinatis ſcabra. Iacqu. l. c.

Seminibus hispidis diverſiſſimum ab Aſperula lævigata, cui ſemina glabra.

Aparine. 12. G. foliis octonis lanceolatis carinatis ſcabris retrorſum aculeatis, geniculis villoſis, fructu hiſpido. *Sp. pl. p.* 157. *Fl. dan. t.* 495. *Scop. carn. tom.* 1. *p.* 105.

Galium caule ſerrato, foliis ſenis, linearibus lanceolatis, ſerratis, petiolis unifloris. *Hall. hiſt. n.* 723.

Le Grateron, ou Rieble. Regn. bot.

In cultis, ruderatis, ad ſepes, vias. Fl. æſtate. Per.

CRUCIANELLA. Corolla monopetala, infundibuliformis, tubo filiformi: limbo unguiculato. Calyx diphyllus. Semina duo, oblonga.

anguſtifolia. 1. C. erecta, foliis ſenis linearibus, floribus ſpicatis. *Sp. pl. p.* 157.

Rubia anguſtifolia ſpicata. *Bauh. pin.* 334. *prodr.* 145.

Tergeſti, in Iſtria veneta. *I. Hoſt.*

RUBIA. Corolla monopetala campanulata. Baccæ duæ monoſpermæ.

tinctorum. 1. R. foliis annuis, caule aculeato. *Syſt. veget. p.* 152. *Scop. carn. tom.* 1. *p.* 106.

Rubia foliis ſenis ellipticis, aſperis. *Hall. hiſt. n.* 708.

Rubia major, ſeu legitima. *Clus. hiſt. p. CLXXVII.*

La Garance. Regn. bot.

In Litorali in ceſpitoſis ad mare. Fl. Majo, Iunio. Per.

PLANTAGO. Calyx quadrifidus. Corolla quadrifida: limbo reflexo. Stamina longiſſima. Capſula bilocularis, circumſciſſa.

major. 1. P. foliis ovatis glabris, ſcapo tereti, ſpica floſculis imbricatis. *Sp. pl. p.* 163. *Curtis lond. Fl. dan. t.* 461. *Scop. carn. tom.* 1. *p.* 107.

Plantago foliis petiolatis, ovatis, glabris, ſpica cylindrica. *Hall. hiſt. n.* 660.

Le Grand Plantin. Regn. bot.

In cultis, ad vias. Fl. æſtate. Per.

media. 2. P. foliis ovato-lanceolatis pubeſcentibus, ſpica cylindrica, ſcapo tereti. *Sp. pl. p.* 163. *Fl. dan. t.* 581. *Scop. carn. tom.* 1. *p.* 108.

Plantago foliis ſubhirſutis, ellipticis, ſpica cylindrica denſa. *Hall. hiſt. n.* 659.

Plantago major incana. *Cluſ. hiſt. p. CIX.*

Plantago major hirſuta, media a nonnullis cognominata. *Bauh. hiſt.* 3. *p.* 504.

In pratis, paſcuis ſterilibus, & alibi. Fl. æſtate. Per.

lanceolata. 3. P. foliis lanceolatis, ſpica ſubovata nuda, ſcapo angulató. *Sp. pl. p.* 164. *Curtis lond. Fl. dan. t.* 437. *Scop. carn. tom.* 1. *p.* 108.

Plantago foliis lanceolatis, quinquenerviis, ſcapo nudo, ſpica ovata. *Hall. hiſt. n.* 656.

Habitat & floret cum præc. Per.

alpina. 4. P. foliis linearibus planis, ſcapo tereti hirſuto, ſpica oblonga erecta. *Sp. pl. p.* 165. *Iacqu. hort. tom.* 2. *p.* 58. *t.* 125.

Plantago foliis linearibus, planis, ſpica cylindrica. *Hall. hiſt. n.* 657.

In Carniolia in monte *Nanas. Zois.* In Carniæ alpibus. *I. Hoſt.* Fl. Iunio, Iulio. Per.

maritima. 5. P. foliis ſemicylindraceis integerrimis baſi lanatis, ſcapo tereti. *Sp. pl. p.* 165. *Fl. dan. t.* 243.

In Auſtria, Pannonia, Litorali locis uliginoſis, paluſtribus, alibique. Fl. æſtate. Per.

Variat foliis denticulatis.

ſubulata. 6. P. foliis ſubulatis triquetris ſtriatis ſcabris, ſcapo tereti. *Sp. pl. p.* 166. *Wulfen in Iacqu. collect. tom.* 1. *p.* 204. *t.* 10.

Holoſteum ſtrictiſſimo folio, minus. *Bauh. pin.* 191.

In Litorali in locis aſperis, herbidis. Fl. Iunio, Iulio. Per.

Coronopus. 7. foliis linearibus dentatis, ſcapo tereti. *Sp. pl. p.* 166.

Plantago foliis ſubhirſutis, ſemipinnatis, pinnis raris, lanceolatis. *Hall. hiſt. n.* 658.

Coronopus hortenſis. *Bauh. pin.* 190.

Herba ſtella, ſive cornu cervinum. *Dod. pempt.* 169. *Zann. iſt. p.* 74. *t.* 207.

In agri fluminenſis herbidis, uliginoſis. Fl. æſtate.

Pſyllium. 8. P. caule ramoſo herbaceo, foliis ſubdentatis recurvatis, capitulis aphyllis. *Sp. pl. p.* 167.

Plantago caulibus erectis, herbaceis, foliis linearibus, patulis, capitulis ovatis hirſutis. *Hall. hiſt. n.* 661.

L' Herbe aux Puces. Regn. bot.

In Auſtria in collibus agri vindobonenſis prope *Weinhaus*. Fl. æſtate.

SANGUISORBA. Calyx tetraphyllus. Germen inter calycem corollamque.

officinalis. 1. S. ſpicis ovatis. *Sp. pl. p.* 169. *Fl. dan. t.* 97.

Pimpinella tetraſtemon, ſpica brevi. *Hall. hiſt. n.* 705.

Pimpinella italica major. Sanguiſorba. *Clus. hiſt. p. CXCVII.*

In pratis humidis, præcipue paluſtribus. Fl. Iulio, Auguſto. Per.

Tota planta glabra. Folia pinnata; foliolis cordatis, ſerratis. Stipulæ duæ reniformes ſerratæ ſupra baſin petioli communis. Stipulæ vero foliolorum ad baſin petiolorum.

EPIMEDIUM. Calyx tetraphyllus, caducus. Corolla tetrapetala. Nectaria quatuor, cyathiformia, petalis incumbentia. Siliqua.

alpinum. 1. Epimedium. *Sp. pl. p.* 171. *Scop. carn. tom.* 1. *p.* 111. *Dod. pempt.* 599.

In Forojulio in ſylvis circa Goritiam. *Scop.* Ego in Croatia circa *Raunagora.* Fl. Aprili, Majo. Per.

CORNUS. Involucrum tetraphyllum ſæpius. Corolla quadripetala, ſupera. Drupa nucleo biloculari.

mascula. 1. C. arborea, umbellis involucrum æquantibus. *Sp. pl. p.* 171. *Scop. carn. tom.* 1. *p.* 112. *Hall. hiſt. n.* 815. *L'Heritier corn. n.* 4.

Cornus. *Clus hiſt. p.* 12.

Le Cornouiller. Regn. bot.

In ſylvis, collibus, ad ſepes. Fl. primo vere.

Liburniæ quidam ruricolæ, fructus conquaſſatos ſemimaturos maturis immixtos, cum ſufficienti quantitate aquæ fermentationi exponunt, peractaque fermentatione acquirunt potum, urente ſirio, palato gratum, & ſalubrem.

ſanguinea. 2. C. arborea, cymis nudis, ramis rectis. *Syſt. veget. p.* 159. *Fl. dan. t.* 481. *Scop. carn. tom.* 1. *p.* 112. *Hall. hiſt. n.* 816.

Cornus ramis rectis, foliis ovatis concoloribus, cymis depreſſis. *L'Héritier corn. n.* 5.

In ſylvis, ad ſepes, & alibi. Fl. Majo.

In Iſtria fructus maturos cum addito pauxillo cinerum coquunt, ac ex coctis oleum exprimunt ad fovendas lampades aptiſſimum.

ISNARDIA. Calyx quadrifidus. Corolla nulla. Capſula quadrilocularis, cincta calyce.

paluſtris. 1. Isnardia. *Sp. pl. p.* 175. *Wulfen in Iacqu. collect. tom.* 1. *p.* 270.

Alſine paluſtris rotundifolia repens, foliis portulacæ pinguibus. *Lind. alſat.* 114. *t.* 2.

In Carinthia prope Clagenfurtum in uliginoſis ſylvarum montis Calvariæ; in paluſtribus zigulnenſibus; & in turfoſis ad lacum *Wertenſee*. In ſimilibus locis agri goritienſis in Forojulio. *Wulfen*. In Carnia in aquis ſtagnantibus non procul a monte *Croſtis*. *I. Hoſt*. Fl. Iulio, Auguſto.

TRAPA. Calyx quadripartitus. Corolla tetrapetala. Nux ſpinis quatuor, oppoſitis cincta, quæ calycis laciniæ fuere.

natans. 1. T. nucibus quadricornibus: ſpinis patentibus. *Syſt. veget. p.* 162.

Trapa foliis natantibus, petiolis ventricoſis. *Hall. hiſt. n.* 527.

La Mâcre, ou Chataigne d'Eau. Regn. bot.

In Auſtria in aquis ſtagnantibus prope Neoſtadium. In Carinthia in lacu non procul a *Dahlenſtein*. Fl. Iulio, Auguſto.

ELÆAGNUS. Calyx quadrifidus, campanulatus, ſuperus. Corolla nulla. Drupa.

anguſtifolia. 1. E. inermis foliis lanceolatis. *Sp. pl. p.* 176. *Pallas roſſic. t.* 4.

Olea ſylveſtris, folio molli incano. *Bauh. pin.* 472.

Ad litus maris adriatici in inſulis venetis, copioſiſſime in inſula Auxerio, vulgo *Iſola di Oſero*. *I. Hoſt*. Fl. Majo. Arb.

Rami plerumque ſpinis armati ſunt in ſylveſtri arbuſcula.

CAMPHOROSMA. Calyx urceolatus: dentibus duobus oppoſitis, alternisque minimis. Corolla nulla. Capſula monoſperma.

acuta. 1. C. foliis ſubulatis rigidis glabris. *Sp. pl. p.* 178.

Camphoratæ congener. *Bauh. pin.* 486.

Anthyllis altera italorum. *Lob. ic.* 404. *Zann. iſt. p.* 65. *t.* 276.

In arenoſis Litoralis. Fl. Iunio, Iulio. Per.

ALCHEMILLA. Calyx octofidus. Corolla nulla. Stylus ad baſin germinis inſertus.

vulgaris. 1. A. foliis lobatis. *Sp. pl. p.* 178. *Fl. dan. t.* 693. *Scop. carn. tom.* 1. *p.* 115. *Clus. hiſt. p.* CVIII.

Alchemilla foliis ſerratis, palmatis, ſeminovemfidis. *Hall. hiſt. n.* 1566.

Le Pied de Lion. Regn. bot.

In pratis. Fl. æſtate. Per.

alpina. 2. A. foliis digitatis ſerratis. *Sp. pl. p.* 179. *Fl. dan. t.* 49.

Alchemilla foliis ſericeis digitatis, apice dentatis. *Hall. hiſt. n.* 1567.

Heptaphyllon. *Clus. hiſt. p.* CVIII.

Tormentilla alpina foliis ſericeis. *Bauh. pin.* 326.

In alpibus tyrolenſibus, carinthiacis, carniolicis. Fl. Iulio, Auguſto. Per.

Aphanes. 3. A. foliis trilobatis, floribus monandris. *Retzii ſcand. p.* 37.

Alchemilla hirſuta, foliis trilobatis, lobis bipartitis & tripartitis. *Hall. hiſt. n.* 1569.

Alchemilla arvenſis. *Scop. carn. tom.* 1. *p.* 115.

Aphanes arvenſis. *Sp. pl. p.* 179. *Schkukr hand. t.* 26.

In arvis Auſtriæ, Carnioliæ, Iſtriæ. Fl. æſtate.

DIGYNIA.

CUSCUTA. Calyx quadri- ſeu quinquefidus. Corolla monopetala, quadri- ſeu quinquefida. Stamina quatuor vel quinque. Capſula bilocularis.

europæa. 1. C. floribus ſeſſilibus. *Sp. pl. p.* 180. *Fl. dan. t.* 199. *Scop. carn. tom.* 1. *p.* 116. *Hall. hiſt. n.* 654.

Cuscute. Regn. bot.

In pratis, collibus, cultis, ad ſepes, alibique. Fl. æſtate.

E ſemine oritur caulis volubilis, ex internodiis radiculas papillares, ſubſecuriformes emittens, quibus plantis adſigitur; ubi autem pars hujus ſuperior ita plantæ vicinæ adhæſit, ut ex illa nutrimentum haurire valeat; tunc pars inferior cum radice perit. Si autem nulla ſtirps in vicinia adſit, perit tota. Folia nulla. Squamula unica ovata ſub ſingulo glomere florum, ubi ſæpe ramum educit. Nectária in primariis floribus fere nulla, in ſeriorihus ſemibifida: laciniis iterum bifidis, incurvis. Linn. Leers. Gouan.

TETRAGYNIA.

ILEX. Calyx quadridentatus. Corolla rotata. Stylus nullus. Bacca quadriſperma.

Aquifolium. 1. I. foliis ovatis acutis ſpinoſis. *Sp. pl. p.* 181. *Fl. dan. t.* 508. *Hall. hiſt. n.* 667.

Aquifolium Ilex. *Scop. carn. tom.* 1. *p.* 116.

Ilex aculeata baccifera folio ſinuato. *Bauh. pin.* 425.

In Litorali in ſaxoſis agri fluminenſis. In Carniolia in ſylvis circa Idriam. In Forojulio circa Goritiam. In Auſtria inferiore *zu Heiligenkreutz im Waldthale.* Fl. Majo, Iunio.

Iunior & pumila arbuscula folia habet ſinuata, undulata, ſpinoſa. Adulta & elata plana, inermia.

POTAMOGETON. Calyx nullus. Corolla tetrapetala. Germina quatuor. Stylus nullus.

natans. 1. P. foliis oblongo-ovatis petiolatis natantibus. *Sp. pl. p.* 182. *I. Miller illuſtr. ic. Fl. dan. t.* 165.

Potamogeton foliis firmis, petiolatis ovatis, natantibus. *Hall. hiſt. n.* 843.

In fluviis, & aquis ſtagnantibus. Fl. Iunio, Iulio. Per.

perfoliatum. 2. P. foliis cordatis amplexicaulibus. *Sp. pl. p.* 182. *Fl. dan. t.* 196. *Scop. carn. tom.* 1. *p.* 118. *Hall. hiſt. n.* 845.

In lacubus, fluviis. Fl. Iunio. Per.

denſum. 3. P. foliis ovatis acuminatis oppoſitis confertis, caulibus dichotomis, ſpica quadriflora. *Sp. pl. p.* 182.

Potamogeton caule dichotomo, foliis conjugatis, ellipticis, complicatis, imbricatis. *Hall. hiſt. n.* 849.

Tribuli aquatici minoris ſpecies altera. *Clus. hiſt. p.* CCLII.

In rivulis, lacubus, foſſis, fluviis. Fl. Iulio, Auguſto. Per.

lucens. 4. P. foliis lanceolatis planis in petiolos deſinentibus. *Sp. pl. p.* 183. *Fl. dan. t.* 195. *Scop. carn. tom.* 1. *p.* 118.

Potamogeton foliis tenuibus, longiſſime lanceolatis. *Hall. hiſt. n.* 847.

Potamogeton foliis anguſtis ſplendentibus. *Bauh. pin.* 193.

In fluviis, foſſis, lacubus. Fl. Iunio, Iulio. Per.

criſpum. 5. P. foliis lanceolatis alternis oppoſitisve undulatis ſerratis. *Syſt. veget. p.* 169. *Curtis lond. Scop. carn. tom.* I. *p.* 118. *Hall. hiſt. n.* 848.

Tribulus aquaticus minor. *Clus. hiſt. p.* CCLII.

Potamogeton foliis criſpis ſeu lactuca ranarum ſarmentis planis. *Bauh. pin.* 193.

Cum præcedente. Fl. Iunio. Per.

Folia ſeſſilia, lanceolata, obtuſa, valde criſpa, nitida, ſerrulata, inferiora alterna, ſuperiora oppoſita.

compreſſum. 6. P. foliis linearibus obtuſis, caule compreſſo. *Sp. pl. p.* 183. *Fl. dan. t.* 203.

Potamogeton caulibus compreſſis, foliis gramineis lanceolatis, ſpicis paucifloris. *Hall. hiſt. n.* 851.

In aquis ſtagnantibus. Fl. Iunio. Per.

gramineum. 7. P. foliis lineari-lanceolatis alternis ſeſſilibus ſtipula latioribus. *Sp. pl. p.* 184. *Fl. dan. t.* 222.

Potamogeton foliis gramineis, petiolis erectis, vaginis laxis numeroſiſſimis. *Hall. hiſt. n.* 850.

In foſſis, rivulis. Fl. Iunio. Per.

pectinatum. 8. P. foliis ſetaceis parallelis approximatis diſtichis. *Sp. pl. p.* 183.

Potamogeton millefolium, ſeu foliis gramineis ramoſum. *Raj. ſyn.* 150.

Petamogeton gramineum ramoſum. *Bauh. pin.* 193. *prodr.* 101.

In fluviis, lacubus, foſſis. Fl. Iunio. Per.

RUPPIA. Calyx nullus. Corolla nulla. Semina quatuor, pedicellata.

mariti-ma. 1. Ruppia. *Sp. pl. p.* 184. *Fl. dan. t.* 364.

Buccaferrea maritima, foliis acutiſſimis. *Mich. gen. p.* 72. *t.* 35.

Tergeſti in foſſis *alle Saule. I. Hoſt.*

SAGINA. Calyx tetraphyllus. Petala quatuor. Capſula unilocularis, quadrivalvis, polyſperma.

procum-bens. 1. S. ramis procumbentibus. *Syſt. veget. p.* 169. *Curtis lond.*

In arenoſis, uliginoſis, paſcuis, montoſis ſylvaticis. Fl. æſtate.

Radix annua: in locis umbroſis, humidis plerumque perennis. Caules in umbroſis procumbentes, in aridis erecti, geniculati, ramoſi. Folia oppoſita, patentia, connata, ſubulata, mucronata, ad ramorum finem faſciculata. Pedunculi axillares, plerumque uniflori, alterni, foliis longiores. Petala quatuor, calyce triplo breviora, alba, patentia, ſæpe nulla. Curtis l. c.

PENTANDRIA.

MONOGYNIA.

HELIOTROPIUM. Corolla hypocrateriformis, quinquefida, interjectis dentibus: faux nuda.

europaeum. 1. H. foliis ovatis integerrimis tomentoſis, ſpicis conjugatis. *Sp. pl. p.* 187. *Iacqu. auſtr. t.* 207. *Scop. carn. tom.* 1. *p.* 120.

Heliotropium foliis ovatis, petiolatis, ſpicis inferioribus ſimplicibus, ſupremis gemellis. *Hall. hiſt. n.* 593.

Heliotropium majus. *Clus. hiſt. p. XLVI.*

In Auſtriae, Pannoniae, Croatiae, Forijulii arvis, alibique. Fl. aeſtate.

Inter lacinias corollae invenitur denticulus unus, vel alter: ſaepe nullus.

MYOSOTIS. Corolla hypocrateriformis, quinquefida, emarginata: fauce clauſa fornicibus.

ſcorpioides. 1. M. ſeminibus nudis: foliorum apicibus calloſis. *Sp. pl. p.* 188. *Fl. dan. t.* 583. *Curtis lond.*

In arvis, campis, collibus, ſylvis, locis paluſtribus, aquoſis, in alpinis, & ſubalpinis. Fl. aeſtate.

In apricis, aridis creſcens annua eſt; perennis contra in humidis, umbroſis, alpinisque habitans. Prout, pro diverſa loci natura, ac elevatione, vitae duratione variat, ſic variat quoque caule ſimplici, vel ramoſo; calyce, foliisque glabris, aut plus, minusve hirſutis; corollis amplioribus, & minoribus. Cl. Linnaeus fecit inde tres varietates. Cl. Allionius in flora pedemontana tres ſpecies. Varietatemque in arvis, aridisque creſcentem, Myoſotim arvenſem: in aquoſis, paluſtrem; nanam, quae celſiſſima alpium juga incolit, vocaverat. At in Auctario ſuo ad dictam florum de nana loquens dicit. „Quamvis magni-

tudine florum, & caulis ſimplicitate diſtincta ſpecies videatur, erit forte tamen habenda pro varietate paluſtris; cum ſucceſſivo gradu, prout ex alpibus ad montes deſcendit, in paluſtrem abeat, atque etiam in ſummis alpibus occurrat eadem planta humilis quidem, ſed ramoſa, minus villoſa, & minoribus floribus donata. Adnotavi etiam arvenſem in agris humidis naſcentem ad paluſtrem accedere, ita ut dicendum ſit, non male a ſummo viro Linnæo in unam ſpeciem eas fuiſſe conjunctas."

Lappula. 2. M. ſeminibus aculeis glochidibus, foliis lanceolatis piloſis. *Sp. pl. p.* 189. *Fl. dan. t.* 692.

Cynogloſſum glandulis faucium glabris, *Hall. hiſt. n.* 589.

Cynogloſſum Lappula. *Scop. carn. tom.* I. *p.* 125.

In arvis, ruderatis, vetuſtis hortorum muris, ad vias. Fl. æſtate.

LITHOSPERMUM. Corolla infundibuliformis fauce perforata, nuda. Calyx quinquepartitus.

officinale. 1. L. ſeminibus lævibus, corollis calycem vix ſuperantibus, foliis lanceolatis. *Sp. pl. p.* 189. *Scop. carn. tom.* I. *p.* 121.

Lithoſpermum caule erecto, ramoſiſſimo, floribus calycem vix ſuperantibus. *Hall. hiſt. n.* 595.

Le Gremil. Regn. bot.

In ruderatis, ſylvis, ad fluvios. Fl. æſtate. Per.

arvenſe. 2. L. ſeminibus rugoſis, corollis vix calycem ſuperantibus. *Sp. pl. p.* 190. *Fl. dan. t.* 456. *Scop. carn. tom.* I. *p.* 122.

Heliotropium foliis lingulatis, floribus tubuloſis, *Hall. hiſt. n.* 594.

In arvis, alibique. Fl. primo vere.

purpuro-cœruleum. 3. L. ſeminibus lævibus, corollis calycem multoties ſuperantibus. *Sp. pl. p.* 190. *Iacqu. auſtr. t.* 14. *Scop. carn. tom.* 1. *p.* 122.

Lithoſpermum caule procumbente radicato, ramis floralibus erectis. *Hall. hiſt. n.* 596.

Lithoſpermum repens majus. *Clus. hiſt. p.* CLXIII.

In ſylvis montoſis, collibus fruticoſis. Fl. Majo, Iunio. Per.

ANCHUSA. Corolla infundibuliformis, fauce clauſa ſquamulis quinque convexis. Semina obtuſa, gibba, in ſinu calycis.

officinalis. 1. A. foliis lanceolatis ſtrigoſis, ſpicis ſecundis imbricatis, calycibus quinquepartitis. *Ait. Kew. tom.* 1. *p.* 178. *Sp. pl. p.* 191. *Fl. dan. t.* 572. *Scop. carn. tom.* 1. *p.* 123.

Bugloſſum foliis linguiformibus aſperis, ſpicis ſupremis gemellis. *Hall. hiſt. n.* 599.

La Bugloſſe. Regn. bot.

Ad vias, agrorum margines, in ruderatis, & alibi. Fl. æſtate. Per.

italica. 2. A. foliis undulatis, ſummis acuminatis, ſquamulis faucis penicilliformibus. *Syſt. veget. p.* 186. *Trew. rar. dec.* 2. *p.* 14. *t.* 28.

In herbidis, ad vias, vineas agri tergeſtini. *I. Hoſt.* Fl. æſtate. Per.

CYNOGLOSSUM. Corolla infundibuliformis. Faux clauſa ſquamulis quinque, convexis. Pericarpium nullum, ſed arilli quatuor, monoſpermi, ſcabri, interiore tantum latere ſtylo affixi.

officinale. 1. C. ſtaminibus corolla brevioribus, foliis lato-lanceolatis baſi attenuatis tomentoſis ſeſſilibus, la-

ciniis calycinis oblongis. *Ait. Kew. tom.* 1. *p.* 179. *Sp. pl. p.* 192. *Curtis lond. Scop. carn. tom.* 1. *p.* 124.

La Cynogloſſe. Regn. bot.

Ad vias, fluvios, in ruderatis, pratis, ſylvis. Fl. æſtate.

cheirifolium. 2. C. corollis calyce duplo longioribus, foliis lanceolatis. *Sp. pl. p.* 193. *Scop. carn. t.* 1. *p.* 125.

Cynogloſſum creticum primum. *Clus. hiſt. p.* CLXII.

Tergeſti *dietro il Lazzaretto ſporco*, ad vias. Fl. Majo, Iunio.

Omphalodes. 3. C. repens, foliis radicalibus cordatis. *Sp. pl. p.* 193. *Curtis magaz.* 7. *Scop. carn. tom.* 1. *p.* 123. *t.* 3.

Symphytum minus, boraginis facie. *Bauh. pin.* 209.

In ſylvis Carnioliæ, Croatiæ. Fl. Aprili, Majo. Per.

PULMONARIA. Calyx prismatico-pentagonus. Corolla infundibuliformis, fauce pervia.

anguſtifolia. 1. P. foliis hirſutis: caulinis oblongo-lanceolatis amplexicaulibus; radicalibus ellipticis. *Ait. Kew. tom.* 1. *p.* 181. *Sp. pl. p.* 194. *Fl. dan. t.* 483. *Hall. hiſt. n.* 598.

Pulmonaria tertia auſtriaca. *Clus. hiſt. p.* CLXIX.

Pulmonaria quinta pannonica. *Clus. hiſt. p.* CLXX.

In ſylvis, & fruticetis. Fl. Aprili, Majo. Per.

officinalis. 2. P. foliis hirſutis: caulinis ovato-oblongis; radicalibus ſubcordatis. *Ait. Kew. tom.* 1. *p.* 181. *Sp. pl. p.* 194. *Hall. hiſt. n.* 597.

Pulmonaria vulgaris maculoſo folio. *Clus. hiſt. p.* CLXIX.

Pulmonaria folio non maculoſo. *Clus. hiſt. p.* CLXIX.

La Pulmonaire. Regn. bot.

In pratis, ſylvis, alibique. Fl. Martio, Aprili. Per.

In utraque ſpecie corollæ faux munita eſt villis conniventibus.

SYMPHYTUM. Corollæ limbus tubulato-ventricoſus: fauce munita radiis quinque, ſubulatis, limbo brevioribus, in conum conniventibus.

officinale. 1. S. foliis ovato-lanceolatis decurrentibus. *Sp. pl. p.* 195. *Curtis lond. Fl. dan. t.* 664. *Scop. carn. tom.* 1. *p.* 126. *Hall. hiſt. n.* 600.

Symphytum tuberoſum majus. *Clus. hiſt. p.* CLXVI.

La grande Conſoude. Regn. bot.

In pratis humidis, pomariis, ad fluvios, rivulos. Fl. Iunio. Per.

tuberoſum. 2. S. foliis ſemidecurrentibus: ſummis oppoſitis. *Syſt. veget. p.* 187. *Iacqu. auſtr. t.* 225. *obs. bot.* 3. *p.* 12. *t.* 63.

In ſylvaticis, fruticoſis Auſtriæ, Pannoniæ. Fl. Aprili, Majo. Per.

Radix eſt verum tuber, in juniori planta ſimplex, in annoſa ramoſum, breve, nunquam in fibras craſſas longasque, ut radix Symphyti officinalis, extenſum. Corollæ ſunt ſemper ochroleucæ, dum in officinali purpureæ, vel, quod rarum, albæ. Iacqu. l. c.

CERINTHE. Calyx pentaphyllus, foliolis inæqualibus. Corollæ limbus tubulato-ventricoſus: fauce pervia. Antheræ ſerrulatæ, baſi biſetæ. Germen didymum. Nuces duæ, biloculares: loculis monoſpermis. *Ehrh. beytr. t. 5. p. 44.*

major. 1. C. foliis amplexicaulibus, fructibus geminis, corollis obtuſiusculis patulis. *Sp. pl. p.* 195.

Cerinthe foliis amplexicaulibus, floris denticulis breviſſimis, revolutis. *Hall. hiſt. n.* 602.

Cerinthe quorundam major verſicolore flore. *Clus. hiſt. p.* CLXVII.

In Carnia. *I. Hoſt.* Fl. Iunio, Iulio.

minor. 2. C. foliis amplexicaulibus integris, fructibus geminis, corollis acutis clauſis. *Sp. pl. p.* 196. *Iacqu. auſtr. t.* 124.

Cerinthe quorundam minor, flavo flore. *Clus. hiſt. p.* CLXVIII.

In agris, ſylvis. Fl. æſtate.

Folia radicalia in utraque ſpecie in petiolum attenuata & obverſe ovata, caulina amplexicaulia. Omnia *obtuſa, integra, modo maculis pallidis, modo nullis adſperſa. In minore ſubtus glauca, ſupra magis virentia. In majore per ætatem utrinque e viridi cæruleſcentia.* Co*rolla in minore flava, baſi, apiceque attenuata. In majore e rubro & purpuraſcente varia, raro flava, baſi attenuata, apice patula.*

ONOSMA. Corolla campanulata: fauce pervia. Semina quatuor.

echioides. 1. O. foliis lanceolatis hiſpidis, fructibus erectis. *Sp. pl. p.* 196. *Iacqu. auſtr. t.* 295.

Symphytum foliis lingulatis hiſpidis. *Hall. hiſt. n.* 601.

Cerinthe echioides. *Scop. carn. tom.* 1. *p.* 128.

Anchuſa exalbido flore. *Clus. hiſt. p.* CLXV.

In collibus apricis, ſaxoſis, aſperis Auſtriæ, Pannoniæ, Croatiæ, Litoralis, Forijulii. Fl. æſtate.

BORAGO. Corollæ limbus quinquepartitus, rotatus: fauce coronata prominentiis quinque, emarginatis.

officinalis. 1. B. foliis omnibus alternis, calycibus patentibus. *Sp. pl. p.* 197. *Plenk ic. t.* 77.

Borago foliis aſperis lanceolatis, palis florum duplicatis. *Hall. hiſt. n.* 607.

Occurrit ad hortos. Fl. æſtate.

ASPERUGO. Corolla infundibuliformis: fauce clauſa ſquamulis quinque. Calyx fructus compreſſus: lamellis plano parallelis, ſinuatis.

procumbens. 1. A. calycibus fructus compreſſis. *Sp. pl. p.* 198. *Fl. dan. t.* 552. *Scop. carn. tom.* 1. *p.* 129. *Hall. hiſt. n.* 606.

Bugloſſum ſylveſtre, caulibus procumbentibus. *Bauh. pin.* 257.

In ruderatis, ad ſepes, vias. Fl. Majo, Iunio.

LYCOPSIS. Corolla tubo incurvato.

pulla. 1. L. foliis integerrimis, caule erecto, calycibus fructescentibus inflatis pendulis. *Sp. pl. p.* 198. *Iacqu. auſtr. t.* 188.

Echium pullo flore. *Clus. hiſt. p.* CLXIV.

In Auſtria ad vias, agrorum margines, in arenoſis, ſterilibus. Fl. Majo, Iunio.

arvenſis. 2. L. foliis lanceolatis hiſpidis, calycibus floreſcentibus erectis. *Sp. pl. p.* 199. *Fl. dan. t.* 435. *Curtis lond.*

Lycopſis foliis aſperrimis, undulatis, ſerratis, linguiformibus. *Hall. hiſt. n.* 605.

In Auſtria in agro vindobonenſi circa *Weinhaus.* In Tyroli prope *Brunegen.* Fl. æſtate.

ECHIUM. Corolla campanulata. Limbus quinquefidus obtuſus: laciniis ſæpius inæqualibus; ſuperioribus duabus longioribus, infima minore, reflexa. Faux nuda.

italicum. 1. E. caule erecto piloſo, ſpicis hirſutis, corollis ſubæqualibus, ſtaminibus longiſſimis. *Sp. pl. p.* 200.

Echium foliis hirtis, lingulatis, calycibus pungentibus, ſpicis laxis, paniculatis. *Hall. hiſt. n.* 604.

Echium altiſſimum. *Iacqu. auſtr. app. t.* 16.

In comitatu Hungariæ nitrienſi, qua Poſonio Chemnitium itur in locis campeſtribus, ſiccis. In Auſtria prope *Schwæchat.* Fl. Iulio.

rubrum. 2. E. caule ex racemulis brevibus longe ſpicato, corollis ſubæqualibus, foliis hiſpidis. *Syſt. veget. p.* 190. *Iacqu. auſtr. app. t.* 3.

Echium rubro flore. *Clus. hiſt. p.* CLXIV.

In Auſtria in pratis, aliiſque herbidis locis circa *Ebergäſſing.* In Pannonia prope Sopronium, alibique. Fl. Iunio.

vulgare. 3. E. caule tuberculato hiſpido, foliis caulinis lanceolatis hiſpidis, floribus ſpicatis lateralibus. *Sp. pl. p.* 200. *Fl. dan. t.* 445. *Scop. carn. tom.* I. *p.* 129. *Clus. hiſt. p.* CLXIII.

Echium alis ſpiciferis, foliis aſperis lingulatis, calycibus hirſutis. *Hall. hiſt. n.* 603.

La Viperine. Regn. bot.

In ruderatis, pratis ſiccioribus, ad vias. Fl. æſtate.

ARETIA. Corolla hypocrateriformis. Limbus quinquefidus. Tubus ovatus, collo coarctatus. Stigma depreſſo capitatum. Capſula unilocularis, globoſa, ſubpentaſperma.

alpina. 1. A. foliis liuearibus patentibus, floribus pedunculatis. *Syſt. veget. p.* 192. *Iacqu. auſtr. app. t.* 18.

Aretia foliis ovatis repandis, ſcapis unifloris. *Hall. hiſt. n.* 618. *tab.* 11.

In Carinthia in ſupremis folkartenſium alpium jugis; in fladnizenſibus ſummitatibus circa *Leiterſteig*; in excelſo *Winterthal*; *Eiſenhut* carinthiaco & ſtyriaco. *Wulfen.* Ego in ſummis alpium jugis prope *Sterzing*, alibique ad marginem æternæ glaciei in rupium fiſſuris. Fl. a Iunio ad Auguſtum. Per.

Variat floribus albis, rubris; foliis glabris, & villoſis.

ANDROSACE. Involucrum umbellulæ. Corollæ tubus ovatus: ore glanduloſo. Capſula unilocularis, globoſa.

maxima. 1. A. perianthiis fructuum maximis. *Sp. pl. p.* 203. *Iacqu. auſt. t.* 331. *Scop. carn. tom.* 1. *p.* 130.

Androſace ſtipulis quinis ampliſſimis. *Hall. hiſt. n.* 624.

Androſaces altera Matthioli. *Clus. hiſt. p.* CXXXIV.

In cultis, alibique. Fl. primo vere.

elongata. 2. A. foliis ſubdentatis, pedicellis longiſſimis, corollis calyce brevioribus. *Syſt. veget. p.* 192. *Iacqu. auſtr. t.* 330.

In arvis, ad agrorum margines, in apricis herbidis Auſtriæ. Fl. primo vere.

obtuſifolia. 3. A. foliis ellipticis lanceolatis glabris, ſcapis umbellatis. *Hall. hiſt. n.* 621. ſub Aretia. *Allion. pedem. tom.* 1. *p.* 90. *t.* 46. *f.* 1. *Hænke in Iacqu. collect. tom.* 2. *p.* 9.

In alpibus Auſtriæ, Styriæ. Fl. Iunio, Iulio. Per.

Folia glabra, ciliata. Scapus, involucri foliola, pedicelli elongati, perianthium villis brevibus pubeſcit.

lactea. 4. A. foliis lanceolatis glabris, umbella involucris multoties longiore. *Sp. pl. p.* 204. *Iacqu. auſtr. t.* 333. *Scop. carn. tom.* 1. *p.* 131.

Aretia foliis gramineis, ſcapis paucifloris. *Hall. hiſt. n.* 622.

Sedum minus nonum. *Clus. hiſt. p.* LXI.

In alpibus. Fl. Iulio, Auguſto. Per.

villoſa. 5. A. foliis piloſis, perianthiis hirſutis. *Sp. pl. p.* 203. *Wulfen in Iacqu. collect. tom.* 1. *p.* 193. *t.* 12. *f.* 3. *Scop. carn. tom.* 1. *p.* 130.

Aretia foliis ovatis villoſis, ſcapis umbellatis. *Hall. hiſt. n.* 206.

In alpibus carniolicis. Fl. Iulio. Per.

Chamæjasme. 6. A. foliis ciliatis, perianthiis hirſutis. *Wulfen in Iacqu. collect. tom.* 1. *p.* 193. ſub A. villoſa.

Androſace villoſa. *Iacqu. auſtr. t.* 332.

Chamæjasme. *Bauh. hiſt.* 3. *p.* 869.

Sedum minus decimum alpinum quartum. *Clus. hiſt. p.* LXII.

In alpinis, & ſubalpinis. Fl. Iunio, Iulio. Per.

Folia obſita ſunt pilis rarioribus, deciduis. Cilia perſiſtunt. Corollæ laciniæ in aliis emarginatæ, apice integræ in aliis. Involucri foliola primo pedicellis æqualia, dein breviora.

PRIMULA. Involucrum umbellulæ. Corollæ tubus cylindricus: ore patulo.

acaulis. 1. P. foliis rugoſis dentatis ſubtus hirſutis, ſcapis unifloris. *Iacqu. miſcell. tom.* 1. *p.* 158.

Primula foliis rugoſis, hirſutis, dentatis, ſcapo unifloro. *Hall. hiſt. n.* 608.

Primula veris, var. acaulis, *Sp. pl. p.* 204. *Curtis lond.*

Primula ſylveſtris. *Scop. carn. tom.* 1. *p.* 132.

Primula veris pallido flore humilis. *Cluſ. hiſt. p.* 302.

In pratis, ſylvis, ad ſepes. Fl. primo vere, quandoque hyeme. Per.

officinalis. 2. P. foliis rugoſis dentatis ſubtus hirſutis, ſcapis multifloris, floribus nutantibus, corollæ limbo brevi. *Iacqu. miſcell. tom.* 1. *p.* 159.

Primula foliis rugoſis, dentatis, hirſutis, ſcapis multifloris, floribus omnibus nutantibus. *Hall. hiſt. n.* 610.

Primula veris, var. officinalis *Sp. pl. p.* 204. *Fl. dan. t.* 433.

Primula veris flavo flore elatior. *Cluſ. hiſt. p.* 301.

La Primevere. Regn. bot.

In montanis ſylvaticis, in pratis, ad ſepes, in fruticoſis, & in ipſis alpibus. Fl. Aprili, Majo, ſerius in alpibus. Per.

elatior. 3. P. foliis rugoſis dentatis hirſutis, ſcapis multifloris, floribus exterioribus nutantibus; medio erecto. *Iacqu. miſcell. tom.* 1. *p.* 158.

Primula foliis rugoſis dentatis, ſcapis multifloris, floribus exterioribus nutantibus. *Hall. hiſt. n.* 609.

Primula veris, var. elatior. *Sp. pl. p.* 204. *Fl. dan. t.* 434.

Primula veris pallido flore elatior. *Clus. hiſt. p.* 301.

In montanis ſylvaticis. Fl. Martio, Aprili. Per.

farinoſa. 4. P. foliis crenatis glabris, florum limbo plano. *Sp. pl. p.* 205. *Fl. dan. t.* 125.

Aretia foliis glabris, ellipticis, rugoſis, ſubtus farinoſis, floribus umbellatis. *Hall. hiſt. n.* 623.

Primula veris rubro flore. *Clus. hiſt. p.* 300.

In pratis humidis in Auſtria inter *Gaden* & *Heiligenkreutz*, prope *Erla*, *Lachſenburg*, *Himberg*. In Tyroli prope *Sterzing* in aſcenſu alpium. Fl. Majo, Iunio. Per.

longiflora. 5. P. foliis ſerratis glabris, umbella nutante, corollæ tubo longiſſimo. *Iacqu. auſtr. app. t.* 46. *miscell. tom.* 1. *p.* 159. *Syſt. veget. p.* 193. *Allion. pedem. tom.* 1. *p.* 91. *t.* 39. *f.* 3.

Primula foliis glabris, ellipticis, rugoſis, ſerratis, floribus umbellatis, tubo longiſſimo. *Hall. hiſt. n.* 611.

In alpibus Tyrolis, Carinthiæ, Carnioliæ, Croatiæ. Fl. Iunio, Iulio. Per.

Auricula. 6. P. foliis ſerratis glabris. *Sp. pl. p.* 205. *Iacqu. auſtr. t.* 415. *Scop. carn. tom.* 1. *p.* 133.

Primula umbellifera, foliis ſerratis, glabris, carnoſis. *Hall. hiſt. n.* 612.

Auricula urſi prima. *Clus. hiſt. p.* 302.

In rupeſtribus alpinis, & ſubalpinis. In demiſſis quoque agri vindobonenſis *in der Briel.* Fl. hic Aprili. Iunio, Iulio ibi. Per.

villoſa. 7. P. foliis ſerrulatis planis hirſutis. *Iacqu. auſtr. app. t.* 27. *miſcell. auſtr. tom.* 1. *p.* 159. *Syſt. veget. p.* 193.

Primula foliis ſubhirſutis, dentatis, ſcapo pauciſloro. *Hall. hiſt. n.* 613.

Primula pubeſcens. *Wulfen in Iacqu. miscell. tom.* 1. *p.* 157. *t.* 18. *f.* 2.

In alpibus carinthiacis, & tyrolenſibus. Fl. Iunio, Iulio. Per.

glutinoſa. 8. P. foliis lanceolatis ſerrulatis glutinoſis, involucro longitudine florum ſeſſilium. *Syſt. veget. p.* 193. *Iacqu. auſtr. app. t.* 26.

Auricula urſi ſeptima. *Clus. hiſt. p.* 304.

In alpibus tyrolenſibus, plerumque in vicinia glaciei. Fl. Iulio, Auguſto. Per.

minima. 9. P. foliis cuneiformibus nitidis, ſcapis unifloris. *Iacqu. auſtr. t.* 273. *en. p.* 202. *Hall. hiſt. n.* 614.

Auricula urſi minima. *Clus. hiſt. p.* 305.

In alpibus. Fl. æſtate. Per.

integrifolia. 10. P. foliis integerrimis glabris oblongis, calycibus tubuloſis obtuſis. *Sp. pl. p.* 205. *Iacqu. auſtr. t.* 327.

Primula foliis ciliatis ellipticis, carnoſis, integerrimis. *Hall. hiſt. n.* 615.

Auricula urſi quarta. *Clus. hiſt. p.* 303.

In alpibus auſtriacis, ſtyriacis, carniolicis. Fl. Iunio, Iulio. Per.

carniolica. 11. P. foliis integerrimis ellipticis, umbella erecta, calycibus acutis breviſſimis. *Iacqu. auſtr. app. t.* 4. *miscell. tom.* 1. *p.* 160. *Syſt. veget. p.* 193.

In Carniolia in agro idrienſi ad pedem collis incolis *Kobila* dicti, non procul a cataracta. Fl. primo vere. Per.

In hac calyx plus quam duplo brevior tubo corollæ. In integrifolia vix ultra unam tertiam ejusdem partem; eſtque præterea in illa anguſtior, in hac tubo corollæ amplior. Laciniæ petali in integrifolia multo profundius ſecantur; tubus ve-

ro baſi anguſtatur. In carniolica folia ex oblongo ovalia & ſubacuta ſunt utplurimum obſoletiſſime repanda, nec ut in integrifolia tam craſſa vereque integerrima, nec habent ad oras cilia ulla. Iacqu. l. c.

CORTUSA. Corolla rotata: fauce annulo elevato. Capſula unilocularis, ovalis, ſemiquinquevalvis.

Matthioli. 1. C. calycibus corolla brevioribus. *Sp. pl. p. 206. Cluſ. hiſt. p. 306. Allion. pedem. tom. 1. p. 93. t. 5. f. 3. Wulfen in Iacqu. collect. tom. 1. p. 236. ic. t. 32.*

Auricula urſi laciniata, ſeu Cortuſa Matthioli. *Tourn. inſt.* 123.

In alpibus infrequens. Copioſe creſcit in agro fluminenſi Terſacti in ſylva P. P. Franciſcanorum. Fl. Majo ad Iunii finem. Per.

SOLDANELLA. Corolla campanulata, lacero-multifida. Capſula unilocularis, apice multidentata.

alpina. 1. Soldanella. *Sp. pl. p. 206. Iacqu. auſtr. t. 13. Scop. carn. tom. 1. p. 135. Hall. hiſt. n. 634.*

Soldanella alpina major & minor. *Cluſ. hiſt. p.* 308. 309.

In alpibus. Fl. æſtate. Per.

Folia ex rotundo reniformia. Scapus biflorus.

CYCLAMEN. Corolla rotata, reflexa, tubo breviſſimo: fauce prominente. Bacca tecta putamine capſulari.

europæum. 1. C. corolla retroflexa. *Sp. pl. p.* 207. *Iacqu. auſtr. t.* 401.

Cyclaminus flore cernuo, ſegmentis revolutis. *Hall. hiſt. n.* 635.

Cyclaminus europæus. *Scop. carn. tom.* 1. *p.* 136.

Cyclaminus odorata purpureo flore. *Clus hiſt. p.* 264.

Le Cyclamen ou Pain de Pourceau. Regn. bot.

In ſylvis montanis, montium declivibus, alibique. Fl. Aprili, Majo, dein ab Auguſto ad Octobrim. Per.

Pedunculus, qui florifer erectus, fructtifer in ſpiras multas ſe convolvit, donec bacca terram attigerit. Semen terræ commiſſum in exiguum tuber convertitur, unde poſtea folia promit. Clus. l. c.

MENYANTHES. Corolla hirſuta. Stigma bifidum. Capſula unilocularis.

trifoliata. 1. M. foliis ternatis. *Sp. pl. p.* 208. *Fl. dan. t.* 541. *Curtis lond. Scop. carn. tom.* 1. *p.* 136. *Hall. hiſt. n.* 633.

La Meniante ou le Trefle d'eau. Regn. bot.

In aquis ſtagnantibus, & pratis udis. Fl. Majo. Per.

HOTTONIA. Corolla hypocrateriformis. Stamina tubo corollæ impoſita. Capſula unilocularis.

paluſtris. 1. H. pedunculis verticillato-multifloris. *Sp. pl. p.* 208. *Curtis lond. Fl. dan. t.* 483. *Scop. carn. tom.* 1. *p.* 137.

Hottonia florum verticillis ſpicatis. *Hall. hiſt. n.* 632.

In Auſtria in aquis ſtagnantibus prope *Bruck an der Leitha*. In Carniolia circa Labacum. Fl. Iunio. Per.

LYSIMACHIA. Corolla rotata. Capſula globoſa, mucronata, decemvalvis.

vulgaris. 1. L. paniculata, racemis terminalibus. *Sp. pl. p.* 209. *Curtis lond. Fl. dan. t.* 689. *Scop. carn. tom.* 1. *p.* 138.

Lyſimachia foliis ovato-lanceolatis, ſpicis paniculatis. *Hall. hiſt. n.* 630.

Lyſimachia lutea communis. *Clus. hiſt. p. L.*

Ad fluvios, foſſas, ad ſepes humidas, in pratis udis. Fl. æſtate. Per.

thyrſiflora. 2. L. racemis lateralibus pedunculatis. *Sp. pl. p.* 209. *Fl. dan. t.* 517.

Lyſimachia racemis alaribus, denſiſſimis, folio brevioribus. *Hall. hiſt. n.* 631.

Lyſimachia lutea tertia ſive minor. *Clus. hiſt. p. LIII.*

In paludoſis Auſtriæ prope *Weitra. R. Liborius Miller.* Fl. æſtate. Per.

punctata. 3. L. foliis ſubquaternis, pedunculis verticillatis unifloris. *Sp. pl. p.* 210. *Iacq. auſtr. t.* 366. *Scop. carn. tom.* 1. *p.* 138.

Lyſimachia lutea ſecunda. *Clus. hiſt. p. LII.*

In udis ſylvaticis, paluſtribusque. Fl. Iunio, Iulio. Per.

Tota planta ſpirat gravem tabaci odorem.

nemorum. 4. L. foliis ovatis acutis, floribus ſolitariis, caule procumbente. *Sp. pl. p.* 211. *Curtis lond. Fl. dan. t.* 174.

Lyſimachia caule decumbente, foliis ovato-lanceolatis, petiolis alaribus unifloris. *Hall. hiſt. n.* 628.

Anagalis flavo flore. *Clus. hiſt. p.* CLXXXII.

In Auſtria, Styria in montanis, ac ſubalpinis, locis umbroſis, & riguis. Fl. æſtate. Per.

Nummularia. 5. L. foliis ſubcordatis, floribus ſolitariis, caule repente. *Sp. pl. p.* 211. *Fl. dan. t.* 493. *Curtis lond. Scop. carn. tom.* I. *p.* 139.

Lyſimachia caule proſtrato, foliis ſubrotundis, petiolis alaribus unifloris. *Hall. hiſt. n.* 629.

La Nummulaire. Regn. bot.

In humidis. Fl. æſtate. Per.

ANAGALLIS. Corolla rotata. Capſula circumſciſſa.

arvenſis. 1. A. foliis indiviſis, caule procumbente. *Sp. pl. p.* 211. *Curtis lond. Fl. dan. t.* 88.

Anagallis caule procumbente, foliis ovato-lanceolatis, petalis ſerratis, calycis ſegmentis ſubulatis. *Hall. hiſt. n.* 625.

Anagallis phœnicea. *Scop. carn. tom.* I. *p.* 139.

Anagallis purpuraſcente flore. *Clus. hiſt. p.* CLXXXIII.

Le Mouron. Regn. bot.

In arvis, hortis, & alibi. Fl. æſtate.

Variat floribus cœruleis, ac phœniceis.

tenella. 2. A. foliis ovatis acutiuſculis, caule repente. *Syſt. veget. p.* 196. *Curtis lond.*

Lyſimachia foliis ſubrotundis, caule repente, filamentis villoſis. *Ger. prov. p.* 307.

Lyſimachia tenella. *Sp. pl. p.* 211.

In Carnia ad litus lacus; qui in viciniis montis *Croſtis. I. Hoſt.* Fl. æſtate. Per.

Caules repentes. Folia oppoſita. Pedunculi uniflori, axillares. Perianthium quinquepartitum. Filamenta, ut in præcedente, pilis articulatis munita.

AZALEA. Corolla campanulata. Stamina receptaculo inſerta. Capſula quinquelocularis.

procumbens. 1. A. ramis diffuſo procumbentibus. *Sp. pl. p.* 215. *Fl. lapp. n.* 90. *t.* 6. *f.* 2. *Fl. dan. t.* 9. *Scop. carn. tom.* 1. *p.* 140.

Azalea caule procumbente, foliis ovatis, margine introverſo. *Hall. hiſt. n.* 666.

Chamæciſtus ſeptimus. *Clus. hiſt. p.* 75.

In alpibus. Fl. Iunio, Iulio. Fr.

CONVOLVULUS. Corolla campanulata, plicata. Stigmata duo. Capſula bilocularis: loculis diſpermis.

arvenſis. 1. C. foliis ſagittatis utrinque acutis, pedunculis unifloris. *Sp. pl. p.* 218. *Fl. dan. t.* 459. *Scop. carn. tom.* 1. *p.* 140.

Convolvulus foliis ſagittatis, latescentibus, petiolis unifloris, ſtipulis remotis, ſubulatis. *Hall. hiſt. n.* 664.

Convolvulus minor. *Clus. hiſt. p. L.*

Le petit Liſeron. Regn. bot.

In cultis, ad vias, & alibi. Fl. æſtate. Per.

ſepium. 2. C. foliis ſagittatis poſtice truncatis, pedunculis tetragonis unifloris. *Sp. pl. p.* 218. *Curtis lond. Fl. dan. t.* 458. *Scop. carn. tom.* 1. *p.* 141.

Convolvulus foliis ſagittatis, hamis emarginatis anguloſis, petiolis unifloris, ſtipulis cordatis maximis. *Hall. hiſt. n.* 663.

Ad ſepes humidas, in fruticoſis. Fl. æſtate. Per.

Cantabrica. 3. C. foliis lineari-lanceolatis acutis piloſis, pedunculis inferioribus foliis longioribus ſubbifloris, calycibus oblongo-lanceolatis hirſutis. *Ait. Kew. tom.* 1. *p.* 213. *Sp. pl. p.* 225. *Iacqu. auſtr. t.* 296. *Scop. carn. tom.* 1. *p.* 141.

Convolvulus foliis lineari-lenceolatis acutis, caule ramoſo erectiusculo, pedunculis ſubbifloris. *Ger. prov. p.* 318.

In apricis aſperis in Litorali, Forojulio, & in Auſtria ad Thermas badenſes. Fl. æſtate. Per.

Soldanella. 4. C. foliis reniformibus, pedunculis unifloris. *Sp. pl. p.* 226. *Scop. carn. tom.* 1. *p.* 242.

Soldanella maritima minor. *Bauh. pin.* 295.

Braſſica marina, ſive Soldanella. *Bauh. hiſt.* 2. *p.* 166. *Zann. iſt. p.* 71. *t.* 6.

La Soldanelle. Regn. bot.

In arenoſis Litoralis. Fl. Majo, Iunio.

POLEMONIUM. Corolla quinquepartita, fundo clauſo valvis ſtaminiferis. Stigma trifidum. Capſula trilocularis, ſupera, apice trifafariam dehiscens.

cœruleum. 1. P. foliis pinnatis, floribus erectis, calycibus tubo corollæ longioribus. *Sp. pl. p.* 230. *Fl. dan. t.* 255.

Polemonium foliis pinnatis, floribus erectis, foliolis elliptico-lanceolatis. *Hall. hiſt. n.* 665.

Valeriana cœrulea. *Bauh. pin.* 164.

In Carnia prope *Mion. I. Hoſt.* Fl. Iunio, Iulio. Per.

Variat colore florum albo, cœruleo.

CAMPANULA. Corolla campanulata fundo clauſo valvis ſtaminiferis. Stigma trifidum. Capſula iufera, nuda, vel obtecta ſinubus calycis reflexis, tri- vel quinquelocularis, totidemque poris lateralibus emittens ſemina.

* *Foliis laevioribus.*

pulla. 1. C. caulibus unifloris, foliis caulinis ovatis crenatis, calycibus cernuis. *Sp. pl. p.* 231. *Iacqu. auſtr.* t. 285.

Campanula alpina latifolia, flore pullo. *Bauh. pin.* 93. *prodr.* 33.

In Auſtriae, Styriae alpinis, & ſubalpinis. Fl. aeſtate. Per.

Zoiſii. 2. C. caulibus ſubtrifloris, foliis integerrimis: radicalibus ovatis petiolatis, caulinis oblongis ſeſſilibus, floribus nutantibus. *Wulfen in Iacqu. collect. tom.* 2. *p.* 122. *ic. t.* 334.

In Carniolia in rupeſtribus alpium wochinenſium. *Zois.* Fl. Iulio. Per.

rotundifolia. 3. C. foliis radicalibus reniformibus, caulinis linearibus. *Sp. pl. p.* 232. *Curtis lond.*

Campanula foliis ſerratis, radicalibus cordatis, caulinis lanceolatis. *Hall. hiſt. n.* 701.

Campanula minor alpina rotundioribus imis foliis. *Clus. hiſt. p. CLXXIII.*

In pratis, collibus, ad vias, in alpinis, & ſubalpinis. Fl. aeſtate. Per.

Variat foliis radicalibus ovatis; caulinis linearibus, & lanceolatis; caule ramoſo, ac ſimplici.

patula. 4. C. foliis ſtrictis: radicalibus lanceolato-ovalibus, panicula patula. *Sp. pl. p.* 232. *Fl. dan. t.* 375. *Scop. carn. tom.* 1. *p.* 145.

Campanula caule brachiato, foliis lanceolato-linearibus glabris, petiolis pauciflor is nudis. *Hall. hiſt. n.* 698.

In pratis. Fl. Majo, Iunio.

Rapunculus. 5. C. foliis undulatis: radicalibus lanceoleto-ovalibus, panicula coarctata. *Sp. pl. p.* 232. *Scop. carn. tom.* 1. *p.* 147.

Campanula foliis ellipticis ſerratis, ſubhirſutis, caule aſpero, floribus paniculatis. *Hall. hiſt. n.* 699.

Campanula radice eſculenta, flore cœruleo. *Tourn. inſt.* 110.

Rapunculus eſculentus. *Bauh. pin.* 92. *Zann. iſt. p.* 47. *t.* 102.

In pratis Forijulii, Litoralis. Fl. Majo, Iunio.

Radix biennis, napiformis. Caulis pedalis, altiorque, ſulcatus, pilis brevibus rigidis, ut & folia, inſperſus, ubi flores glaber. Folia ſunt undulata: radicalia lanceolato-ovata, in petiolum decurrentia; caulina lanceolata, ſeſſilia. Flores paniculati. Laciniæ calycis ſetaceæ, uno alterove denticulo notatæ. Pericarpium triloculare.

Folia cum radice in cibum cedunt incolis ſub acetarii forma primo vere.

perſicifolia. 6. C. foliis radicalibus obovatis, caulinis lanceolato-linearibus ſubſerratis ſeſſilibus remotis. *Syſt. veget. p.* 206. *Scop. carn. tom.* 1. *p.* 145.

Campanula foliis radicalibus ovatis, caulinis linearibus, petiolis longiſſimis paucifloris. *Hall. hiſt. n.* 697.

Campanula perſicæ folio. *Clus. hiſt. p. CLX.*

In ſylvaticis. Fl. æſtate. Per.

pyramidalis. 7. C. foliis lævibus ſerratis cordatis: caulinis lanceolatis, caulibus junceis ſimplicibus, umbellis ſeſſilibus lateralibus. *Syſt. veget. p.* 206. *Scop. carn. tom.* 1. *p.* 146.

Campanula lactescens pyramidalis. *Cluſ. hiſt. p. CLXXI.*

In Carniolia, Forojulio, Litorali in rupibus, maceriis, muris veſtutis. Fl. Iulio, Auguſto. Per.

carpatica. 8. C. foliis omnibus cordatis ſerratis petiolatis glabris, ramis filiformibus unifloris. *Ait. Kew. tom. 1, p. 219. Iacqu. hort. tom. 1. p. 22. t. 37. Suppl. p. 140.*

In Pannonia in montibus carpaticis. Fl. Iulio. Per.

** *Foliis ſcabris.*

rapunculoides. 9. C. foliis cordato-lanceolatis, caule ramoſo, floribus ſecundis ſparſis, calycibus reflexis. *Syſt. veget. p. 207.*

Campanula foliis cordato-lanceolatis, ſubhirſutis, petiolis unifloris. *Hall. hiſt. n.* 692.

Campanula hortenſis, rapunculi radice. *Bauh. pin.* 94. *Moris. hiſt.* 2. *p.* 460. *ſect.* 5. *t.* 3. *f.* 32.

In cultis, pratis, collibus, ad vias. Fl. æſtate. Per.

bononienſis. 10. C. foliis ovato-lanceolatis ſubtus ſcabris: radicalibus petiolatis, caulinis ſeſſilibus, caule paniculato. *Sp. pl. p.* 234. *Scop. carn. tom.* 1. *p.* 147.

Campanula caule erecto ramoſiſſimo, foliis ovato-lanceolatis, ſubtus tomentoſis, petiolis multifloris erectis. *Hall. hiſt. n.* 689.

Campanula ſeu Cervicaria bononienſis flore parvo. *Bauh. hiſt.* 2. *p.* 806.

In pratis ſiccioribus Auſtriæ, Pannoniæ, Litoralis. Fl. Iulio, Auguſto. Per.

Folia radicalia petiolata plerumque, antequam planta florere incipit, marceſcunt.

gramini-folia. 11. C. foliis lineari-ſubulatis, capitulo terminali. *Sp. pl. p.* 234.

Trachelium tragopogi folio, montanum. *Col. phyt.* 2. *p.* 25.

Trachelium minus gramineum cœruleo-violaceum. *Barr. ic.* 332.

In agro fluminenſi in herbidis, aſperisque collibus ad latera viæ, quæ prope Draga a via regia corolina Grobnicum ducit, nec non ad pedem montium *mala Planina* dictorum. Fl. Iunio, Iulio. Per.

Radix promit caules plures palmares, vel ſpithamalas, alios erectos, procumbentes alios. Folia ſunt ſeſſilia, ſubulata, pleraque ob margines conniventes canaliculata. Caulis pilis ſcaber deſinit in capitulum florum, quod involvitur foliis baſi valde dilatatis, cæterum caulinis ſimilibus. Flores ſeſſiles. Corolla interne, & valvulæ nectariferæ pilis obſitæ ſunt. Stigma bifidum.

Trachelium. 12. C. caule angulato, foliis petiolatis, calycibus ciliatis, pedunculis trifidis. *Sp. pl. p.* 235. *Scop. carn. tom.* 1. *p.* 148.

Campanula foliis cordato-lanceolatis, acute ſerratis, hirſutis, petiolis ex alis paucifloris, calycibus hiſpidis. *Hall. hiſt. n.* 690.

Trachelium vulgare. *Clus. hiſt. p. CLXX.*

In fruticoſis, ſylvis, ad ſepes. Fl. Iunio, Iulio. Per.

glomerata. 13. C. caule angulato ſimplici, floribus ſeſſilibus, capitulo terminali. *Sp. pl. p.* 235. *Scop. carn. tom.* 1. *p.* 148.

Campanula caule ſimplici, aſpero, foliis amplexicaulibus, floribus capitatis. *Hall. hiſt. n.* 685.

In pratis ſiccioribus, aliisque herbidis, ac aſperis locis. Fl. æſtate. Per.

Cervicaria. 14. C. hiſpida, floribus ſeſſilibus, capitulo terminali, foliis lanceolato-linearibus undulatis. *Sp. pl. p.* 235. *Fl. dan. t.* 787.

Campanula caule ſimplici, aſpero, foliis lingulatis, aſperis, floribus capitatis. *Hall. hiſt. n.* 686.

In Pannonia in agro poſonienſi ad margines ſylvarum. Fl. Iunio, Iulio.

thyrſoidea. 15. C. hiſpida, racemo ovato-oblongo terminali, caule ſimpliciſſimo, foliis lanceolato-linearibus. *Sp. pl. p.* 235. *Iacqu. auſtr. t.* 411. *Scop. carn. tom.* 1. *p.* 148.

Campanula foliis lingulatis aſperis, floribus hirſutis, denſiſſime ſpicatis. *Hall. hiſt. n.* 688.

Trachelium thyrſoides. *Clus. hiſt. p. CLXXII.*

In alpibus Carnioliæ; in pratis Auſtriæ ſubalpinis. Fl. Iulio.

*** *Capſulis obtectis calycis ſinubus reflexis.*

Medium. 16. C. capſulis quinquelocularibus obtectis, caule indiviſo, erecto, folioſo, floribus erectis. *Sp. pl. p.* 236. *Scop. carn. tom.* 1. *p.* 149.

Viola mariana quorundam. *Clus. hiſt. p. CLXXII.*

Campanula hortenſis, folio & flore oblongo. *Bauh. pin.* 94.

In aſperis Litoralis, Forijulii. Fl. Iunio.

barbata. 17. C. capſulis quinquelocularibus obtectis, caule ſimpliciſſimo ſubunifolio, foliis lanceolatis, corollis barbatis. *Syſt. veget. p.* 208. *Iacqu. obs.* 1. *p.* 14. *t.* 37.

Campanula caule ſimplici, foliis lingulatis, aſperis, petiolis unifloris, floribus hirſutis. *Hall. hiſt. n.* 694.

Campanula foliis Echii, floribus villoſis. *Bauh. pin.* 94. *prodr.* 36.

In ſubalpinis, & alpinis. Fl. æſtate. Per.

ſpicata. 18. C. hiſpida, ſpica laxa floribus alternis, foliis linearibus. *Sp. pl. p.* 234. *Allion. pedem. tom.* 1. *p.* 112. *t.* 46. *f.* 2. *t.* 47. *f.* 1.

Campanula caule ſimplici aſpero, foliis linearibus, floribus laxe, & longiſſime ſpicatis. *Hall. hiſt. n.* 687.

Campanula alpina altiſſima hirſuta parvo flore. *Tourn. inſt.* 109.

In Carniolia in aſperis montium comilatus tolmienſis. Fl. Iunio, Iulio.

Tota planta eſt pilis albidis hiſpida. Folia alia ſunt integerrima, alia margine undulata, obiter crenata: radicalia in petiolum decurrentia; caulina ſeſſilia, baſi ampliata caulem amplexantia. Flores axillares in ſpicam longiſſimam digeſti.

alpina. 19. C. caule ſimplici, pedunculis unifloris axillaribus diphyllis. *Sp. pl. app. p.* 1669. *Iacqu. auſtr. t.* 118.

Campanula foliis ellipticis hirſutis, petiolis alaribus unifloris, floribus glabris. *Hall. hiſt. n.* 695.

Trachelium pumilum alpinum. *Clus. hiſt. p. CLXXI.*

In ſummis alpium jugis. Fl. Iunio, Iulio, Auguſto. Per.

ſibirica. 20. C. capſulis trilocularibus obtectis, caule paniculato. *Sp. pl. p.* 236. *Iacqu. auſtr. t.* 200. *Gmel. ſib.* 3. *p.* 154. *t.* 29.

In Auſtriæ, Pannoniæ pratis, collibus. Fl. Majo, Iunio.

Speculum. 21. C. caule ramoſiſſimo diffuſo, foliis oblongis ſubcrenatis, floribus ſolitariis, capſulis prismaticis. *Syſt. veget. p.* 209. *Curtis magaz.* 102. *Scop. carn. tom.* 1. *p.* 149.

Campanula caule procumbente, fructibus prismaticis, floribus ſolitariis, rotatis, longiſſime petiolatis. *Hall. hiſt. n.* 703.

Inter ſegetes. Fl. Iunio, Iulio.

hybrida. 22. C. caule baſi ſubramoſo ſtricto, foliis oblongis crenatis, calycibus aggregatis corolla longioribus, capſulis prismaticis. *Sp. pl. p.* 239.

Campanula caule erecto, fructibus priſmaticis, floribus rotatis, ſeſſilibus, congeſtis. *Hall. hiſt. n.* 704.

In vineis agri fluminenſis. Fl. Majo, Iunio.

PHYTEUMA. Corolla rotata, quinquepartita: laciniis linearibus. Stigma bi- ſeu trifidum. Capſula infera, bi- ſeu trilocularis.

pauciflora. 1. P. capitulo ſubfolioſo, foliis omnibus lanceolatis. *Sp. pl. p.* 241.

Rapunculus foliis ovatis, obtuſe-dentatis, bracteis maximis obtuſis. *Hall. hiſt. n.* 680.

Rapunculus pauciflorus. *Scop. carn. tom.* 1. *p.* 152.

Rapunculus alpinus parvus comoſus. *Bauh. hiſt.* 2. *p.* 811.

In alpibus Styriæ, Carnioliæ, Carinthiæ, Tyrolis. Fl. Iulio. Per.

hemisphærica. 2. P. capitulo ſubrotundo, foliis linearibus ſubintegerrimis. *Syſt. veget. p.* 211. *Hænke in Iacqu. collect. tom.* 2. *p.* 61. *ic. t.* 333.

Rapunculus foliis linearibus, bracteis ovato-lanceolatis. *Hall. hiſt. n.* 679.

Rapunculus umbellatus folio gramineo. *Bauh. pin.* 92.

In alpibus tyrolenſibus, carinthiacis, ſtyriacis. Fl. Iulio, Auguſto. Per.

comoſa. 3. P. faſciculo terminali ſeſſili, foliis dentatis: radicalibus cordatis. *Sp. pl. p.* 242. *Iacqu. auſtr. t.* 50.

Trachelium petræum minus. *Pona in Clus. hiſt. p. CCCXXVI.*

In Carnia Malborghetti in rupeſtribus vallis frigidæ, qua torrens malborghettanus erumpit. *Wulfen; frà i Ponti. I. Hoſt.* Fl. Iulio. Per.

orbicularis. 4. P. capitulo ſubrotundo, foliis ſerratis radicalibus cordatis. *Sp. pl. p.* 242. *Iacqu. auſtr. t.* 437.

Rapunculus foliis imis cordatis, oblongis, caulinis ſeſſilibus acutis, bracteis ovato-lanceolatis. *Hall. hiſt. n.* 681.

Rapunculus orbicularis. *Scop. carn. tom.* 1. *p.* 150.

In collibus apricis, in montibus, in alpinis, & ſubalpinis. Fl. Iunio, tardius in editioribus. Per.

ſpicata. 5. P. ſpica oblonga, capſulis bilocularibus, foliis radicalibus cordatis. *Sp. pl. p.* 242. *Fl. dan. t.* 362.

Rapunculus foliis radicalibus cordatis, tubis bicornibus revolutis. *Hall. hiſt. n.* 684.

Rapunculus ſpicatus. *Scop. carn. tom.* 1. *p.* 150.

In ſylvis montanis, & ſubalpinis. Fl. a Majo ad Iulium. Per.

In Carnioliæ alpibus wochinenſibus in herbidis montis czerna Perſt *dicti inveni varietatem floribus cœruleis.*

SAMOLUS. Corolla hypocrateriformis. Stamina munita ſquamulis ád baſin ſinus limbi corollæ conniventibus. Capſula unilocularis, infera.

Valerandi. 1. Samolus. *Sp. pl. p.* 243. *Fl. dan. t.* 198.

Anagallis aquatica, folio rotundo non crenato. *Bauh. pin.* 252.

In Auſtria in paluſtribus prope *Schwadorf, Himberg.* In Pannonia ad litus Lacus Peiſonis. Fl. æſtate.

LONICERA. Corolla monopetala, irregularis. Bacca polyſperma, bilocularis, infera.

* *Caule volubili.*

Caprifolium. 1. L. floribus verticillatis terminalibus ſeſſilibus; foliis ſummis connato-perfoliatis. *Sp. pl. p.* 246. *Iacqu. auſtr. t.* 357. *Scop. carn. tom.* 1. *p.* 152.

Periclymenum perfoliatum. *Bauh. pin.* 302.

Periclymenum vulgare alterum. *Cluſ. hiſt. p.* 57.

Il Caprifoglio, o Madreſelva. Zann. iſt. p. 48. *t.* 130.

In ſylvaticis. Fl. Majo, Iunio. Fr.

Periclymenum. 2. L. capitulis ovatis imbricatis terminalibus, foliis omnibus diſtinctis. *Sp. pl. p.* 247. *Curtis lond. Scop. carn. tom.* 1. *p.* 153. *Hall. hiſt. n.* 673.

Periclymenum non perfoliatum germanicum. *Bauh. pin.* 302.

In Forojulio in agro goritienſi. *Scop.* Fl. æſtate. Fr.

** *Pedunculis bifloris.*

nigra. 3. L. pedunculis bifloris, baccis diſtinctis, foliis ellipticis integerrimis. *Sp. pl. p.* 247. *Iacqu. auſtr. t.* 314.

Caprifolium foliis ovato-lanceolatis, glabris, baccis gemellis, calyce quinquefido. *Hall. hiſt. n.* 676.

Periclymenum rectum ſecundum. *Cluſ. hiſt. p.* 58.

Chamæceraſus alpina fructu nigro gemino. *Bauh. pin.* 451.

In ſylvis ſubalpinis. Fl. Majo, Iunio. Fr.

Xyloſte-um. 4. L. pedunculis bifloris, baccis diſtinctis, foliis integerrimis pubeſcentibus. *Sp. pl. p.* 248. *Fl. dan. t.* 808. *Scop. carn. tom.* 1. *p.* 153.

Caprifolium foliis ovatis, acuminatis, ſubhirſutis, baccis gemellis. *Hall. hiſt. n.* 677.

Chamæceraſus dumetorum, fructu gemino rubro. *Bauh. pin.* 451.

In ſylvis, fruticoſis. Fl. Majo. Fr.

alpigena. 5. L. pedunculis bifloris, baccis coadunatis didymis, foliis ovali-lanceolatis. *Syſt. veget. p.* 216. *Iacqu. auſtr. t.* 274. *Scop. carn. tom.* 1. *p.* 154.

Caprifolium foliis ovato-lanceolatis, ſubhirſutis, bacca ſingulari, bicolli, biflora. *Hall. hiſt. n.* 675.

Periclymenum rectum quartum. *Clus. hiſt. p.* 59.

Chamæceraſus alpina, fructu rubro gemino: duobus punctis notato. *Bauh. pin.* 451.

In alpinis, & ſubalpinis. Fl. Iunio, Iulio. Fr.

cærulea. 6. L. pedunculis bifloris, baccis coadunatis globoſis, ſtylis indiviſis. *Sp. pl. p.* 249. *Iacqu. auſtr. app. t.* 17. *Scop. carn. tom.* 1. *p.* 154.

Caprifolium foliis ovatis, bacca ſingulari, ovata, biflora. *Hall. hiſt. n.* 674.

Chamæceraſus montana, fructu ſingulari cæruleo. *Bauh. pin.* 451.

Periclymenum rectum tertium. *Clus. hiſt. p.* 58.

In alpinis, & ſubalpinis Carnioliæ, Styriæ, Carinthiæ, Tyrolis. Fl. Iunio, Iulio. Fr.

CORIS. Calyx ſpinoſus. Corolla monopetala, irregularis. Capſula quinquevalvis, ſupera.

monſpeli-enſis. 1. Coris. *Sp. pl. p.* 252.

Coris quorundam. *Clus. hiſt. p.* CLXXIV.

Coris cœrulea maritima. *Bauh. pin.* 280.

In aſperis agri tergeſtini. *I. Hoſt.* Fl. Majo, Iunio. Per.

VERBASCUM. Corolla rotata, ſubinæqualis. Capſula unilocularis, bivalvis.

Thapſus. 1. V. foliis decurrentibus utrinque tomentoſis, caule ſimplici. *Syſt. veget. p.* 219. *Fl. dan. t.* 361. *Scop. carn. tom.* 1. *p.* 155.

Le Bouillon blanc. Regn. bot.

In glareoſis, ad fluvios, vias, & alibi. Fl. æſtate.

Thapſoides. 2. V. foliis decurrentibus, caule ramoſo. *Syſt. veget. p.* 219.

Habitat & floret cum præcedente.

Lychnitis. 3. V. foliis cuneiformi-oblongis. *Sp. pl. p.* 253. *Fl. dan. t.* 586. *Scop. carn. tom.* 1. *p.* 155.

Verbaſcum ſpica ramoſa, foliis ovato-lanceolatis, inferioribus petiolatis. *Hall. hiſt. n.* 583.

In Carniolia in cultis circa *Gartſchereutz. Scop.*

nigrum. 4. V. foliis cordato-oblongis petiolatis. *Sp. pl. p.* 253. *Plenk. ic. t.* 110. *Scop. carn. tom.* 1. *p.* 156.

Verbaſcum foliis imis petiolatis, cordato-lanceolatis, ſuperioribus ſeſſilibus, ovato-lanceolatis. *Hall. hiſt. n.* 584.

Ad radices montium, ad pagos, vias. Fl. æſtate. Per.

phœniceum. 5. V. foliis ovatis nudis crenatis radicalibus, caule ſubnudo racemoſo. *Sp. pl. p.* 254. *Iacqu. auſtr. t.* 125.

In agri vindobonenſis arvis reſtibilibus. In Pannonia in pratis inter *Bahrendorf* & *Bruck an der Leitha.* Fl. Majo, Iunio. Per.

Blattaria. 6. V. foliis amplexicaulibus oblongis glabris, pedunculis ſolitariis. *Sp. pl. p.* 254. *Scop. carn. tom.* 1. *p.* 157.

Verbaſcum ſpica rariſſima, foliis glabris, ſerratis, cordato-lanceolatis, imis ſemipinnatis, ſuperioribus amplexicaulibus. *Hall. hiſt. n.* 585.

Blattaria flore luteo. *Besl. aichſt. æſt. ord.* 10. *t.* 4. *f.* 2.

In uliginoſis, udis, ad vias. Fl. æſtate.

Variat floribus flavis, & albis.

DATURA. Corolla infundibuliformis plicata. Calyx tubuloſus, angulatus, deciduus. Capſula quadrivalvis.

Stramonium. 1. D. pericarpiis ſpinoſis erectis ovatis, foliis ovatis glabris. *Sp. pl. p.* 255. *Fl. dan. t.* 436.

Stramonium foliis angulоſis, fructu erecto, muricato, calyce pentagono. *Hall. hiſt. n.* 586.

Stramonium fœtidum. *Scop. carn. tom.* 1. *p.* 157.

La Stramoine. Regn. bot.

In ruderatis, ad fimeta, pagos, vias. Fl. æſtate.

HYOSCIAMUS. Corolla infundibuliformis, obtuſa. Stamina inclinata. Capſula circumſciſſa operculo horizontaliter dehiſcente, bilocularis.

niger. 1. H. foliis amplexicaulibus ſinuatis, floribus ſeſſilibus. *Sp. pl. p.* 257. *Plenk ic. t.* 97. *Scop. carn. tom.* 1. *p.* 158. *Hall. hiſt. n.* 580.

Hyoſciamus niger vulgaris. *Clus. hiſt. p.* LXII.

In cultis, ruderatis, ad pagos, vias. Fl. æſtate.

Scopolia. 2. H. foliis ovatis integris, calycibus inflatis campanulatis lævibus. *Syſt. veget. p.* 221. *Scop. carn. tom.* 1. *p.* 158.

Scopolia carniolica. *Iacqu. obſ.* 1. *p.* 32. *t.* 20.

In Carniolia in ſylvis umbroſis circa Idriam. *Scop.* Ego in Croatia ad pedem montis a flumineniſibus *Risnik* vocati, aliisque vicinis locis. Fl. Majo, Iunio. Per.

ATROPA. Corolla campanulata. Stamina diſtantia. Bacca globoſa, bilocularis.

Belladonna. 1. A. caule herbaceo, foliis ovatis integris. *Sp. pl. p.* 260. *Iacqu. auſtr. t.* 309. *Curtis lond.*

Belladonna caule herbaceo brachiato, foliis ovato-lanceolatis, integerrimis. *Hall. hiſt. n.* 579.

Belladonna trichotoma. *Scop. carn. tom.* 1. *p.* 160.

Solanum lethale. *Cluſ. hiſt. p. LXXXVI.*

In ſylvis montanis & ſubalpinis. Fl. Iunio, Iulio. Per.

PHYSALIS. Corolla rotata. Stamina conniventia. Bacca intra calycem inflatum, bilocularis.

Alkekengi. 1. P. foliis geminis integris acutis, caule herbaceo inferne ſubramoſo. *Sp. pl. p.* 262. *Plenk ic. t.* 124.

Phyſalis foliis conjugatis, cordatis ſinuatis, caule ramoſo. *Hall. hiſt. n.* 577.

Phyſalis Halicacabum. *Scop. carn. tom.* 1. *p.* 160.

Le Coqueret. Regn. bot.

In ſylvis, ad ſepes, pagos, & in cultis. Fl. æſtate. Per.

SOLANUM. Corolla rotata. Antheræ ſubcoalitæ, apice poro gemino dehiſcentes. Bacca bilocularis.

Dulcamara. 1. S. caule inermi frutescente flexuoſo, foliis ſuperioribus haſtatis, racemis cymoſis. *Sp. pl. p.* 264. *Curtis lond. Fl. dan. t.* 607. *Scop. carn. tom.* 1. *p.* 161.

Solanum caule flexuoſo, frutescente, foliis cordatis & tripartitis. *Hall. hiſt. n.* 575.

La Morelle grimpante. Regn. bot.

In fruticoſis, ad ſepes. Fl. æſtate.

nigrum. 2. S. caule inermi herbaceo, foliis ovatis dentato-angulatis, racemis diſtichis nutantibus. *Syſt. veget. p.* 224. *Curtis lond. Fl. dan. t.* 460. *Scop. carn. tom.* 1. *p.* 162. *Hall. hiſt. n.* 576.

La Morelle a fruit noir. Regn. bot.

In cultis, ruderatis, ad vias, & alibi. Fl. æſtate.

RHAMNUS. Calyx nullus. Corollæ tubus turbinato-cylindraceus. Limbus patens, diviſus. Squamulæ ad baſin ſingulæ diviſuræ. Stamina tot, quot laciniæ corollæ, petalo ſub ſquamula inſerta. Bacca.

catharticus. 1. R. ſpinis terminalibus, floribus quadrifidis dioicis, foliis ovatis, caule erecto. *Syſt. veget. p.* 232. *Fl. dan. t.* 850. *Scop. carn. tom.* 1. *p.* 162.

Rhamnus ramis ſpinoſis, foliis ovato-lanceolatis, ſerratis. *Hall. hiſt. n.* 824.

Rhamnus ſolutivus. *Clus. hiſt. p.* 111.

Spina infectoria pumila prima. *Clus. hiſt. p.* 111.

Le Nerprun. Regn. bot.

In ſylvis, collibus, ad ſepes. Fl. Majo, Junio. Arb.

Spina infectoria pumila prima Cluſii eſt Rhamnus catharticus fruticis modo creſcens, præditus caule brevi, erecto, ramis vel adſcendentibus, vel erectis, foliis anguſtioribus, floribus minoribus. Non tantum in Auſtriæ monte Leithaberg, *ſed alibi etiam in ſylvis, collibus, ad vineas creſcit hæc varietas.*

ſaxatilis. 2. R. ſpinis terminalibus, floribus quadrifidis dioicis, caulibus procumbentibus. *Iacqu. auſtr. t.* 53. *en. p.* 212. *Sp. pl. app. p.* 1671.

Rhamnus infectorius. *Wulfen in Iacqu. collect. tom.* 3. *p.* 17.

Rhamnus Lycium. *Scop. carn. tom.* 1. *p.* 163.

Spina infectoria pumila ſecunda. *Cluſ. hiſt. p.* 111.

In apricis, ſaxoſis aſperis Auſtriæ, Pannoniæ, Litoralis, Carnioliæ. Fl. Majo, Iunio. Fr.

Caules exeunt ex unica radice plures, procumbentes & diffuſi. Folia alia lanceolata, quædam ovato-lanceolata, nec rarum eſt invenire plura ovata. Omnia dorſo villoſula, dum juniora; glabra, dum adultiora. Flores in aliis individuis ſteriles, fertiles in aliis. Sterilium corolla paulo longior, quadrifida. Stamina quatuor. Germen ſubrotundum & parvum. Styli duo, breves, truncati. In floribus fertilibus tubus corollæ brevior, limbo magis erecto. Filamenta, ſi adſunt, antheris orbata ſunt. Squamæ, ut in floribus ſterilibus, ſæpe deſiderantur. Germen tubi corollæ magnitudine. Styli duo ſubulati, ſuperne reflexi. Baccæ nigræ. An Rhamnus infectorius, cui baccæ flavæ, differat a noſtro ſaxatili, doceant autoptæ.

alpinus. 3. R. inermis, floribus dioicis, foliis duplicato-crenatis. *Sp. pl. p.* 280. *Scop. carn. tom.* 1. *p.* 163.

Rhamnus inermis, foliis ovatis crenulatis. *Hall. hiſt. n.* 823. *t.* 40.

In montanis, atque ſubalpinis Carnioliæ, Croatiæ. Fl. Majo, Iunio. Fr.

pumilus. 4. R. inermis repens, floribus hermaphroditis, foliis ſerratis. *Syſt. veget. p.* 234.

Rhamnus pumilio. *Wulfen in Iacqu. collect. tom.* 2. *p.* 141. *t.* 11.

Rhamnus rupeſtris. *Scop. carn. tom.* 1. *p.* 164. *t.* 5.

In rupeſtribus Forijulii, Litoralis, Carnioliæ calidioris. Fl. Majo, Iunio. Fr.

Frangula. 5. R. inermis floribus monogynis hermaphroditis, foliis integerrimis. *Sp. pl. p.* 280. *Fl. dan. t.* 278. *Scop. carn. tom.* 1. *p.* 164.

Rhamnus inermis, foliis ovato-lanceolatis, integris, floribus quinquefidis androgynis. *Hall. hiſt. n.* 821.

In nemoribus, ad ſepes humidas. Fl. æſtate. Fr.

Paliurus. 6. R. aculeis geminatis: inferiore reflexo, floribus trigynis. *Sp. pl. p.* 281. *Pallas roſſic. t.* 64.

Rhamnus folio ſubrotundo, fructu compreſſo. *Bauh. pin.* 477.

In Litorali, Forojulio. Fl. Majo, Iunio. Fr.

Frequentatur, dum floret, a Cerambyce Kæhleri.

Zizyphus. 7. R. aculeis geminatis: altero recurvo, floribus digynis, foliis ovato-oblongis. *Syſt. veget. p.* 235. *Pallas roſſic. t.* 59. *Scop. carn. tom.* 1. *p.* 165.

Rhamnus aculeis geminis ſubæqualibus, foliis oblique ovatis trinerviis ſerratis, pedunculis confertis. *Gouan hort. p.* 113.

Le Iujubier. Regn. bot.

Occurrit in Forojulio, Iſtria, Litorali in hortis, & vineis neglectis. Fl. Iulio. Arb.

EVONYMUS. Perianthium quadri- ſeu quinquefidum. Petala, & ſtamina quatuor ſeu quinque. Capſula penta- vel tetragona, quinque- ſeu quadrilocularis, in quovis loculamento recondens ſemen unicum, arillo baccato obvolutum.

latifolius. 1. E. floribus plerisque pentandris, cortice lævi, pedunculis filiformibus teretibus multifloris. *Ait. Kew. tom.* 1. *p.* 273. *Syſt. veget. p.* 238. *Iacqu. auſtr. t.* 289. *Scop. carn. tom.* 1. *p.* 165.

Evonymus pedunculis lateralibus patulis, petalis ſubrotundis, fructibus alatis. *Scop. carn. ed.* 1. *p.* 325. *n.* 3. *Hall. hiſt. n.* 830.

Evonymus foliis acute, & argute ſerratis, capſularum angulis acutis. *Gouan. hort. p.* 113.

Evonymus primus latifolius. *Clus. hiſt. p.* 56.

In Auſtriæ, Styriæ, Carnioliæ, Croatiæ, & Pannoniæ ſubalpinis. Fl. Iunio. Fr.

europæus. 2. E. floribus plerisque tetrandris, pedunculis compreſſis multifloris, ſtigmatibus ſubulatis, foliis glabris. *Ait. Kew. tom.* 1. *p.* 273. *Syſt. veget. p.* 238..

Evonymus pedunculis ſolitariis, petalis oblongis, fructibus apteris. *Scop. carn. ed.* 1. *p.* 324. *n.* 1.

Evonymus foliis obtuſe ſerratis, capſularum angulis obtuſis. *Gouan. hort. p.* 113.

Evonymus foliis ovato-lanceolatis ſerratis, ramis tetragonis. *Hall. hiſt. n.* 829.

Evonymus vulgaris. *Clus. hiſt. p.* 57. *Scop. carn. tom.* 1. *p.* 166.

Ad ſepes, & in ſylvis. Fl. Majo. Fr.

verrucoſus. 3. E. floribus tetrandris, cortice verrucoſo, pedunculis filiformibus teretibus ſubtrifloris. *Ait. Kew. tom.* 1. *p.* 273. *Iacqu. auſtr. t.* 49. *Syſt. veget. p.* 238. *Scop. carn. tom.* 1. *p.* 166.

Evonymus pedunculis lateralibus; petalis ſubrotundis, fructibus apteris. *Scop. carn. ed. 1. p.* 324. *n.* 2.

Evonymus ſecundus. *Clus. hiſt. p.* 57.

In ſylvis montoſis. Fl. Majo. Fr.

RIBES. Petala, & ſtamina quinque calycis margini inſerta. Stylus bifidus. Bacca infera, polyſperma.

rubrum. 1. R. inerme, racemis glabris pendulis, floribus planiusculis. *Sp. pl. p.* 290. *Fl. dan. t.* 967.

Ribes inerme foliis planiusculis, ſtipulis minimis. *Hall. hiſt. n.* 818.

Le Groſeiller à Grappes & fruit rouge. Regn. bot.

Ad pagos. Fl. Aprili, Majo. Fr.

petræum. 2. R. inerme, racemis florentibus erectis, fructeſcentibus pendulis. *Syſt. veget. p.* 243. *Wulfen in Iacqu. miscell. tom.* 2. *p.* 36. *ic. t.* 49. *Retzii prodr. p.* 60. *Fl. dan. t.* 968.

In Carinthiæ alpibus reichenavienſibus ſupra lacum *Petaunſee* ſub rupibus. In Styriæ alpibus ad lacum *Tieſingſee* intra rupes, ſub *Winterthal* & *Turnau*, magisque infra verſus *Bergenhütten* ad petras prope caſas equiſonum; denique qua ex *Turach* verſus *Stangalpen* itur. *Wulfen.* In Carniolia in rupeſtribus circa Idriam, alibique. Fl. Iunio, Iulio. Fr.

alpinum. 3. R. inerme, racemis erectis, bracteis flore longioribus. *Sp. pl. p.* 291. *Iacqu. auſtr. t.* 47.

Ribes inerme floribus planis, ſtipulis florum longitudine. *Hall. hiſt. n.* 817.

Ribes vulgare dulci fructu. *Clus. hiſt. p.* 120.

In alpinis, ac ſubalpinis. Fl. Iunio. Fr.

Dantur individua, in quibus flores carent germine, qui cum bracteis decidunt omnes. Iacqu. l. c.

Groſſularia. 4. R. ramis aculeatis, petiolorum ciliis piloſis, baccis hirſutis. *Sp. pl. p.* 291.

Groſſularia Uva. *Scop. carn. tom.* 1. *p.* 168.

In Styria, Carinthia, Carniolia ad vias, montium pedes, in ſubalpinis. Fl. Iunio. Fr.

Uva criſpa. 5. R. ramis aculeatis, baccis glabris, pedicellis bractea monophylla. *Sp. pl. p.* 292. *Fl. dan. t.* 546.

Ribes ramis aculeatis, foliis rotunde lobatis. *Hall. hiſt. n.* 820.

In Auſtria ad ſepes. Fl. Aprili, Majo. Fr.

HEDERA. Petala quinque, oblonga. Pomum quinqueloculare: loculis monoſpermis. Semina arillata. *Ehrh. beytr. tom.* 4. *p.* 165.

Helix. 1. H. foliis ovatis lobatisque. *Sp. pl. p.* 292. *Curtis lond. Scop. carn. tom.* 1. *p.* 169.

Hedera foliis ſterilibus trilobatis, fructiferis ovato-lanceolatis. *Hall. hiſt. n.* 826.

La Lierre. Regn. bot.

In ſylvis, ad arborum truncos; muros, maceries. Fl. Auguſto, Septembri. Fr.

In Liburnia, vicinisque locis ruricolarum quidam e ligno cochlearia condunt; folia hyemis tempore capris, & ovibus, alio deficiente cibo, propinant. Flores ab apibus frequentantur. Turdus Merula, in Litoroli hybernans, fructibus victitat.

VITIS. Petala apice cochærentia, caduca. Bacca pentaſperma, ſupera.

vinifera. 1. V. foliis lobatis ſinuatis nudis. *Sp. pl. p.* 293. *Iacqu. collect. tom.* 1. *p.* 160. *ic. t.* 50.

Vitis foliis ſemitrilobatis, hirſutis, dentatis. *Hall. hiſt. n.* 825.

Vitis Labrusca. *Scop. carn. tom.* 1. *p.* 169.

La Vigne. Regn. bot.

In Pannonia, Auſtria, aliisque provinciis viniferis ad ſepes, in ſylvis demiſſis. Fl. Aprili, Majo. Fr.

ILLECEBRUM. Calyx pentaphyllus, cartilagineus. Corolla nulla. Stigma ſimplex. Capſula quinquevalvis, monoſperma.

Paronychia. 1. I. floribus bracteis nitidis obvallatis, caulibus procumbentibus. *Sp. pl. p.* 299.

Illecebrum Kapela. *Hacquet. pl. alp. t.* 2. *f.* 1.

Paronychia hiſpanica. *Clus. hiſt. p. CLXXXII.*

In Litorali in montibus circa Flumen Seniam, Grobnicum. Fl. Iulio, Auguſto. Per.

GLAUX. Calyx monophyllus. Corolla nulla. Capſula unilocularis, quinquevalvis, pentaſperma.

maritima. 1. Glaux. *Sp. pl. p.* 301. *Fl. dan. t.* 549.

Glaux foliis elliptico-oblongis. *Fl. lapp. n.* 72.

In arenoſis Tergeſti *alle Saule. I. Hoſt.* Fl. Majo, Iunio. Per.

THESIUM. Calyx monophyllus, cui Stamina inſerta. Semen unicum, inferum.

Linophyllum. 1. T. panicula foliata, foliis lineari-lanceolatis. *Sp. pl. p.* 301. *Scop. carn. tom.* 1. *p.* 113.

Theſium caule erecto, paniculato, foliis lanceolatis. *Hall. hiſt. n.* 1573.

Anonymos Lini folio. *Cluſ. hiſt. p.* 324.

In pratis ſiccioribus, collibus. Fl. Majo, Iunio. Per.

alpinum. 2. T. racemo foliato, foliis linearibus. *Sp. pl. p.* 301. *Iacqu. auſt. t.* 416. *en. p.* 213. *Scop. carn. tom.* 1. *p.* 114.

Theſium caule diffuſo, floribus alaribus, foliis linearibus. *Hall. hiſt. n.* 1574.

In montanis, ſubalpinis, alpinis. Fl. æſtate. Per.

VINCA. Folliculi duo, erecti. Semina nuda.

minor. 1. V. caulibus procumbentibus, foliis lanceolato-ovatis, floribus pedunculatis. *Sp. pl. p.* 304. *Curtis lond.*

Pervinca caulibus procumbentibus, foliis ovato-lanceolatis, petiolis unifloris. *Hall. hiſt. n.* 572.

Pervinca minor. *Scop. carn. tom.* 1. *p.* 170.

Le petite Pervenche. Regn. bot.

In montanis ſterilibus, alibique. Fl. primo vere. Per.

Vel folia baſi, vel petioli apicem verſus tam in hac, quam ſequente ſpecie ſunt utrinque denticulo notati.

major. 2. V. caulibus erectis, foliis ovatis ciliatis. *Sibthorp. oxon. p.* 79. *Sp. pl. p.* 304. *Wulſen in Iacqu. collect. tom.* 2. *p.* 210.

Pervinca caulibus erectis, foliis ovato-lanceolatis, ciliatis, petiolis unifloris. *Hall. hiſt. n.* 573.

Pervinca major. *Scop. carn. tom.* 1. *p.* 170.

Clematis Daphnoides latifolia. *Cluſ. hiſt. p.* 121.

La grande Pervenche. Regn. bot.

In Forojulio in agro goritienſi ad ſepes. Fl. Martio, Aprili. Per.

Folia cordato-ovata, ciliata. Calyx longitudine piſtilli: laciniis biglanduloſis, ciliatis. Ehrh. beytr. tom. 5. p. 45. *tom.* 6. *p.* 105.

DIGYNIA.

ASCLEPIAS. Calyx quinquefidus. Corolla quinquepartita. Filamenta in tubum connexa. Antheræ membrana ſtigmati incumbente terminatæ, utrinque ala auctæ. Nectaria quinque, tubo filamentorum intra antheras adnata, e fundo corniculum exerentia. Folliculi duo. Semina pappo coronata.

Vincetoxicum. 1. A. foliis cordato-ovatis baſi ſubpiloſis, caule erecto, floribus faſciculatis. *Wild. prodr. p.* 93. *Sp. pl. p.* 314. *Fl. dan. t.* 498. *Scop. carn. tom.* 1. *p.* 171.

Asclepias ex alis racemoſa, caulibus ſimplicibus erectis, foliis ovato-lanceolatis. *Hall. hiſt. n.* 571.

Le Domte-venin. Regn. bot.

In fruticoſis, montoſis aſperis. Fl. Iunio, Iulio. Per.

HERNIARIA. Calyx quinquepartitus. Corolla nulla. Stamina decem; quorum quinque intra calycis lacinias, & fertilia, alia quinque ſterilia. Capſula calyce perſiſtente tecta, monoſperma.

glabra. 1. H. glabra. *Syſt. veget. p.* 261. *Fl. dan. t.* 529. *Scop. carn. tom.* 1. *p.* 172.

Herniaria glabra, glomerulis multifloris. *Hall. hiſt. n.* 1552.

La Turquette. Regn. bot.

In arenoſis. Fl. æſtate.

hirſuta. 2. H. hirſuta. *Syſt. veget. p.* 261.

Herniaria hirſuta, glomerulis paucifloris. *Hall. hiſt. n.* 1553.

Herniaria hirſuta. *Bauh. hiſt.* 3. *p.* 379. *Zann. iſt. p.* 138. *t.* 254.

In Litorali in arenoſis ad mare; in Tyroli ad fluvios. Fl. Iunio, Iulio.

CHENOPODIUM. Calyx pentaphyllus, pentagonus. Corolla nulla. Semen unicum, lenticulare, ſuperum.

* *Foliis angulatis.*

Bonus Henricus. 1. C. foliis triangulari-ſagittatis integerrimis, ſpicis compoſitis aphyllis axillaribus. *Syſt. veget. p.* 261. *Curtis lond. Fl. dan. t.* 579. *Scop. carn. tom.* 1. *p.* 172.

Chenopodium foliis triangularibus, undulatis, integerrimis, ſubtus farinoſis. *Hall. hiſt. n.* 1578.

Le Bon Henri. Regn. bot.

In ruderatis, ad pagos. Fl. æſtate. Per.

murale. 2. C. foliis ovatis nitidis dentatis acutis, racemis ramoſis nudis. *Sp. pl. p.* 318. *Curtis lond.*

In cultis, ruderatis, muris hortorum vetuſtis. Fl. Iulio, Auguſto.

viride. 3. C. foliis rhomboideis dentato-ſinuatis, racemis ramoſis ſubfoliatis. *Sp. pl. p.* 319. *Curtis lond. Retzii prodr. p.* 62.

In cultis, ruderatis, ad vias. Fl. æſtate.

Cum Chenopodio viridi conjungunt C. album Cl. Viri, Retzius, Mœnchius.

hybridum. 4. C. foliis cordatis angulato-acuminatis, racemis ramoſis nudis. *Sp. pl. p.* 319. *Curtis lond.*

Chenopodium foliis glabris ſeptangulis, floribus paniculatis. *Hall. hiſt. n.* 1581.

In cultis, & alibi. Fl. æſtate.

Botrys. 5. C. foliis oblongis ſinuatis, racemis nudis multifidis. *Sp. pl. p.* 320. *Blakw. herb. t.* 314.

Chenopodium foliis oblongis, ſemipinnatis, viscidis, rotunde dentatis. *Hall. hiſt. n.* 1585.

In arenoſis Auſtriæ, Tyrolis. Fl. æſtate.

glaucum. 6. C. foliis ovato-oblongis repandis, racemis nudis ſimplicibus glomeratis. *Sp. pl. p.* 320.

Chenopodium foliis oblongis, ſinuatis, ſubtus incanis. *Hall. hiſt. n.* 1584.

Atriplex anguſtifolia laciniata. *Bauh. hiſt.* 2. *p.* 472. *t.* 473.

In cultis, ad fimeta, vias. Fl. Iulio, Auguſto, Septembri.

** *Foliis integris.*

olidum. 7. C. foliis integerrimis rhombeo-ovatis, floribus conglomeratis axillaribus. *Curtis lond.*

Chenopodium caule diffuſo, foliis obtuſe lanceolatis. *Hall. hiſt. n.* 1577.

Chenopodium Vulvaria. *Sp. pl. p.* 321. *Scop. carn. tom.* 1. *p.* 174.

L'Arroche fetide. Regn. bot.

In cultis, ruderatis. Fl. æſtate.

polyſpermum. 8. C. foliis integerrimis ovatis, caule decumbente, cymis dichotomis aphyllis axillaribus. *Sp. pl. p.* 321. *Curtis lond. Scop. carn. tom.* 1. *p.* 173.

Chenopodium caule erecto, foliis ovatis integris. *Hall. hiſt. n.* 1576.

In cultis, ad ſepes humidas, foſſas, & in paluſtribus. Fl. æſtate.

maritimum. 9. C. foliis ſubulatis ſemicylindricis. *Sp. pl. p.* 321. *Fl. dan. t.* 489. *Scop. carn. tom.* 1. *p.* 174.

In Litorali in arenoſis ad mare. In Pannoniæ comitatu jaurinenſi in ſterilibus circa *Sövényháza.* Fl. Auguſto.

BETA. Calyx pentaphyllus. Corolla nulla. Semen reniforme, intra ſubſtantiam baſeos calycis.

maritima. 1. B. floribus geminis. *Syſt. veget. p.* 262.

Beta ſylveſtris maritima. *Bauh. pin.* 118. *Zann. iſt. p.* 36.

In Litorali in paluſtribus aquilejenſibus ad fluvium *Anfora.* Fl. Auguſto.

SALSOLA. Calyx pentaphyllus. Corolla nulla. Capſula monoſperma. Semen cochleatum.

Tragus. 1. S. herbacea erecta, foliis ſubulatis ſpinoſis lævibus, calycibus ovatis. *Sp. pl. p.* 322.

Kali Tragus. *Scop. carn. tom.* 1. *p.* 175.

In collibus, arvis, ad fluvios. Fl. æſtate.

Soda. 2. S. herbacea patula, foliis inermibus. *Sp. pl. p.* 323. *Iacqu. hort. tom.* 1. *p.* 28. *t.* 68.

Kali magnum, ſedi medii folio, ſemine cochleato. *Lob. ic. p.* 394.

Kali majus cochleato ſemine. *Bauh. pin.* 289.

La Soude. Regn. bot.

In paluſtribus aquilejenſibus. Fl. Auguſto.

ſativa. 3. S. herbacea diffuſa, foliis teretibus glabris, floribus conglomeratis. *Sp. pl. p.* 323.

Habitat & floret cum præcedente.

proſtrata. 4. S. fruteſcens, foliis linearibus piloſis inermibus. *Sp. pl. p.* 223. *Iacqu. auſtr. t.* 294.

In Auſtria circa oppidum. *Rœtz.* Fl. a Iulio ad Septembrim.

ULMUS. Calyx quinquefidus. Corolla nulla. Capſula monoſperma, alata. *Wild. prodr. p.* 94.

campeſtris. 1. U. foliis duplicato-ſerratis, baſi inæqualibus. *Sp. pl. p.* 327. *Fl. dan. t.* 632. *Scop. carn. tom.* 1. *p.* 176.

Ulmus foliis ſcabris, ovato-lanceolatis, dentibus ſerratis. *Hall. hiſt. n.* 1586.

In hortis, ſylvis, ad ſepes, pagos, alibique. Fl. Martio, Aprili. Arb.

Folia cocta cum pauca furfurum, vel farinæ quantitate pro cibo ſuibus dant Iſtriæ incolæ.

SWERTIA. Corolla rotata. Nectariferi pori ad baſin laciniarum corollæ Capſula unilocularis, bivalvis.

perennis. 1. S. corollis quinquefidis, foliis radicalibus ovalibus. *Sp. pl. p.* 328. *Iacqu. auſtr. t.* 243.

Gentiana foliis radicalibus petiolatis, ovatis, floribus paniculatis, rotatis, ſpicatis. *Hall. hiſt. n.* 636.

Gentiana duodecima punctato flore. *Cluſ. hiſt. p.* 316.

In Auſtria ad rivulos, & aliis udis locis montis *Oetſcherberg*; in Styria in alpibus judenburgenſibus. Fl. æſtate. Per.

GENTIANA. Corolla monopetala. Capſula bivalvis, unilocularis. Receptacula duo, ſingula valvulæ longitudinaliter adnata.

lutea. 1. G. corollis ſubquinquefidis rotatis verticillatis, calycibus ſpathaceis. *Sp. pl. p.* 329. *Scop. carn. tom.* 1. *p.* 182.

Gentiana caule folioſo, foliis ovatis, nervoſis, floribus verticillatis, rotatis. *Hall. hiſt. n.* 637.

Gentiana veterum. *Cluſ. hiſt. p.* 311.

Gentiana major lutea. *Bauh. pin.* 187.

La grande Gentiane. Regn. bot.

In alpinis, & ſubalpinis Carnioliæ, Forijulii, Litoralis, Croatiæ. Fl. Iulio, Auguſto. Per.

Equi ſpernunt folia, ſpicas contra floribus onuſtas libentiſſime vorant.

campanulata. 2. G. corollis campanulatis ſeptemfidis, ſtriatis, immaculatis, calycibus ſeptemfidis. *Syſt. veget. p.* 267. *Iacqu. auſtr. app. t.* 29.

In Carinthiæ alpe, *Garten* dicta, in *Reichenau. Wulfen.* Fl. Iulio. Per.

punctata. 3. G. corollis ſubquinquefidis campanulatis punctatis, calycibus quinquedentatis. *Syſt. veget. p.* 267. *Iacqu. auſtr. app. t.* 28.

Gentiana foliis ovato-lanceolatis, nervoſis, floribus campaniformibus, fasciculatis, creberrime punctatis. *Hall. hiſt. n.* 638.

Gentiana major pallido flore, punctis diſtincto. *Cluſ. hiſt. p.* 312.

In alpibus Styriæ, Carinthiæ, Carniæ, Tyrolis. Fl. Iulio, Auguſto. Per.

algida. 4. G. corollis quinquefidis campanulatis punctatis terminalibus ſeſſilibus, foliis radicalibus faſciculatis. *Pallas it.* 3. *app. p.* 724. *t.* 50. *Fl. roſſic. t.* 95.

Gentiana floribus terminantibus diaphanis. *Gmel. ſib.* 4. *p.* 106. *n.* 73.

Gentiana frigida. *Hænke in Iacqu. collect. tom.* 2. *p.* 13.

In Styria ſuperiore in abruptis præcipitiis alpium ſeckavienſium. *Hænke.* In Pannonia in montibus carpaticis. *Iacquin.* Fl. Iulio. Per.

pannonica. 5. G. corollis campanulatis ſex- vel ſeptemfidis punctatis, calycibus ſubſexfidis, foliis caulinis lanceolatis acuminatis. *Syſt. veget. p.* 267. *Iacqu. auſtr. t.* 136. *Scop. carn. tom.* 1. *p.* 182.

Gentiana major purpureo flore prima. *Cluſ. hiſt. p.* 312.

In alpinis, & ſubalpinis. Fl. Auguſto, Septembri. Per.

Hæc ſpecies habet calycem campanulatum, obſolete hexagonum, circa medietatem in lacinias tot, quot ſunt corollæ laciniæ, duas oppoſitas majores, diviſum. Diverſa ergo a Gentiana purpurea, cui calyx ſpathaceus, bivalvis. Iacqu. l. c.

aſclepiadea. 6. G. corollis quinquefidis campanulatis oppoſitis ſeſſilibus, foliis amplexicaulibus. *Sp. pl. p.* 329. *Iacqu. auſtr. t.* 328. *Scop. carn. tom.* 1. *p.* 181.

Gentiana floribus campaniformibus, alis paucifloris, foliis ovato-lanceolatis. *Hall. hiſt. n.* 640.

Gentiana ſecunda cœruleo flore. *Cluſ. hiſt. p.* 312.

In ſubalpinis, alpinis. Fl. Auguſto, Septembri. Per.

Pneumonanthe. 7. G. corollis quinquefidis campanulatis oppoſitis pedunculatis, foliis linearibus. *Sp. pl. p.* 330. *Fl. dan. t.* 269. *Scop. carn. tom.* 1. *p.* 181.

Gentiana alis floriferis, foliis linearibus. *Hall. hiſt. n.* 641.

Gentianæ quarta ſpecies. *Clus. hiſt. p.* 313.

In pratis humidis. Fl. Iulio, Auguſto, Septembri. Per.

Variat foliis lanceolatis, & ovatis; obtuſis, ac acutis.

acaulis. 8. G. corolla quinquefida campanulata caulem excedente. *Sp. pl. p.* 330. *Iacqu. auſtr. t.* 135. *Scop. carn. tom.* 1. *p.* 180.

Gentiana caule unifloro, foliis lanceolatis, flore maximo, campaniformi. *Hall. hiſt. n.* 642.

Gentiana quinta, ſive Gentianella major verna. *Clus. hiſt. p.* 314.

Gentiana alpina latifolia, magno flore. *Bauh. pin.* 187. *prodr.* 97.

In ſummis alpium jugis. Fl. Iulio, Auguſto. Per.

verna. 9. G. corolla quinquefida infundibuliformi caulem excedente, foliis radicalibus confertis majoribus. *Sp. pl. p.* 331. *Scop. carn. tom.* 1. *p.* 178.

Gentiana foliis ovato-lanceolatis, caule unifloro. *Hall. hiſt. n.* 644.

Gentiana ſexta, ſive Gentianella minor verna. *Clus. hiſt. p.* 314.

In pratis montanis, ſubalpinis, & paſcuis alpinis. Fl. pro diverſa elevatione ab Aprili ad Auguſtum. Per.

pumila. 10. G. corolla quinquefida infundibuliformi ſubſerrata, foliis lanceolato-linearibus, *Syſt. veget. p.* 268. *Iacqu. auſtr. t.* 302.

Gentiana undecima minima. *Clus. hiſt. p.* 316.

In ſummis alpium jugis. Fl. æſtate.

bavarica. 11. G. corolla quinquefida infundibuliformi ſerrata, foliis ovatis obtuſis. *Sp. pl. p.* 331.

Gentiana foliis ovatis, caule unifloro. *Hall. hiſt. n.* 645.

Gentianella elegantiſſima. *Cam. hort. ic.* 65. *t.* 15. *f.* 2.

In alpibus ſtyriacis, carinthiacis, tyrolenſibus. Fl. Iulio, Auguſto. Per.

nivalis. 12. G. corollis quinquefidis infundibuliformibus, ramis unifloris alternis. *Sp. pl. p.* 332. *Fl. dan. t.* 17.

Gentiana caule ramoſo, foliis ovato-lanceolatis, floribus infundibuliformibus. *Hall. hiſt. n.* 647.

Gentiana nana. *Wulfen in Iacqu. miscell. tom.* 1. *p.* 161. *t.* 18. *f.* 3. *Syſt. veget. p.* 267.

In alpinis, & ſubalpinis. Fl. Iulio, Auguſto.

In ſummis carniolicarum alpium jugis occurrit caule breviſſimo, ſimplici, unifloro. Conjunxi G. nanam Wulfenii cum nivali Linnæi: nam Cl. Allionius in Auctario ad Floram pedemontanam dicit: „Collatis multis ſpeciminibus Gentianæ nivalis eodem loco collectis, video G. nanam Iaquinii non eſſe diſtinguendam a nivali. Linn.“

utriculoſa. 13. G. corollis quinquefidis hypocrateriformibus, calycibus plicato-alatis. *Sp. pl. p.* 332. *Scop. carn. tom.* 1. *p.* 179.

Gentiana caule ramoſo, calycibus alatis. *Hall. hiſt. n.* 646.

In alpinis, & ſubalpinis. Fl. æſtate.

Variat caule ſimplici, unifloro, & ramoſo.

Centaurium. 14. G. corollis quinquefidis infundibuliformibus, caule dichotomo, piſtillo ſimplici. *Syſt. veget. p.* 268. *Fl. dan. t.* 617. *Scop. carn. tom.* 1. *p.* 180.

Gentiana caule dichotomo, floribus infundibuliformibus, ſtrictis. *Hall. hiſt. n.* 648.

Chironia Centaurium. *Curtis lond.*

In ſylvis, pratis. Fl. æſtate.

Amarella. 15. G. corollis quinquefidis hypocrateriformibus fauce barbatis. *Sp. pl. p.* 334. *Fl. dan. t.* 328.

Gentiana faucibus barbatis, calycis ſegmentis quinis æqualibus. *Hall. hiſt. n.* 651.

In pratis demiſſis, montanis, ſubalpinis, alpinisque, nec non in collibus herbidis. Fl. a Iulio ad ſerum Autumnum.

campeſtris. 16. G. corollis quadrifidis fauce barbatis. *Sp. pl. p.* 334. *Fl. dan. t.* 367.

Gentiana faucibus barbatis, calycis foliis quaternis, alterne majoribus. *Hall. hiſt. n.* 650.

Habitat & floret cum præcedente.

tenella. 17. G. corollis quadrifidis fauce barbatis, caule filiformi tetragono, foliis ovatis baſi ſolutis. *Retzii prodr. p.* 64. *Fl. dan. t.* 318.

Swertia carinthiaca. *Wulfen in Iacqu. miscell. tom.* 2. *p.* 53. *t.* 6.

In Tyroli in Groſskirchheimii alpibus, ſupra *Sagritz* & *Döllach* in altiſſimis alpium jugis apricis lapidoſis inter *Mocker* & *Waſchgang. Wulfen.* In Carnia in ſummis montis *Croſtis* jugis. *I. Hoſt.* Fl. Iulio, Auguſto.

Variat corollis & calycibus quinquefidis, quadrifidisque.

ciliata. 18. G. corollis quadrifidis margine ciliatis. *Sp. pl. p.* 334. *Iacqu. auſtr. t.* 113. *Scop. carn. tom.* 1. *p.* 176.

Gentiana flore quadrifido, oris ciliatis. *Hall. hiſt. n.* 653.

In herbidis montanis, & ſubalpinis. Fl. Auguſto, Septembri, Octobri.

cruciata. 19. G. corollis quadrifidis imberbibus, floribus verticillatis ſeſſilibus. *Sp. pl. p.* 334. *Iacqu. auſtr. t.* 372. *Scop. carn. tom.* I. *p.* 177.

Gentiana foliis ovato-lanceolatis, nervoſis, vaginalibus, floribus verticillatis & umbellatis. *Hall. hiſt. n.* 643.

Gentiana tertia, cruciata. *Cluſ. hiſt. p.* 313.

In collibus herbidis, pratis montanis. Fl. Auguſto, Septembri. Per.

ERYNGIUM. Flores capitati, Receptaculum paleaceum.

planum. 1. E. foliis radicalibus ovalibus planis crenatis, capitulis pedunculatis. *Sp. pl. p.* 336. *Iacqu. auſtr. t.* 391.

Eryngium pannonicum latifolium. *Cluſ. hiſt. p.* CLVII.

Eryngium latifolium planum. *Bauh. pin.* 386.

In Auſtria, Pannonia in herbidis, ceſpitoſis ad Danubium & Leitham, alibique. Fl. a Iunio ad Auguſtum. Per.

maritimum. 2. E. foliis radicalibus ſubrotundis plicatis ſpinoſis, capitulis pedunculatis, paleis tricuſpidatis. *Sp. pl. p.* 337. *Fl. dan. t.* 875. *Scop. carn. tom.* I. *p.* 185.

Eryngium marinum. *Cluſ. hiſt. p.* CLIX. *Zann. iſt. p.* 95. *t.* 94.

Infrequens in arenoſis Litoralis. Fl. æſtate. Per.

campeſtre. 3. E. foliis amplexicaulibus pinnato-laciniatis. *Sp. pl. p.* 337. *Iacqu. auſtr. t.* 155. *Fl. dan. t.* 554. *Scop. carn. tom.* I. *p.* 185.

Eryngium foliis ſemipinnatis, pinnis pinnatis, pinnulis lobatis, ſpinoſis. *Hall. hiſt. n.* 735.

Eryngium campeſtre vulgare. *Cluſ. hiſt. p.* CLVII.

Le Panicaut. Regn. bot.

In campis, collibus herbidis. Fl. æſtate. Per.

Folia, quæ primis annis prodeunt, ſunt oblonga, apice magis minusve trifida, aut quinquefida. In florente planta ſunt pleraque ternata: foliolis bipinnatis; pinnulis oblongis, mucronatis, per coſtam decurrentibus. Involucrum pentaphyllum mucronatum; foliolis ſubenſiformibus, patentiſſimis, capituloque duplo, triplove longioribus. Iacqu. l. c.

alpinum. 4. E. foliis radicalibus cordatis oblongis, caulinis pinnatifidis, capitulis ſubcylindricis. *Sp. pl. p.* 337. *Iacqu. ic. t.* 55. *Scop. carn. tom.* 1. *p.* 184.

Eryngium foliis radicalibus petiolatis, cordatis, involucro pinnato, ciliato. *Hall. hiſt. n.* 736.

In Carniolia in alpibus wochinenſibus, tolmienſibusque. Fl. Iulio, Auguſto. Per.

SANICULA. Umbellæ confertæ, ſubcapitatæ. Fructus ſcaber. Flores disci abortientes.

europæa. 1. S. foliis radicalibus ſimplicibus, floſculis omnibus ſeſſilibus. *Sp. pl. p.* 339. *Fl. dan. t.* 283.

Sanicula foliis radicalibus trilobatis, lateralibus lobis tripartitis, umbellis globoſis. *Hall. hiſt. n.* 737.

Sanicula vulgaris. *Clus. hiſt. p.* 308.

Aſtrantia Diapenſia. *Scop. carn. tom.* 1. *p.* 186.

Caucalis Sanicula officinalis. *Crantz faſc. p.* 228.

In ſylvis montoſis. Fl. Majo, Iunio. Per.

ASTRANTIA. Involucra partialia lanceolata patentia, æqualia, longiora, colorata. Flores plurimi abortientes.

major. 1. A. foliis quinquelobis: lobis trifidis. *Sp. pl. p.* 339. *Hall. hiſt. n.* 790.

Aſtrantia floribus pedunculatis, ſeminum angulis denticulatis. *Scop. carn. ed.* 1. *p.* 337. *n.* 3.

Aſtrantia nigra. *Blakw. herb. t.* 470. *Scop. carn. tom.* 1. *p.* 188.

Helleborus niger, ſaniculæ folio. *Bauh. pin.* 186.

In ſubalpinis, & alpinis; occurrit quoque ad margines ſylvarum montoſarum. Fl. æſtate. Per.

minor. 2. A. foliis digitatis ſerratis. *Sp. pl. p.* 340. *Hall. hiſt. n.* 791. *Scop. carn. tom.* 1. *p.* 187. *t.* 7.

Aſtrantia floribus pedunculatis, ſeminum angulis alato-plicatis. *Scop. carn. ed.* 1. *p.* 337. *n.* 4.

Aſtrantia carniolica. *Iacqu. auſtr. app. t.* 10.

Helleborus niger, ſaniculæ facie, minor. *Bauh. pin.* 186. *prodr.* 97.

In Carniolia in ſylveſtribus circa Idriam; *in der Wochein* locis alpinis, & ſubalpinis. In Carinthia in monte *Loibl.* Fl. æſtate. Per.

In planta in demiſſioribus creſcente folia partita obſervavi. Specimina contra, quæ in ſummis alpium jugis apricis legi, habent folia digitata, foliis Aſtrantiæ minoris helveticæ perfecte ſimilia.

Epipactis. 3. A. foliis quinquepartitis obtuſis ſerratis. *Syſt. veget. p.* 273. *Scop. carn. tom.* 1. *p.* 185. *t.* 6. *Iacqu. auſtr. app. t.* 11.

Aſtrantia floribus maſculis pedunculatis; femineis ſeſſilibus; ſeminum angulis obtuſis, glabris. *Scop. carn. ed.* 1. *p.* 337. *n.* 2.

Sanicula involucris maximis quinquefoliis, ſeminibus tuberculoſis. *Hall. nom. n.* 737.

In ſylvis montanis, & ſubalpinis Carinthiæ, Carnioliæ, Forijulii, Croatiæ, Pannoniæ. Fl. a Martio ad Iunium. Per.

BUPLEURUM. Involucra umbellulæ majora, pentaphylla. Petala involuta. Fructus ſubrotundus, compreſſus, ſtriatus.

rotundifolium. 1. B. involucris univerſalibus nullis, foliis perfoliatis. *Sp. pl. p.* 340. *Scop. carn. tom.* 1. *p.* 209.

Bupleurum caule brachiato, ramoſiſſimo, foliis ovato-lanceolatis, amplexicaulibus. *Hall. hiſt. n.* 767.

La Perce-Feuille. Regn. bot.

Inter ſegetes, alibique. Fl. Iunio, Iulio.

petræum. 2. B. involucellis coadunatis, univerſali pentaphyllo. *Sp. pl. p.* 340. *Wulfen in Iacqu. collect. tom.* 1. *p.* 209. *ic. t.* 56.

Bupleurum alpinum foliis anguſtis gramineis ſeſſilibus, caule fere nudo. *Segu. ver.* 2. *p.* 15.

Sedum petræum, Bupleuri folio. *Pon. in Cluſ. hiſt. p.* CCCXLVII.

In Carnioliæ alpibus wochinenſibus, lithopolitanisque, innaſcitur rupium fiſſuris. Fl. Iulio, Auguſto. Per.

falcatum. 3. B. involucellis pentaphyllis acutis, univerſali ſubpentaphyllo, foliis lanceolatis, caule flexuoſo. *Sp. pl. p.* 341. *Iacqu. auſtr. t.* 158. *Scop. carn. tom.* 1. *p.* 210.

Bupleurum foliis radicalibus petiolatis, elliptico-lanceolatis, caulinis falcatis, caule flexuoſo. *Hall. hiſt. n.* 776.

In aſperis, ad ſepes, & in aliis locis. Fl. ab Auguſto ad Octobrim. Per.

Folia radicalia obverſe ovata. Caulina, & ramea ſeſſilia, obverſe lanceolata, aut oblonga, aut ſublinearia; quandoque ſubfalcata. Iacqu. l. c.

odontites. 4. B. involucellis pentaphyllis acutis; univerſali triphyllo, floſculo centrali altiore, ramis divaricatis. *Sp. pl. p.* 342. *Iacqu. hort. tom.* 3. *p.* 47. *t.* 91. *Scop. carn. tom.* 1. *p.* 210.

Bupleurum caule brachiato, involucris utrisque pentaphyllis, ariſtatis, petiolos excedentibus. *Hall. hiſt. n.* 772.

In Carnioliæ calidioris, Forijulii, agri fluminenſis herbidis, aſperisque locis. Fl. Iulio, Auguſto.

Planta in aſperis ſaxoſis Carnioliæ prope Prewald, Vipacum *creſcens nana eſt. Pedalis contra, pedeque longior, quam herbida agri fluminenſis loca alunt.*

ſemicompoſitum. 5. B. umbellis compoſitis ſimulque ſimplicibus. *Sp. pl. p.* 342.

Bulpleurum caule herbaceo, foliis lanceolatis, umbellis axillaribus terminalibusque, ſeminibus ſcabris. *Gouan. illuſtr. p.* 9. *t.* 7. *f.* 1.

In agro tergeſtino, totaque Iſtria. Fl. Iunio, Iulio.

Gerardi. 6. B. caule erecto paniculato, involucris & involucellis pentaphyllis. *Syſt. veget. p.* 274. *Iacqu. auſtr. t.* 256.

Bupleurum involucris & involucellis pentaphyllis, foliolis lineari-ſubulatis. *Ger. prov. p.* 233. *t.* 9.

In Auſtria inferiore, & Litorali ad ſepes, vineas. Fl. Iulio, Auguſto.

junceum. 7. B. caule erecto paniculato, foliis linearibus, involucris triphyllis, involucellis pentaphyllis. *Sp. pl. p.* 343.

In agro fluminenſi ad ſepes, in fruticoſis. Fl. Iulio, Auguſto.

Vel ſolo caule ſexpedali, altioreque diſtinguitur facillime a reliquis ſpeciebus.

baldenſe. 8. B. involucro univerſali tetraphyllo, partialibus pentaphyllis acuminatis, ramis foliisque alternis. *Syſt. nat. ed.* 13. *p.* 470. *Turra giorn. p.* 47. *t.* 1.

In Croatiæ, Litoralis, Carnioliæ calidioris montanis, & ſubalpinis. Fl. Iulio, Auguſto. Per.

ECHINOPHORA. Flores laterales masculi; centrali hermaphrodito. Semen unicum, involucello immerſum.

ſpinoſa. 1. E. foliis ſubulato-ſpinoſis integerrimis. *Sp. pl. p.* 344.

Echinophora maritima, ſpinoſa. *Tourn. inſt.* 566.

Crithmum maritimum ſpinoſum. *Bauh. pin.* 288.

Paſtinaca marina. *Lob. ic. p.* 710. *Zann. iſt. p.* 90. *t.* 15.

In arenoſis agri aquilejenſis ad mare. Fl. Iunio, Iulio. Per.

TORDYLIUM. Corollæ radiatæ omnes hermaphroditæ. Fructus ſuborbiculatus, margine crenatus. Involucra longa indiviſa.

maximum. 1. T. umbellis confertis radiatis, foliolis lanceolatis inciſo-ſerratis. *Sp. pl. p.* 345. *Iacqu. auſtr. t.* 142. *Scop. carn. tom.* 1. *p.* 195. *Crantz faſc. p.* 153.

Tordylium foliis pinnatis, pinnis ſemipinnatis, ſeminis limbo lævi. *Hall. hiſt. n.* 811.

Caucalis major. *Clus. hiſt. p.* CCI.

In Auſtriæ, Litoralis fruticoſis, ad ſepes. Fl. Majo, Iunio.

Foliorum radicalium pinnæ ſunt ſeſſiles, ſubovatæ, inæqualiter ſerrato-crenatæ, obtuſiſſimæ, extrema majore, ac ad baſin attenuata. Extimorum floſculorum petalum extimum eſt maximum, atque profundiſſime bifidum. Petalum laterale alterutrum vel utrumque laciniam priori petalo proximam etiam talem, ſed paulo minorem habet. Inque hujusmodi lacinias geminas dividuntur petala duo reliqua. Semina ſunt ſuborbicularia, compreſſo-plana. Iacqu. l. c.

Anthriscus. 2. T. umbellis confertis, foliolis ovato-lanceolatis pinnatifidis. *Sp. pl. p.* 346. *Iacqu. auſtr. t.* 261. *Crantz faſc. p.* 226.

Caucalis foliis duplicato-pinnatis, nervo multoties latioribus. *Hall. hiſt. n.* 741.

Caucalis Anthriscus. *Scop. carn. tom.* 1. *p.* 191.

Caucalis ſemine aſpero, floſculis rubentibus. *Bauh. pin.* 153. *prodr.* 80.

In ſylvis, ad ſepes. Fl. Iulio, Auguſto.

Caulis erectus. Folia ſubbipinnata: foliolis pinnatifidis aut ſerrato-inciſis, acutis, etiam mucronatis. Petala ob apicem inflexum cordata. Fructus ovatus dispescitur in duo ſemina hinc gibba, & pilis ſubulatis ſurſumque arcuatis hispida, illinc linea glabra longitudinaliter concava; ad convexas oras piloſa. Iacqu. l. c.

nodoſum. 3. T. umbellis ſimplicibus ſeſſilibus, ſeminibus exterioribus hiſpidis. *Sp. pl. p.* 346. *Iacqu. auſtr. app. t.* 24.

Caucalis nodoſa. *Scop. carn. tom.* 1. *p.* 192.

Caucalis nodoſa, echinato ſemine. *Bauh. pin.* 153.

In Forojulio, Iſtria, Litorali ad vias, vineas. Fl. Iunio, Iulio.

Folia ſunt ſubbipinnata: pinnulis laciniatis, acutis. Umbellæ ad nodos caulis ſeſſiles, oppoſitifoliæ. Involucrum univerſale nullum. Partiale polyphyllum. Petala apice inflexa. Semina hinc convexa, & pilis uncinatis hispida, inde plana. Iacqu. l. c.

CAUCALIS. Corollæ radiatæ; diſci maſculæ. Petala inflexo-emarginata. Fructus ſetis hiſpidus. Involucra integra.

grandiflora. 1. C. involucris pentaphyllis: foliolo unico duplo majore. *Sp. pl. p.* 346. *Iacqu. auſtr. t.* 54. *Crantz faſc. p.* 224.

Caucalis umbellis planis, petalis extremis & involucris maximis. *Hall. hiſt. n.* 740.

Caucalis albo flore. *Cluſ. hiſt. p.* CCI.

Caucalis arvenſis echinata, magno flore. *Bauh. pin.* 152. *Zann. iſt. p.* 54. *t.* 209.

Daucus grandiflorus. *Scop. carn. tom.* 1. *p.* 189.

In ſaxoſis aſperis, inter ſegetes in Auſtria, Forojulio, Iſtria, Litorali. Fl. Majo, Iunio.

Folia multiplicato-pinnata: pinnulis exiguis, lanceolatis, confluentibus. Involucri univerſalis foliola ſunt ut plurimum quinque. Partiale ſenis conſtat foliolis, quorum tria exteriora duplo longiora ſunt. Petalum exterius floſculorum, qui in umbellulis exterioribus externe locantur, multo majus eſt reliquis, & ferme ad baſin usque bifidum cum lacinia intermedia exigua, lineari, inflexa. Cæteris exterioribus corollulis petalum idem duplo dumtaxat majus eſt. Interiorum petala ſunt æqualia. Semina ſunt ovata, compreſſa, extus ſtriata, ac ſetis numeroſis rigidis uncinatisque armata. Iacqu. l. c.

daucoides. 2. C. umbellis trifidis aphyllis, umbellulis triſpermis triphyllis. *Syſt. veget. p.* 276. *Iacqu. auſtr. t.* 157.

Caucalis leptophylla. *Crantz faſc. p.* 225.

Daucus leptophyllus. *Scop. carn. tom.* 1. *p.* 190.

In arvis, ad vineas, vias. Fl. Iunio.

Folia multiplicato-pinnata. Involucrum quandoque nullum, quandoque unifolium ac breviſſimum. Particulare ſæpius triphyllum. Umbella univerſalis radios habet tres, vel quatuor; quarto plerumque umbellulam floſculis maſculis onuſtam

gerente. Flores a tribus ad ſeptem in qualibet umbellula, tres vel duo inter hos utplurimum hermaphroditi; reliqui mares. Semina ovata, hinc plana & ſulcata, inde ſpinis albidis uncinatis munita. Iacqu. l. c.

DAUCUS. Corollæ ſubradiatæ, omnes hermaphroditæ. Fructus pilis hiſpidus.

Carota. 1. D. ſeminibus hiſpidis, petiolis ſubtus nervoſis. *Sp. pl. p.* 348. *Fl. dan. t.* 723. *Scop. carn. tom.* 1. *p.* 189.

Daucus involucris cavis, communibus pinnatis, peculiaribus lineari-lanceolatis. *Hall. hiſt. n.* 746.

In pratis, cultis, alibique. Fl. Iunio, Iulio.

Umbella primo plana, dein hemiſphærii cavi formam induit.

CONIUM. Involucella dimidiata, ſubtriphylla. Fructus ſubgloboſus, quinqueſtriatus, utrinque crenatus.

maculatum. 1. C. ſeminibus ſtriatis. *Sp. pl. p.* 349. *Iacqu. auſtr. t.* 156. *Curtis lond. Scop. carn. tom.* 1. *p.* 206. *Hall. hiſt. n.* 766.

Cicutaria vulgaris. *Clus. hiſt. p.* CC.

Coriandrum Cicuta. *Crantz faſc. p.* 211.

La grande Ciguë. Regn. bot.

Ad ſepes rivulos, in cultis, cœmeteriis. Fl. Iunio, Iulio.

Tota planta fœtidiſſimum murium odorem ſpirat. Caulis teres, lineatus, fiſtuloſus, rore glauco tectus, virens cum plerumque adſperſis maculis ſanguineis. Folia ſunt multiplicato-pinnata: pinnulis oblongis, obtuſis, inciſo-ſerratis. Involucri univerſalis foliola circiter ſeptem ſunt, inæqualia, lanceolato-acuminata, reflexa, margine membranacea. Partiale ſimile eſt, at dimidiatum, ac patens. Petala inæqualia. Fructus compreſſo-ſubrotundus conſtat binis ſeminibus hinc planiuſculis, illinc gibbis. Iacqu. l. c.

SELINUM. Fructus ovali-oblongus, compreſſo-planus, in medio ſtriatus. Involucrum reflexum. Petala cordata, æqualia.

ſylveſtre. 1. S. radice fuſiformi multiplici, ſtylis erectis, corollulis explicatis. *Syſt. veget. p.* 278. *Iacqu. auſtr. t.* 152.

Selinum Thyſſelinum. *Crantz faſc. p.* 170.

In paluſtribus Auſtriæ prope *Himberg.* Fl. Iulio. Per.

Radix fuſiformis, læſa lac fundens. Folia multiplicato-pinnata: pinnulis linearibus, vel lanceolatis. Germinis vertex pentagonus. Semina glabra hinc planiſſima, inde convexula tribusque ſtriis elevatis notata, cum margine tenui membranaceo. Iacqu. l. c.

auſtriacum. 2. S. caule ſulcato, foliolis ovatis inciſis. *Syſt. veget. p.* 278. *Iacqu. auſtr. t.* 71.

Selinum argenteum. *Crantz faſc. p.* 174. *t.* 4. *f.* 2.

In Auſtria inferiore in fruticoſis, ſaxoſis aſperis, copioſiſſime prope *Bertholdsdorf.* Fl. Iulio, Auguſto. Per.

Folia multiplicato-pinnata: foliolis inciſis, mucrone exiguo donatis. Involucri univerſalis foliola circiter decem, lanceolato-linearia, reflexa, margine membranacea. Cum his foliola involucri partialis congruunt. Seminum nuclei ſunt elliptici, hinc plani, illinc convexuli, & tribus quatuorve ſtriis elevatis notati, excurrentes in marginem membranaceum. Iacqu. l. c.

carvifolia. 3. S. caule ſulcato acutangulo, involucro univerſali caduco, ſtylis erectis, petalis conniventibus. *Syſt. veget. p.* 278. *Iacqu. auſtr. t.* 16.

Laſerpitium ſelinoides. *Scop. carn. tom.* 1. *p.* 198.

In montoſis Auſtriæ, Carnioliæ, Fl. a Iulio ad Septembrim. Per.

Caulis membranis longitudinalibus ubique alatus. Folia multiplicato-pinnata. Petiolus communis ſtriatus: in radicalibus longus, antice ſulcatus; in caulinis dilatatus, at vix vaginans. Foliola ex oblongo elliptica, mucronata, apice albida. Involucrum univerſale conſtat foliolo unico, rariſſime pluribus, aliquando nullo. Partialis foliola a quinque ad decem. Seminum oblongi nuclei excurrunt in alam undique latam, membranaceam; & alias tres ſimiles in dorſo gerunt minutas. Iacqu. l. c.

Chabraei. 4. S. vaginis foliorum caulinorum planis flaccidis pendulis. *Syſt. veget. p.* 279. *Iacqu. auſtr. t.* 72. *Crantz faſc. p.* 162. *t.* 3. *f.* 2.

In fruticoſis agri vindobonenſis circa *Dornbach*, in *Gallitzinberg*. Fl. Auguſto, Septembri. Per.

Folia radicalia caulinaque ima ſunt compoſita ex foliolis oppoſitis pinnatis, vel tantum pinnatifidis: pinnulis lanceolato-linearibus, planis, acutis. Caulina ſuperiora conſtant foliolis minus diviſis, & longioribus. Summa ſunt ſimpliciter pinnata. Involucrum univerſale nullum. Particulare rarum quoque, unifolium & ſetaceum. Semina ſubovata & compreſſiſſima, hinc plana, illinc in medio ſtriis tribus parum elevatis notata, in marginem excurrentia membranaceum. Iacqu. l. c.

Seguieri. 5. S. involucro univerſali nullo, piſtillis divaricatiſſimis. *Syſt. veget. p.* 279. *Iacqu. hort. tom.* I. *p.* 24. *t.* 61.

Liguſticum alpinum perenne, ferulæ folio, floribus albis. *Segu. ver.* 2. *p.* 41. *t.* 13.

In montanis Carnioliæ. Fl. Iulio, Auguſto. Per.

Folia radicalia cum caulinis imis ſunt ſupradecompoſita: pinnnulis lanceolatis, apice mucrone albido terminatis. Involucrum univerſale nullum, aut conſtans foliolo uno alterove ſubulato. Partiale polyphyllum. Fructus ſubrotundo-

ovatus, conſtans ſeminibus hinc convexis & tri-vel quadriſulcatis, inde planis, angulis menbranaceis. Iacqu. l. c.

ATHAMANTA. Fructus ovato-oblongus, ſtriatus. Petala inflexa, emarginata.

Libanotis. 1. A. foliis bipinnatis planis, umbella hemisphærica; ſeminibus hirſutis. *Sp. pl. p.* 351. *Iacqu. auſtr. t.* 392.

Liguſticum foliis triplicato-pinnatis, pinnulis primis decuſſatis, ſeminibus hirſutis. *Hall. nom. n.* 757.

Libanotis Riviniana. *Scop. carn. tom.* 1. *p.* 193.

Libanotis montana. *Crantz faſc. p.* 222.

In apricis aſperis, fruticoſis, pratis montanis & ſubalpinis. Fl. Iulio, Auguſto. Per.

Folia radicalia, caulinaque inferiora ſunt pinnata aut bipinnata. pinnis ſubovatis, ſeſſilibus, ſerrato-inciſis, acutiuſculis. Pinnæ ad coſtam principem ſedent decuſſantes, quæ tamen in pluribus foliis, etiam plantis integris deſiderantur. Involucrum univerſale, & partiale conſtat foliolis pluribus, integris, ſubulatis, patentiſſimis. Radii ſunt ſtriati, villoſi. Umbella univerſalis denſiſſima, dum floret, hemisphærica, fructeſcens plana. Semina cinerea, ovata, hinc plana, illinc tribus ſtriis parum elevatis notata, villoſulaque. Iacqu. l. c.

Cervaria. 2. A. foliis pinnatis decuſſatis inciſo-angulatis, ſeminibus nudis. *Sp. pl. p.* 352. *Iacqu. auſtr. t.* 69.

Selinum foliis duplicato-pinnatis, pinnis ſemibilobatis, circumſerratis. *Hall. hiſt. n.* 804.

Selinum Cervaria. *Scop. carn. tom.* 1. *p.* 201. *Crantz faſc. p.* 167. *t.* 3. *f.* 1.

Seſeli montanum ſecundum pannonicum. *Cluſ. hiſt. p.* CXCIII.

In montoſis fruticoſis, collibusque aſperis. Fl. Auguſto, Septembri. Per.

Folia radicalia ſunt terpinnata. Caulina bipinnata: foliolis oppoſitis, ſeſſilibus aut breviſſime petiolatis, ſubovatis, acutis, inæqualiter ſerratis cum dentibus mucronatis, aliis integris, aliis lobato-inciſis, extremis plerumque trilobis. Involucrum univerſale componitur ex foliolis circiter decem, lanceolato-linearibus in ſetulam deſinentibus, reflexis, margine membranaceis. Partiale fit ex foliolis ſetaceis. Semina ſubrotunda, extus compreſſo-plana, in marginem attenuata membranaceum. Iacqu. l. c.

Oreoſelinum. 3. A. foliolis divaricatis. *Sp. pl. p.* 352. *Iacqu. auſtr. t.* 68.

Selinum foliis triplicato-pinnatis, refractis. *Hall. nom. n.* 803.

Selinum Oreoſelinum. *Scop. carn. tom.* 1. *p.* 201. *Crantz faſc. p.* 169.

Oreoſelinum. *Cluſ. hiſt. p.* CXCV.

In collibus, montaniſque herbidis. Fl. Auguſto, Septembri. Per.

Folia multiplicato-pinnata coſtam habent teretem, ex baſi membranacea dilatata atque amplexante exortam, in foliis radicalibus ſæpe longam, quæ in ramos ſubdividitur, ad angulos rectos iisve majores egreſſos, ad nodos, ac ſi fracti forent, plerumque diſtortos. Foliola ſunt pinnatifida, acuta, ſemitriloba, aut ſolo apice trifida. Involucri foliolorum numerus varius. Fructus complanatus conſtat duobus ſeminibus orbicularibus, interdum baſi magis elongatis, planis, nucleo tribus ſtriis variegato, margine membranaceo. Iacqu. l. c.

cretenſis. 4. A. foliis linearibus planis hirſutis, petalis bipartitis, ſeminibus oblongis hirſutis. *Sp. pl. p.* 352. *Iacqu. auſtr. t.* 62.

Libanotis foliis hirſutis, triplicato-pinnatis, lobulis profundiſſime bipartitis. *Hall. hiſt. n.* 745.

Libanotis cretenſis. *Scop. carn. tom.* 1. *p.* 192.

Libanotis Daucus creticus off. *Cranz faſc. p.* 219.

Daucus creticus. *Cluſ. hiſt. p.* CXCVIII.

In rupeſtribus alpinis. Fl. Iulio, Auguſto. Per.

Folia multiplicato-pinnata: pinnulis linearibus, brevibus, acutis, planis. Involucrum univerſale aut nullum, aut conſtans foliolis ab uno ad ſex, lanceolatis. Partiale ſit ex foliolis a quatuor ad ſeptem, lanceolatis, concavis. Semina ovalia, hirſuta, hinc concava, illinc convexa, & quadriſulcata cum angulis, ſinubusque obtuſis. Iacqu. l. c.

Matthioli. 5. A. foliolis capillaribus, ſeminibus oblongis hirſutis.

Athamanta foliis ſupradecompoſitis diffuſis: foliolis elongato-ſetaceis, ſeminibus acutis hirſutis. *Wulfen in Iacqu. collect. tom.* 1. *p.* 211. *ic. t.* 57.

Libanotis rupeſtris. *Scop. carn. tom.* 1. *p.* 192.

Meon quod Athamanticum vocant. *Matthiol. comm. p.* 20. *f.* 29.

In rupeſtribus Croatiæ, Carnioliæ calidioris, Litoralis. Fl. Iunio, Iulio. Per.

Differt a præcedente caule magis ramoſo, pinnulis anguſtioribus, magisque elongatis.

PEUCEDANUM. Fructus ovatus, utrinque ſtriatus, ala cinctus.

officinale. 1. P. foliis quinquies tripartitis linearibus integerrimis. *Hort. cliff.* 93.

Peucedanum. *Bauh. hiſt.* 3. *p.* 36.

Le Fenouil. Regn. bot.

In pratis ſiccioribus, ad ſylvarum margines in Pannonia inter *Bahrendorf* & *Bruck an der Leitha.* Fl. Iunio, Iulio. Per.

Radix valde craſſa, externe nigricans, interne flaveſcens, vulnerata flavum balſameum ſuccum fundens, qui mox in reſinam ſulphurei coloris

concreſcit. Foliola ſunt linearia, vel lineari-lanceolata. Filiformia in noſtra planta obſervare nondum licuit.

Silaus. 2. P. foliolis pinnatifidis: laciniis oppoſitis, involucro univerſali diphyllo. *Sp. pl. p.* 354. *Iacqu. auſt. t.* 15.

Peucedanum foliis triplicato-pinnatis, pinnulis nervo diſtinctis, lanceolatis, imparibus trilobatis. *Hall. hiſt. n.* 797.

Seſeli pratenſe. *Crantz faſc. p.* 209. *t.* 6.

In Auſtria, Styria, Pannonia in pratis præcipue udis. Fl. Iunio, Iulio. Per.

Pinnulæ ſunt mucronatæ. Involucrum univerſale aut nullum, aut conſtans foliolo lineari unico, ſimilibusve paucis. Partiale componitur ex pluribus, lanceolato-linearibus. Semina ovata, aut oblonga, hinc plana, illinc convexa, ac quadriſulcata, ſive tribus ſtriis elevatis notata. Iacqu. l. c.

alſaticum. 3. P. foliis pinnatifidis: laciniis trifidis obtuſiuſculis. *Sp. pl. p.* 354. *Iacqu. auſtr. t.* 70.

Selinum foliis quadruplicato-pinnatis, nervis canaliculatis. *Hall. hiſt. n.* 798.

Selinum alſaticum. *Crantz faſc. p.* 159.

Umbellifera alſatica magna, umbella parva ſublutea. *Bauh. hiſt.* 3. *p.* 106.

In apricis aſperis, ad ſepes Auſtriæ, Styriæ, Pannoniæ. Fl. Auguſto, Septembri. Per.

Caulis ramoſiſſimus. Folia ſunt multiplicato-pinnata. Foliola inciſa; laciniis lanceolato-oblongis, acutis. Involucrum tam univerſale quam partiale conſtat foliolis circiter quinis. Semina compreſſa, nucleo nigricante extus ſtriis tribus in membranulam elevatis notato, & excurrente in marginem undique membranaceum. Iacqu. l. c.

CRITHMUM. Fructus ovalis, compreſſus. Floſculi æquales.

mariti-mum. 1. C. foliolis lanceolatis, carnoſis. *Sp. pl. p.* 354. *Iacqu. hort. tom.* 2. *p.* 88. *t.* 187. *Scop. carn. tom.* 1. *p.* 219.

Crithmum multis ſive Fœniculum marinum. *Bauh. hiſt.* 3. *p.* 194. *Zann. iſt. p.* 74. *t.* 9.

In Litorali in rupeſtribus ad mare. Fl. a Iunio ad Septembrim. Per.

FERULA. Fructus ovalis, compreſſo-planus, ſtriis utrinque tribus.

nodiflo-ra. 1. F. foliolis appendiculatis, umbellis ſubſeſſilibus. *Sp. pl. p.* 356. *Iacqu. auſtr. app. t.* 5. *Scop. carn. tom.* 1. *p.* 204.

In pratis, aliisque herbidis locis Litoralis. Fl. Iulio. Auguſto. Per.

Folia caulina infima alterna, ſubſequa oppoſita, ſuperiora verticillata. Omnia ſupradecompoſita: pinnulis extremis ſetaceo-linearibus, acutis, integerrimis & planis. Involucri univerſalis foliola reflexa; particularis patentiſſima. Utriusque plura, lanceolata & acuta. Iacqu. l. c.

LASERPITIUM. Fructus oblongus: angulis octo membranaceis. Petala inflexa, emarginata, patentia.

latifo-lium. 1. L. foliolis cordatis inciſo-ſerratis. *Sp. pl. p.* 356. *Iacqu. auſtr. t.* 146. *Scop. carn. tom.* 1. *p.* 195. *Hall. hiſt. n.* 792.

Laſerpitium glabrum. *Crantz faſc. p.* 181.

Laſerpitium aſperum. *ejusd. faſc. p.* 179. *t.* 1. *f.* 2.

Seſeli æthiopicum Matthioli. *Cluſ. hiſt p.* CXCIV.

In montanis, & ſubalpinis. Fl. Iulio. Per.

Folia ſunt duplicato-pinnata: foliolis cordatis, integris, vel rarius lobatis, dentato-ſerratis, obtuſis, vel acutis, inferioribus petiolatis, ſummis ſæpe ſeſſilibus, vel utrinque glaberrimis, vel ſolo dorſo aſperis. Semina glabra, nucleo oblongo, hinc plana, illinc quadrialata. Iacqu. l. c.

aquilegifolium. 2. L. foliolis obtuſis baſi ovatis lobatis. *Syſt. veget. p.* 281. *Iacqu. auſtr. t.* 147.

Siler trilobum. *Crantz faſc. p.* 186.

Seſeli æthyopici aliud genus. *Clus. hiſt. p.* CXCV.

In Auſtria in collibus ſaxoſis, in montoſis inter frutices. Fl. Iunio. Per.

Folia ſupradecompoſita: foliolis lobatis & inciſis, minime cordatis. Fructus ex ovato-oblongus, compreſſus, obtuſus, in ſemina bina diſpeſcitur, hinc plana, inde convexa, aliſque membranaceis brevibus quinis inſtructa, cum interjectis alternantibus quatuor alis anguſtioribus. Iacqu. l. c.

Arcangelica. 3. L. foliolis ovatis lanceolatisque ſerratis hirſutis integris labatisque. *Wulfen in Iacqu. collect. tom.* I. *p.* 214. *ic. t.* 58.

In Carniolia in rupeſtribus, umbroſis montis *Slivinza* prope *Zirchnizerſee*, & in monte *Fridrichſtein*. *Wulfen.* Ego in Croatia in ſaxoſis, & ad fruticeta inter *Fucine* ac *Merslavodizza*. Fl. Iulio. Per.

Caulis hirſutus, ſtriatus. Folia ſupradecompoſita. Foliola ſunt ovata, aut lanceolata, ſerrata, integris lobata intermixta. Omnia ſupra glabra, ſubtus hirſuta. Involucrum univerſale conſtat foliolis pluribus, lineari lanceolatis, integris, inciſisque. Partiale item polyphyllum. Fructus oblongus alis membranaceis quatuor inſtructus. Wulfen, l. c.

ſilaifolium. 4. L. caule glabro, radice multicipite, foliolis pinnatifidis, laciniis lanceolatis. *Syſt. veget. p.* 281. *Iacqu. auſt. app. t.* 44.

In Carnioliæ calidioris, Litoralis ſaxoſis, aſperis. Fl. Iulio, Auguſto. Per.

Folia multiplicato - pinnata, glabra: pinnis profunde inciſis planis; laciniis oblongis, breviter mucronatis. Pedunculi oppoſitifolii, erecti, ſtriati. Involucri univerſalis foliola pauca, linearia, integra, quandoque pinnatifida, umbella duplo breviora. Particularis plura, umbellulæ longitudine. Semina elliptica, glabra, hinc plana, illinc convexa, atque alis quinque membranaceis munita. Iacqu. l. c.

prutenicum. 5. L. foliolis lanceolatis integerrimis: extimis coalitis. *Sp. pl. p.* 357. *Iacqu. auſtr. t.* 153.

Laſerpitium gallicum. *Scop. carn. tom.* 1. *p.* 196.

Laſerpitium ſelinoides. *Crantz faſc. p.* 182.

In Auſtria in fruticoſis, ſylvaticis inter *Bertholdsdorf* & *Giſshübel*, in monte *Gallitzinberg*. In Pannonia in ſylvis montanis prope Poſonium. In Forojulio circa Goritiam. Fl. Auguſto.

Folia radicalia ac ſubſequa inferiora ſunt ſubbipinnata, ſuperiora ſucceſſive minus diviſa. Pinnæ ex lanceolato oblongæ, confluentes, prona parte glabræ, ad nervos dorſales piloſæ, breviter mucronatæ, in adultis foliis ſæpe ex fuſco maculatæ. Semina excurrunt in alam utrinque membranaceam, in parte convexa adhuc duabus alis inſtructa ſunt. Iacqu. l. c.

peucedanoides. 6. L. foliolis lineari - lanceolatis venoſo - ſtriatis diſtinctis. *Sp. pl. p.* 358. *Wulfen in Iacqu. collect. tom.* 1. *p.* 220. *ic. t.* 349.

Laſerpitium peucedanoides foliorum ſegmentis anguſtiſſimis. *Segu. veron.* 3. *p.* 227. *t.* 7.

Laſerpitium anguſtifolium. *Scop. carn. tom.* 1. *p.* 197.

In Carnioliæ, Croatiæ ſubalpinis & alpinis. Fl. Iulio, Auguſto. Per.

Folia ſunt multiplicato - pinnata. Pinnulæ lineari - lanceolatæ. Involucrum univerſale & partiale conſtat foliolis pluribus. Fructus oblongus,

alatus. Non differt a Laſerpitio anguſtifolio Linn., *nam foliola ſunt vel anguſtiora vel latiora, prout diverſa plantæ ætas, ac diverſa ſoli natura, in quo creſcit.* Wulfen.

Siler. 7. L. foliolis ovali-lanceolatis integerrimis petiolatis. *Sp. pl. p.* 357. *Iacqu. auſtr. t.* 145. *Scop. carn. tom.* 1. *p.* 196.

Siler montanum. *Clus. hiſt. p.* CXCV. *Crantz faſc. p.* 185.

Laſerpitium foliis duplicato-pinnatis, pinnulis integerrimis lanceolatis, ſimplicibus & ternatis. *Hall. hiſt. n.* 794.

In Auſtriæ, Carnioliæ montanis & ſubalpinis. Fl. Iulio, Auguſto. Per.

Pinnulæ ſunt anguſte lanceolatæ, plurimæ ternatæ. Fructus oblongus, ſubteres. Semina hinc plana & ſtriata, inde convexa & inſtructa alis quatuor membranaceis, quas inter lineæ elevatæ decurrunt. Iacqu. l. c.

ſimplex. 8. L. ſcapo nudo aphyllo ſimplici, foliis pinnatis multifidis acutis linearibus, umbella ſemigloboſa. *Syſt. veget. p.* 282. *Wulfen in Iacqu. miscell. tom.* 2. *p.* 40. *t.* 2.

Laſerpitium foliis triplicato-pinnatis, pinnulis lanceolatis; involucris ſemitrifidis. *Hall. hiſt. n.* 796.

In Carinthia in ſuperioribus apricis jugis alpis melbingenſis in diſtrictu gmündenſi; tum in Taurero naſsfeldenſi. *Wulfen.* Per.

HERACLEUM. Fructus ellipticus, emarginatus, compreſſus, ſtriatus, marginatus. Corollæ difformis petala inflexo-emarginata. Involucrum caducum.

Sphondylium. 1. H. foliolis pinnatifidis, floribus difformibus, *Retzii ſcand. p.* 66. *Sp. pl. p.* 358.

Sphondylium *Branca*, foliis pinnatis, pinnis pinnatifidis, floribus radiatis. *Mœnch. meth. p.* 83. *Scop. carn. tom.* 1. *p.* 204.

La Berce: Regn. bot.

In nemoroſis, pratis humidis, & alibi. Fl. Iulio. Per.

anguſti-folium. 2. H. foliis cruciato-pinnatis: foliolis linearibus, corollis floſculoſis. *Syſt. veget. p.* 282. *Iacqu. auſtr. t.* 173.

Heracleum longifolium. *Iacqu. auſtr. t.* 174.

In Auſtriæ, Styriæ alpinis. Fl. Auguſto. Per.

auſtria-cum. 3. H. foliis pinnatis utrinque rugoſis ſcabris, floribus ſubradiatis. *Syſt. veget. p.* 282. *Iacqu. auſtr. t.* 61. *en. p.* 216. *Crantz faſc. p.* 153. *t.* 1.

Sphondylium auſtriacum. *Scop. carn. tom.* 1. *p.* 204.

In Auſtria, Styria, Carniolia in apricis ſaxoſis herbidis alpinis. Fl. a Iunio ad Septembrim. Per.

LIGUSTICUM. Involucrum univerſale, & involucella membranacea. Flores æquales, fertiles. Fructus oblongus, utrinque quinqueſulcatus.

pelopon-nenſe. 1. L. foliis multplicato-pinnatis: foliolis pinnatim inciſis. *Iacqu. auſtr. app. t.* 13. *Scop. carn. tom.* 1. *p.* 205.

Liguſticum foliis duplicato-pinnatis, pinnulis acute dentatis, longe lanceolatis. *Hall. hiſt. n.* 758.

In montibus Forijulii. Fl. Majo, Iunio. Per.

Folia radicalia, & caulina inferiora ſunt tripinnata. Superiora bipinnata. Pinnæ ſunt pinnatifidæ, longe lanceolatæ, confluentes & acutæ. Involucri foliola quina. Fructus compreſſus &

glaber diſpeſcitur in ſemina oblonga, intus plana, breviterque bimarginata, extus tribus alis donata. Iacqu. l. c.

auſtriacum. 2. L. foliis bipinnatis: foliolis confluentibus inciſis integerrimis. *Syſt. veget. p.* 283. *Iacqu. auſtr. t.* 151. *Crantz faſc. p.* 197.

Seſeli primum montanum pannonicum. *Cluſ. hiſt. p.* CXCIII.

In Alpium auſtriacarum declivibus. Fl. Iulio, Auguſto. Per.

Folia ſunt multiplicato-pinnata: foliolis ultimis lanceolatis, aut ovatis, inciſis & ſubſerratis, confluentibus. Petala ſubovata, parum concava, patentiſſima. Fructus ovatus & acutus conſtat ſeminibus duobus, hinc planis, inde convexis & profunde quadriſulcatis. Iacqu. l. c.

ANGELICA. Umbellulæ globoſæ. Corollæ æquales. Fructus ſubrotundus, angulatus, ſtylis reflexis.

Archangelica. 1. A. foliorum impari lobato. *Sp. pl. p.* 360. *Fl. dan. t.* 206. *Scop. carn. tom.* 1. *p.* 199.

Angelica *officinalis*, foliis pinnatis, pinnis ovato-lanceolatis, impari triloba, umbellis globoſis. *Mœnch. meth. p.* 81.

Angelica foliis duplicato pinnatis, ovato-lanceolatis ſerratis. *Hall. hiſt. n.* 807.

L' Angelique. Regn. bot.

In montibus editioribus Litoralis, Forijulii, Carnioliæ, Carinthiæ. Fl. Iulio, Auguſto.

ſylveſtris. 2. A. foliolis æqualibus ovato-lanceolatis ſerratis. *Sp. pl. p.* 361. *Scop. carn. tom.* 1. *p.* 199.

Angelica foliis duplicato-pinnatis, ovatis ſerratis. *Hall. hiſt. n.* 806.

Selinum ſylveſtre. *Crantz faſc. p.* 177.

L' Angelique ſauvage. Regn. bot.

In pratis udis, ſylvis demiſſis umbroſis, ad fluvios, & in alpium declivibus humidis. Fl. Iulio, Auguſto.

verticillaris. 3. A. foliis divaricatiſſimis: foliolis ovatis ſerratis, caule pedunculis verticillato. *Syſt. veget. p.* 284. *Iacqu. hort. tom.* 2. *p.* 60. *t.* 130.

In Auſtria *in Reichenau an der Schwarza.* Fl. Iulio.

SIUM. Involucrum & involucella polyphylla. Petala cordata. Fructus ſubovatus, ſtriatus.

latifolium. 1. S. foliis pinnatis, umbellis terminalibus. *Sp. pl. p.* 361. *Iacqu. auſtr. t.* 66.

Sium foliis pinnatis, argute dentatis, umbellis erectis. *Hall. hiſt. n.* 777.

Coriandrum latifolium. *Crantz faſc. p.* 212.

In aquis ſtagnantibus, inundatis, inter arundines Auſtriæ, Pannoniæ. Fl. Iulio, Auguſto.

Foliola ſunt lanceolata, argute ſerrata, & acuta, tria vel quatuor cum terminante impari, interdum ſine. Sub aqua folia quandoque egrediuntur difformia in capillares ferme pinnas diviſa. Atque etiam folia radicalia extra aquam elevata variare ſolent. Iacqu. l. c.

anguſtifolium. 2. S. foliis pinnatis: foliolis inæqualiter argute que ſerratis, auriculatis, umbellis oppoſitifoliis pedunculatis. *Retzii obs. faſc.* 1. *p.* 14. *prodr. ſcand. p.* 67. *Syſt. veget. p.* 284. *Iacqu. auſtr. t.* 67.

Sium foliis radicalibus ovatis, pinnulis dentatis, caulinis appendiculatis, umbellis alaribus. *Hall. hiſt. n.* 778.

Seſeli nodiflorum. *Scop. carn. tom.* 1. *p.* 213.

Apium Sium. *Crantz faſc. p.* 215.

In rivulis, & aquis ſtagnantibus. Fl. Iulio, Auguſto. Per.

Foliola ſunt ex lanceolato oblonga, argute ſerrata; alia ſunt ex utraque parte ad baſin auriculato-inciſa, alia magis ſimplicia. Impar foliolum vel eſt ſimplex, vel appendiculatum, vel ſemitrilobum. Umbellæ omnes pedunculatæ, primo terminales, dein evadunt pleræque oppoſitifoliæ, quia novus ramus folium inter & pedunculum egreditur. Involucri univerſalis foliola ſunt circiter ſena, integra, vel inciſa. Iacqu. l. c.

repens. 3. S. caule repente, foliolis ſubrotundis dentato-angulatis. *Syſt. veget. p.* 284. *Iacqu. auſtr. t.* 260.

In pratis humidis agri vindobonenſis ad Danubium, & Viennam. Fl. a Iulio, ad Octobrim. Per.

Folia ſunt pinnata. Foliola oppoſita, ſeſſilia, ex ovato ſubrotunda, inæqualiter ſerrata; terminali utplurimum magis, minusve profunde trilobo. Indolucrum conſtat foliolis ſex circiter, ſic & involucellum.

Falcaria. 4. S. foliolis linearibus decurrentibus connatis. *Sp. pl. p.* 362. *Iacqu. auſtr. t.* 257.

Sium foliis firmis, ſerratis, pinnatis, nervo folioſo latescente. *Hall. hiſt. n.* 782.

Seſeli Falcaria. *Crantz fasc. p.* 208. *Scop. carn. tom.* 1. *p.* 213.

Iu pratis, cultis, ad vias, alibique in Litorali, Forojulio, Tyroli calidiore, Auſtria, & Pannonia. Fl. ab Auguſto ad Octobrim. Per.

Radix repens. Folia radicalia ſunt ternata, aut ſimplicia, aut difformiter pinnatifida. Coſta compreſſa ex vagina oritur membranacea & canaliculata. Pinnæ ſunt ex lineari enſiformes, acutæ, planæ, argute ſerratæ, per coſtas particulares magis minusve decurrentes. Involucri univerſalis foliola octona, ſubſetacea, longitudine inæqualia, unico plerumque reliquis duplo longiore. Iacqu. l. c.

OENANTHE. Floſculi difformes: in disco ſeſſiles, ſteriles. Fructus calyce & piſtillo coronatus.

fiſtuloſa. 1. Oe. ſtolonifera, foliis caulinis pinnatis filiformibus fiſtuloſis. *Sp. pl. p.* 365. *Fl. dan. t.* 846.

Oenanthe foliis radicalibus rotunde lobatis, planis, caulinis fiſtuloſis. *Hall. hiſt. n.* 755.

In Auſtria in pratis udis, & ad foſſas prope *Schwadorf*, *Fiſchamend*. Fl. Iunio, Iulio. Per.

Folia radicalia, & caulina ima ſunt bipinnata: pinnulis profunde inciſis. Superiora ſunt pinnata. Flores radii ſunt difformes, disci regulares.

pimpinelloides. 2. Oe. foliolis radicalibus cuneatis fiſſis; caulinis integris linearibus longiſſimis ſimplicibus. *Syſt. veget. p.* 286. *Iacqu. auſtr. t.* 394. *Scop. carn. tom.* 1. *p.* 217. *Crantz faſc. p.* 201.

In udis Auſtriæ, Pannoniæ. In pratis agri fluminenſis, goritienſisque. Fl. Majo, Iunio. Per.

Radix tuber quoddam exhibens, ex quo undique fibræ nigræ exoriuntur, abeuntes pleræque in tuberculum ovatum aut ſubrotundum. Folia radicalia & caulina infima ſunt bipinnata, vel tripinnata; aut etiam dumtaxat pinnata in individuis minoribus. Caulina ſuperiora pinnata; ſumma ſimplicia. Pinnulæ in foliis inferioribus ſunt ſubovatæ, pinnatifidæ, vel laciniatæ, aut furcatæ. In ſuperioribus autem foliis ſunt integræ, & lineari-lanceolatæ. Petioli fiſtuloſi. Iacqu. l. c.

PHELLANDRIUM. Floſculi disci minores. Fructus ovatus, lævis, coronatus perianthio & piſtillo.

aquaticum. 1. P. foliorum ramificationibus divaricatis. *Sp. pl. p.* 366. *Scop. carn. tom.* 1. *p.* 216.

Phellandrium foliis refractis. *Hall. hiſt. n.* 757.

Cicutaria paluſtris. *Lob. ic.* 735.

La Ciguë aquatique. Regn. bot.

In aquis ſtagnantibus. Fl. æſtate.

Mutellina. 2. P. caule ſubnudo, foliis bipinnatis. *Sp. pl. p.* 366. *Iacqu. auſtr. t.* 56. *Scop. carn. tom.* 1. *p.* 216.

Seſeli foliis duplicato-pinnatis, pinnulis ſemipinnatis, lobulis lanceolatis. *Hall. hiſt. n.* 763.

Liguſticum Mutellina. *Crantz faſc. p.* 198.

Daucus montanus. *Cluſ. hiſt. p.* CXCVIII.

In alpium herbidis. Fl. æſtate. Per.

CICUTA. Fructus ſubovatus, ſulcatus.

viroſa. 1. C. umbellis oppoſitifoliis, petiolis marginatis obtuſis. *Sp. pl. p.* 366. *Fl. dan. t.* 208.

Sium foliis duplicato-pinnatis, pinnulis acute ſerratis, trifidis & ſimplicibus. *Hall. hiſt. n.* 781.

Cicuta aquatica. *Geſn. hort. p.* 253.

In agro tergeſtino in aquoſis inter *Saule & Dolline.* Fl. æſtate. Per.

Folia glabra, multiplicato-pinnata: foliolis ſerratis, decurrentibus. Involucrum univerſale aut nullum, aut conſtans foliolo unico, aut tribus. Involucellum vero habet foliola plura. Fructus eſt ſubovatus, ſulcatus, conſtans ſeminibus duobus hinc convexis, inde planis.

AETHUSA. Involucella dimidiata, triphylla, pendula. Fructus ſtriatus.

Cynapium. 1. Ae. foliis conformibus. *Syſt. veget. p.* 286. *Curtis lond. Scop. carn. tom.* 1. *p.* 206. *Hall. hiſt. n.* 765.

Cicuta minor, petroſelino ſimilis. *Bauh. pin.* 160.

La petite Ciguë. Regn. bot.

In cultis. Fl. æſtate.

Involucrum univerſale nullum. Partiale dimidiatum, extus poſitum. Fructus ovato-ſubrotundus dividitur in ſemina duo, hinc plana, inde convexa, & ſtriata.

Meum. 2. Ae. foliis omnibus multipartito-ſetaceis. *Syſt. veget. p.* 287.

Meum athamanticum. *Iacqu. auſtr. t.* 303.

Athamanta Meum. *Sp. pl. p.* 353.

Seſeli Meum. *Scop. carn. tom.* 1. *p.* 212.

Liguſticum Meum. *Crantz faſc. p.* 198.

Meum. *Dod. purg. p.* 478. *t.* 62. *Hall. hiſt. n.* 761.

Daucus Meum. *Cluſ. hiſt. p.* CXCVIII.

In alpibus. Fl. Iulio, Auguſto. Per.

Involucrum univerſale vel nullum, vel monophyllum. Particulare dimidiatum ad extremam umbellulæ dimidiam partem locatum. Flores plerique ob germinis defectum ſteriles. Semina oblonga, acuta, glabra, ſtriata. Iacqu. l. c.

SCANDIX. Corolla radiata. Fructus ſubulatus. Petala emarginata. Floſculi disci ſæpe maſculi.

odorata. 1. S. ſeminibus ſulcatis angulatis. *Sp. pl. p.* 368. *Iacqu. auſtr. app. t.* 37.

Myrrhis foliis triplicato-pinnatis, mollibus, ſeminibus ſulcatis, maximis. *Hall. hiſt. n.* 753.

Myrrhis odorata. *Scop. carn. tom.* 1. *p.* 207.

Le Cerfeuil muſqué. Regn. bot.

In paſcuis Carnioliæ, Croatiæ ſubalpinis. Fl. Iunio, Iulio. Per.

Folia multiplicato-pinnata; mollia, ſubtus villoſula: pinnulis ultimis ovatis, vel oblongis, inciſo-ſerratis, inque foliolum longe lanceolatum acutumque ſubconfluentibus. Involucrum univerſale nullum. Partiale pentaphyllum. Floſculi plerique ſunt mares; pauci hermaphroditi & fertiles in cen-

tro. Petala inflexo-cordata, patentiſſima, alba, extima umbellularum majora. Semina linearia, longa, fuſca, nitida, acuminata, hinc plana, illinc acute pentagona. Iacqu. l. c.

Pecten. 2. S. ſeminibus roſtro longiſſimo. *Syſt. veget. p.* 287. *Iacqu. auſtr. t.* 263. *Curtis lond. Fl. dan. t.* 844. *Scop. carn. tom.* 1. *p.* 211.

Myrrhis ſeminis cornu longiſſimo. *Hall. hiſt. n.* 754.

Chærophyllum Pecten Veneris. *Crantz faſc. p.* 189.

Inter ſegetes. Fl. æſtate.

Involucrum nullum. Involucella conſtant foliolis utplurimum quinque, bis vel ter dentatis. Petala in floſculis centralibus æqualia, in exterioribus exteriora longiora. Germen oblongum, ſubhirſutum. Semina ſcabra, hinc ſulcata, illinc convexa & ſtriata, in roſtrum excurrunt longiſſimum. Iacqu. l. c.

Cerefolium. 3. S. ſeminibus nitidis ovato-ſubulatis, umbellis ſeſſilibus lateralibus. *Sp. pl. p.* 368. *Iacqu. auſtr. t.* 390.

Cerefolium foliis glabris, triplicato-pinnatis, lobulis obtuſis. *Hall. hiſt. n.* 747.

Chærophyllum Cerefolium. *Crantz faſc. p.* 191.

Le Cerfeuil. Regn. bot.

In Auſtria ad ſylvarum margines, ſepes. Fl. Majo, Iunio.

Folia tenuia, nitida, triplicato-pinnata: foliolis ſubovatis, ſerrato-inciſis. Umbellæ componuntur e radiis paucis. Umbellulæ ſunt inæquales. Involucrum univerſale plerumque nullum. Particulare conſtat foliolo unico, ſetaceo. Flores pauci carent germine. Semina ſunt ſubulata, fuſca, ſemiteretia. Iacqu. l. c.

Anthriſcus. 4. *S.* ſeminibus ovatis hiſpidis, corollis uniformibus, caule lævi. *Sp. pl. p.* 368. *Iacqu. auſtr. t.* 154. *Fl. dan. t.* 863. *Curtis lond.*

Caucalis vaginis lanuginoſis, foliis triplicato-pinnatis, ſeminibus roſtratis. *Hall. hiſt. n.* 743.

Caucalis Scandix. *Scop. carn. tom.* 1. *p.* 191.

In agro vindobonenſi ad ſepes, in fruticoſis, aggeribus. Fl. Majo, Iunio.

Folia ſunt multiplicato-pinnata, ſubhirſuta, aut etiam glabra. Semina oblonga, teretia, interne ſulco longitudinali exarata, undique pilis brevibus, hamatis, rigidis hirta, attenuata in apicem gracilem, glabrum. Iacqu. l. c.

nodoſa. 5. S. ſeminibus ſubcylindricis hiſpidis, caule hiſpido, geniculis tumidis. *Sp. pl. p.* 369. *Iacqu. hort. tom.* 3. *p.* 16. *t.* 25.

Cerefolium annuum nodoſum, ſemine aſpero majore. *Moris. hiſt.* 3. *p.* 303. *ſect.* 9. *t.* 10. *f.* 4.

In agro fluminenſi ad ſepes, in fruticoſis, copioſiſſime Terſacti in ſylva P. P. Franciſcanorum. Fl. Majo.

Caulis fiſtuloſus, pilis reflexis hiſpidus, infra ramorum exortum tumidus. Folia ſubtriplicato-pinnata: foliolis inciſo-ſerratis, lobatisque, obtuſis. Umbella conſtat radiis plerumque ternis. Involucrum univerſale vel nullum, vel unifolium. Partiale fit e foliolis circiter quinis. Flores omnes fertiles. Fructus ſeparatur in ſemina hinc ſulcata, inde teretia & hiſpida. Iacqu. l. c.

infeſta. 6. S. ſemine exteriore hiſpido, umbellulis confertiſſimis hemiſphæricis. *Syſt. veget. p.* 287. *Iacqu. auſtr. t.* 46.

In Auſtriæ, Pannoniæ agris. Fl. Iulio, Auguſto.

Folia ſunt pinnata. Foliola in plantis majoribus modo ſimiliter pinnata, modo pinnatifide inciſa; utrinque ob pilos minutos ſubhirſuta, & ſcabriuſcula. Involucrum univerſale vel nullum, vel unifolium. Partiale conſtat foliolis quatuor, vel quinque, parvis. Semina ſunt oblonga, obtuſa, fuſca, ac ſetis uncinatis munita, quarum extimæ in ſeminibus fructuum exteriorum ſunt longæ, reliquæ breviſſimæ. Iacqu. l. c.

CHÆROPHYLLUM. Involucrum reflexum, concavum. Petala inflexo-cordata. Fructus oblongus, lævis.

ſylveſtre. 1. C. caule lævi ſtriato, geniculis tumidiuſculis. *Syſt. veget. p.* 288. *Iacqu. auſtr. t.* 149. *Curtis lond. Scop. carn. tom.* I. *p.* 218. *Crantz faſc. p.* 192.

Cerefolium foliis triplicato-pinnatis, acute dentatis, glabris, nervis hirſutis. *Hall. hiſt. n.* 748.

In ſylvis, ad ſepes, pagos. Fl. a Majo ad Auguſtum. Per.

Folia ſunt multiplicato-pinnata: pinnulis lanceolatis, acutis, ſerrato-inciſis, extremis longe confluentibus, utrinque ſæpe nitidis, facie glabris & ex atro virentibus, dorſo pallidioribus nervisque præprimis & ad oras pubeſcentibus, quandoque totis hirſutulis. Petala plana ac obverſe ovata: centralium floſculorum ſubæqualia; exteriorum vero extimum majus & ſubcordato-emarginatum. Fructus oblongus, acutus, nitidus, niger, lævis, ſeparabilis in ſemina oblonga, antice ſulcata. Iacqu. l. c.

bulboſum. 2. C. caule lævi geniculis tumido, baſi hirto. *Syſt. veget. p.* 288. *Iacqu. auſtr. t.* 63. *Crantz faſc. p.* 189.

Myrrhis radice bulboſa, caule inferne hirto, ſuperne lævi, foliis triplicato-pinnatis, acute inciſis. *Hall. hiſt. n.* 752.

Cicutaria pannonica. *Clus. hiſt. p.* CC.

In Pannonia, Auſtria in fruticoſis, ad ſepes. Fl. Majo, Iunio.

Radix napiformis. Caulis teres, virens, punctis, maculisque ferrugineis adſperſus, inferioribus duobus, tribusve articulis hirtis, reliquis ſuperioribus glabris, ac infra folia magis minusve tumentibus. Folia ſunt multiplicato-pinnata: pinnulis pinnatifide inciſis, acutis, ſupra glabris, una cum petiolis dorſo pilis albis, vagis, longisque adſperſis. In quibusdam indivi-

duis pinnarum laciniæ ferme capillaceæ. Involucrum univerſale vel nullum, vel monophyllum. Particulare conſtat foliolis paucis, acutis. Semen aciculare, hinc planum cum ſulco medio, illinc convexum, & quinque lineis elevatis notatum. Iacqu. l. c.

temulum. 3. C. caule ſcabro: geniculis tumidis. *Sp. pl. p.* 370. *Iacqu. auſtr. t.* 65. *Curtis lond. Fl. dan. t.* 918. *Crantz faſc. p.* 190.

Myrrhis foliis hirſutis, laciniis obtuſis, caule geniculato. *Hall. hiſt. n.* 750.

In ſylvis, ad ſepes. Fl. Majo.

Caulis punctis atropurpureis ſæpe adſperſus, ſub geniculis tumens, breviſſimis pilis veſtitus, ad tactum modice ſcaber. Folia duplicato-pinnata, hirſutula. Pinnæ ſubovatæ, obtuſæ, lobato-inciſæ. Involucrum univerſale raro adeſt. Partiale fit e foliolis circiter ſenis, lanceolatis, acutis, reflexis. Petala bifida. Flos centralis, atque extimi fructum gerunt. Reliqui intermedii plures germine carent. Semina oblonga, glabra, fuſca, bilinearia, acuta, intus concava, extus obſolete, obtuſeque quinqueſulcata. Iacqu. l. c.

hirſutum. 4. C. caule æquali, foliolis inciſis acutis, fructibus biariſtatis. *Syſt. veget. p.* 288. *Iacqu. auſtr. t.* 148. *Crantz faſc. p.* 194.

Myrrhis radice prælonga, foliis triplicato-pinnatis, acutis, ſeminibus cylindricis. *Hall. hiſt. n.* 751.

Scandix hirſuta. *Scop. carn. tom.* 1. *p.* 211.

In ſubalpinis, & alpinis. Fl. Iulio, Auguſto. Per.

Folia ſunt fere biternata: foliolis oblongis, pinnatifide inciſis, argute ſerratis, acutis, facie glabris, dorſo magis dilute virentibus, lucidis & ad venas piloſis. Floſculi circumferentiæ, & unicus centralis ſoli germen poſſident oblongum, compreſſum. Qui vero in diſco floſculi reliqui ſunt carent germine & ſtylis, Iacqu. l. c.

aromaticum. 5. C. caule æquali, foliolis ſerratis integris, fructibus biariſtatis. *Syſt. veget. p.* 288. *Iacqu. auſtr. t.* 150.

Scandix tinctoria. *Scop. carn. tom.* 1. *p.* 212.

In ſylvaticis, udis, pomariis Auſtriæ, Pannoniæ. Fl. Iunio, Iulio. Per

Folia omnia ſunt ſubbiternata; pinnulis oblongo-lanceolatis, acutis, ſerratis, ſubrugoſis, facie glabris & atro-virentibus, ſubtus ſcabriusculis, & ad nervos inprimis, venaſque piloſis, quibusdam pinnula minore pedatis. Floribus fertilibus immiſcentur flores germine carentes. Semina oblonga, glabra, fuſca, ſubacuta, hinc canaliculata, inde convexa, ſubteretia & obſolete quadriſulcata. Iacqu. l. c.

aureum. 6. C. caule æquali, foliolis inciſis, ſeminibus coloratis ſulcatis muticis. *Syſt. veget. p.* 288. *Iacqu. auſtr. t.* 64.

Cerefolium foliis hirſutis, petalis cordatis, ſeminibus obscure ſtriatis. *Hall. hiſt. n.* 749.

In Auſtriæ, Styriæ ſubalpinis, umbroſis. Fl. Iunio, Iulio. Per.

Caulis ſubſtriatus, ſub geniculis intumeſcens. Foliola ex oblongo lanceolata, pinnatifida, acuta, ſubtus hirſutula, extrema longe confluentia. Involucrum univerſale vel nullum, vel unifolium. Partiale habet foliola quinque aut ſex, lanceolato-acuminata, reflexa. Flos centralis breviſſime pedunculatus, pluresque ex circumferentia alii, ſunt hermaphroditi & fertiles. Hisce vero immiscentur alii germine deſtituti. Semina oblonga, glabra & quadrilinearia, intus, qua ſe mutuo tangunt, ſulco profundo exarantur; extus ſunt obſoletius angulata, fuſceſcentia, cum angulis pallidioribus. Iacqu. l. c.

IMPERATORIA. Fructus ſubrotundus, compreſſus, medio gibbus, marginatus. Petala inflexo-emarginata.

Oſtruthium. 1. Imperatoria. *Sp. pl. p.* 371. *Scop. carn. tom.* 1. *p.* 200. *Hall. hiſt. n.* 805. *Bauh. pin.* 156.

Selinum Imperatoria. *Crantz faſc. p.* 174.

Aſtrantia. *Cluſ. hiſt. p.* CXCIV.

L' Imperatoire. Regn. bot.

In alpibus ſtyriacis, carniolicis, carinthiacis, tyrolenſibus. Fl. Iulio, Auguſto. Per.

Radix articulata. Folia ternata: foliolis vel integris, vel lobatis, quandoque & partitis. Involucrum univerſale plerumque nullum. Partiale componitur ex foliolis tribus, vel quatuor, ſetaceis, caducis. Semina hinc ſulcata.

SESELI. Umbellæ globoſæ. Involucrum foliolo uno alterove. Fructus ovatus, ſtriatus.

glaucum. 1. S. petiolis ramiferis membranaceis oblongis integris: foliolis ſingularibus binatisque canaliculatis lævibus petiolo longioribus. *Sp. pl. p.* 372. *Iacqu. auſtr. t.* 144. *Scop. carn. tom.* 1. *p.* 214.

Seſeli oſſeum. *Crantz faſc. p.* 207.

In Carnioliæ, Auſtriæ ſaxoſis, herbidisque ſiccioribus. Fl. Iulio, Auguſto.

Folia radicalia ſunt bipinnata, aliqua tripinnata. Superiora pinnata, dein ternata, tandem ſimplicia. Petioli communes membranacei, canaliculati, lineati, reliqui ſubteretes. Pinnulæ ſunt lineares, anguſtæ, acutæ, ſubcanaliculatæ: plures etiam planiſſimæ. Involucrum univerſale nullum. Partiale conſtat foliolis aliquot ſetaceis. Floſculi omnes fertiles. Semina ovata, hinc plana, illinc convexa, haud profunde quinqueſulcata. Iacqu. l. c.

annuum. 2. S. petiolis rameis membranaceis ventricoſis emarginatis. *Sp. pl. p.* 373. *Iacqu. auſtr. t.* 55. *Scop. carn. tom.* 1. *p.* 213.

Seſeli bienne. *Crantz faſc. p.* 204.

In pratis, collibus montibusque apricis herbidis. Fl. Auguſto, Septembri.

Folia ſunt multiplicato - pinnata, facie glabra, dorſo margineque villoſa: pinnulis lanceolato - linearibus, acutis, planis. Involucrum univerſale ſæpius nullum, quandoque conſtans foliolo unico. Involucri univerſalis foliola ſunt plura, lanceolata, acuminata, margine membranacea. Flores omnes fertiles ſemina ferunt fusca, ovata, parva, hinc plana, illinc gibba atque ſtriis tribus elevatis notata, lateribusque inſuper membrana exigua acuta. Iacqu. l. c.

Hippomarathrum. 3. S. involucellis connato - monophyllis. *Sp. pl. p.* 374. *Iacqu. auſtr. t.* 143. *Scop. carn. tom.* 1. *p.* 214.

Seſeli articulatum. *Crantz faſc. p.* 205. *t.* 5. *f.* 1. 2.

Saxifraga pannonica. *Clus. hiſt. p.* CXCVI.

In apricis, aridis. Fl. Iulio, Auguſto. Per.

Involucrum partiale monophyllum & pelviforme, margine multidentatum. Fructus ſubovatus, ex fusco pallens ſecedit in duo ſemina oblonga vel ovata, obtuſiſſima & veluti truncata, vel acuta, hinc plana, illinc quinque coſtis extantibus notata. Iacqu. l. c.

PASTINACA. Fructus ellipticus, compreſſo-planus. Petala involuta, integra.

ſativa. 1. P. foliis ſimpliciter pinnatis. *Sp. pl. p.* 290. *Scop. carn. tom.* 1. *p.* 202.

Paſtinaca *ſativa*, foliis pinnatis ſubbipinnatisque auriculatis. *Retzii obs. faſc.* 5. *p.* 3. *prodr. p.* 69.

Paſtinaca foliis pinnatis. *Hall. hiſt. n.* 808.

Paſtinaca. *Riv. pent. t.* 6.

In pratis. Fl. Iulio, Auguſto.

SMYRNIUM. Fructus oblongus, ſtriatus. Petala acuminata, carinata.

perfoliatum. 1. S. foliis caulinis ſimplicibus amplexicaulibus. *Sp. pl. p.* 376. *Iacqu. collect. tom.* 4. *p.* 93.

In Pannonia, in monte *Kobel* prope Poſonium. *Iacquin filius.* Ego in agro fluminenſi in vineis, fruticoſis, & Terſacti in ſylva P. P. Franciſcanorum. Fl. Aprili, Majo, Iunio.

Folia radicalia ſupradecompoſita. Caulina ima triternata, ſuperiora ternata, reliqua, ut & ramea, ſimplicia, amplexicaulia. Umbellæ axillares, ſolitariæ, vel geminæ. Involucrum, & involucellum nullum. Fructus oblongus, ſtriatus.

ANETHUM. Fructus ſubovatus, compreſſus, ſtriatus. Petala involuta, integra.

Fœniculum. 1. A. fructibus ovatis. *Sp. pl. p.* 377. *Mill. illuſtr. ic.*

Fœniculum. *Riv. pent. t.* 60. *Hall. hiſt. n.* 760.

Le Fœnouil. Regn. bot.

In vineis, ſaxoſis aſperis Litoralis. Fl. æſtate. Per.

CARUM. Fructus ovato-oblongus, ſtriatus. Involucrum monophyllum. Petala carinata, inflexo-emarginata.

Carvi. 1. Carum. *Sp. pl. p.* 378. *Iacqu. auſtr. t.* 393.

Carum. *Riv. pent. t.* 54. *Hall. hiſt. n.* 789.

Seſeli Carum. *Scop. carn. tom.* I. *p.* 215.

Apium Carvi, *Crantz faſc. p.* 218.

In pratis, aliisque herbidis locis. Fl. Iunio, Iulio.

Folia ſunt plus minus bipinnata, aucta ad coſtam principem foliolis multifidis, cruciatim poſitis. Involucrum univerſale vel nullum, vel conſtans foliolo uno alterove ſubulato. Particulare nullum. Omnes floſculi ſunt fertiles. Semina oblonga, verſus utrumque finem attenuata & quinqueſtriata. Iacqu. l. c.

PIMPINELLA. Fructus ovato-oblongus. Stigmata ſubgloboſa.

Saxifraga. 1. P. foliis pinnatis: foliolis radicalibus ſubrotundis; ſummis linearibus. *Sp. pl. p.* 378. *Iacqu. auſtr. t.* 395.

Tragoſelinum foliis pinnatis, pinnis ovatis. *Hall. hiſt. n.* 786.

Pimpinella rotundifolia. *Scop. carn. tom.* I. *p.* 208.

In pratis ſiccioribus, collibus apricis, ad vias. Fl. Auguſto, Septembri. Per.

magna. 2. P. foliolis omnibus lobatis: impari trilobo. *Syſt. veget. p.* 291.

Tragoſelinum foliis pinnatis, pinnis lanceolatis ſerratis, extremis trilobatis. *Hall. hiſt. n.* 785.

Pimpinella major. *Iacqu. auſtr. t.* 396.

Pimpinnella orientalis. *Gouan. illuſtr. p.* 21. *t.* 15. *Iacqu. auſtr. t.* 397.

In ſylvaticis. Fl. æſtate. Per.

diſſecta. 3. P. foliis omnibus pinnatis, pinnis multipartitis, ſegmentis ſubfalcatis acutis. *Retzii obs. faſc.* 3. *p.* 30. *t.* 2.

Tragoſelinum foliis duplicato-pinnatis, pinnulis profundiſſime lobatis. *Hall. hiſt. n.* 787.

In alpibus ſtyriacis, auſtriacis. Fl. Iulio, Auguſto. Per.

glauca. 4. P. foliis ſupradecompoſitis, caule angulato ramoſiſſimo. *Sp. pl. p.* 378.

Paſtinaca glauca. *Scop. carn. tom.* 1. *p.* 203.

In Carniolia in alpibus wochinenſibus. Fl. Iulio, Auguſto. Per.

Folia ſunt lineari-lanceolata, glauca. Involucra nulla. Flores omnes fertiles ferunt fructum ellipticum conſtantem ſeminibus duobus, hinc planis, inde convexis, & tribus lineis elevatis notatis.

dioica. 5. P. umbellis numeroſiſſimis compoſitis ſimplicibusque, *Syſt. veget. p.* 291.

Pimpinella pumila. *Iacqu. auſtr. t.* 28.

Apium pumilum. *Crantz faſc. p.* 213.

Selinum montanum pumilum. *Cluſ. hiſt. p.* CC.

In collibus, montoſisque herbidis, nec non in pratis ſiccioribus Auſtriæ, Pannoniæ. Fl. Majo.

Foliola linearia, craſſa, antice ſulcata. Involucrum nullum. Floribus maribus piſtillum deeſt. Hermaphroditi ferunt fructum ſubrotundum, obtuſum, glabrum, plerumque decem-ſulcatum. Iacqu. l. c.

APIUM. Fructus ovatus, ſtriatus. Involucrum monophyllum. Petala æqualia.

graveolens. 1. A. foliis caulinis cuneiformibus. *Syſt. veget. p.* 292. *Scop. carn. tom.* 1. *p.* 215. *Fl. dan. t.* 790.

L'Ache. Regn. bot.

In paluſtribus aquilejenſibus, duinenſibusque. Occurrit quoque in humidis Auſtriæ ad Thermas badenſes, circa *Galbrun*, Fl. Iunio.

AEGOPODIUM. Fructus ovato-oblongus, ſtriatus.

Podagraria. 1. Ae. foliis caulinis ſummis ternatis. *Sp. pl. p.* 379. *Scop. carn. tom.* 1. *p.* 215. *Fl. dan. t.* 670.

Podagraria. *Riv. pent. t.* 46. *Hall. hiſt. n.* 759.

In pratis udis, cultis, ſylvis, ad rivulos, ſepes. Fl. Iunio. Per.

TRIGYNIA.

RHUS. Calyx quinquepartitus. Petala quinque. Bacca monoſperma.

Cotinus. 1. R. foliis ſimplicibus obovatis. *Sp. pl. p.* 383. *Iacqu. auſtr. t.* 210.

Rhus racemis plumoſis, foliis ovatis. *Hall. hiſt. n.* 827.

Cotinus Coggygria. *Scop. carn. tom.* 1. *p.* 220.

Coggygria. *Cluſ. hiſt. p.* 16.

In Auſtria in montanis ad Thermas badenſes. In Forojulio, Iſtria, Litorali ad ſepes, vineas, alibique. Fl. Majo. Fr.

Foliis ramisque conſciſſis ad coria perficienda utuntur coriarii liburni.

VIBURNUM. Calyx quinquepartitus, ſuperus. Corolla quinquefida. Bacca monoſperma.

Lantana. 1. V. foliis cordatis ſerratis venoſis, ſubtus tomentoſis. *Sp. pl. p.* 384. *Iacqu. auſtr. t.* 341. *Scop. carn. tom.* 1. *p.* 221.

Viburnum foliis ovato-ſerratis, ſubtus tomentoſis. *Hall. hiſt. n.* 669.

La Viorne. Regn. bot.

In ſylvis, fruticoſis, ad ſepes. Fl. Majo. Fr.

Opulus. 2. V. foliis lobatis, petiolis glanduloſis. *Sp. pl. p.* 384. *Fl. dan. t.* 661. *Scop. carn. tom.* 1. *p.* 297. *Hall. hiſt. n.* 668.

In ſylvis. Fl. Majo, Iunio. Fr.

SAMBUCUS. Calyx quinquepartitus. Corolla quinquefida. Bacca triſperma.

Ebulus. 1. S. cymis tripartitis, ſtipulis foliaceis, caule herbaceo. *Sp. pl. p.* 385. *Curtis lond. Scop. carn. tom.* 1. *p.* 222.

Sambucus herbacea, floribus umbellatis. *Hall. hiſt. n.* 671.

L' Hieble. Regn. bot.

Ad vias, pagos, agrorum margines, in ſylvis. Fl. Iunio, Iulio. Per.

nigra. 2. S. cymis quinquepartitis, foliis pinnatis, foliolis ſubovatis ſerratis, caule arboreo. *Syſt. veget. p.* 295. *Fl. dan. t.* 545. *Scop. carn. tom.* 1. *p.* 222.

Sambucus arborea floribus umbellatis. *Hall. hiſt. n.* 670.

Le Sureau. Regn. bot.

In ſylvis, ad ſepes, pagos. Fl. Majo, Iunio.

racemoſa. 3. S. racemis compoſitis ovatis, caule arboreo. *Sp. pl. p.* 386. *Scop. carn. tom.* 1. *p.* 223. *Iacqu. collect. tom.* 1. *p.* 36. *ic. t.* 59.

Sambucus arborea floribus ſpicatis. *Hall. hiſt. n.* 672.

In ſubalpinis. Fl. Majo, Iunio.

STAPHYLEA. Calyx quinquepartitus. Petala quinque. Nectarium e receptaculo fructificationis, concavum, urceolatum. Capſulæ inflatæ, connatæ. Semina duo, globoſa cum cicatrice.

pinnata. 1. S. foliis pinnatis. *Sp. pl. p.* 386. *Hall. hiſt. n.* 831.

Staphyllodendron pinnatum. *Scop. carn. tom.* 1. *p.* 223.

Staphyllodendron. *Besl. aichſt. vern. ord. frut. t.* 8. *f.* 1.

In ſylvis. Fl. Majo, Iunio. Fr.

Numerus ternarius piſtilli & pericarpii evadit binarius. Linn.

TAMARIX. Calyx quinquepartitus. Petala quinque. Capſula unilocularis, trivalvis. Semina pappoſa.

gallica. 1. T. floribus pentandris. *Sp. pl. p.* 386. *Miller. ic. t.* 262. *f.* 1.

Tamarix altera, folio tenuiore, ſive gallica. *Bauh. pin.* 485. *Zann. iſt. p.* 256. *t.* 186.

Myrica ſylveſtris prima. *Clus. hiſt. p.* 40.

Ad litus maris adriatici in inſulis venetis. Fl. æſtate. Fr.

germanica. 2. T. floribus decandris. *Sp. pl. p.* 387. *Fl. dan. t.* 234. *Miller. ic. t.* 262. *f.* 2.

Tamariscus ſpicis folioſis. *Hall. hiſt. n.* 948.

Tamariscus germanica. *Scop. carn. tom.* 1. *p.* 224.

Tamarix fruticoſa folio craſſiore, ſeu germanica. *Bauh. pin.* 485.

Myrica ſylveſtris ſecunda. *Clus. hiſt. p.* 40.

In arenoſis ad fluviorum ripas. Fl. Iunio, Iulio. Fr.

Tamarix gallica habet folia lanceolato - acuminata, ſtamina quinque, libera; germanicæ contra ſunt folia linearia, obtuſa, ſtamina decem, baſi connata.

CORRIGIOLA. Calyx pentaphyllus. Petala quinque. Semen unicum, triquetrum.

litoralis. 1. Corrigiola. *Sp. pl. p.* 388. *Fl. dan. t.* 334. *Hall. hiſt. n.* 842.

Polygonum litoreum minus, floſculis ſpadiceo-albicantibus. *Bauh. pin.* 281. *prodr.* 131.

Tergeſti in arenoſis *alle Saule. I. Hoſt.* Fl. æſtate.

ALSINE. Calyx pentaphyllus. Petala quinque, æqualia. Capſula unilocularis, trivalvis.

media. 1. A. petalis bipartitis, foliis ovato - cordatis. *Sp. pl. p.* 389. *Curtis lond. Fl. dan. t.* 525 & 438. *Scop. carn. tom.* 1. *p.* 224.

Alſine foliis petiolatis, ovato - lanceolatis, petalis bipartitis. *Hall. hiſt. n.* 880.

La Morgeline. Regn. bot.

In cultis, ruderatis, & alibi. Fl. a primo vere ad hyemem.

DRYPIS. Calyx quinquedentatus. Petala quinque. Capſula circumſciſſa, monoſperma.

ſpinoſa. 1. Drypis. *Sp. pl. p.* 390. *Iacqu. hort. tom.* 1. *p.* 19. *t.* 49. *Scop. carn. tom.* 1. *p.* 225.

Drypis italica aculeata, floribus albis umbellatis compactis. *Mich. gen. p.* 24. *t.* 23.

Spina alba, foliis vidua. *Bauh. pin.* 388.

In agro fluminenſi ad litus maris *al Ponſal, nella valle di S. Martin.* In Carniolia in monte *Nanas*, & *Grindovitz*. Fl. Iunio, Iulio. Per.

Folia ſunt ſubulata, mucronato-pungentia. Superiora ad oras ciliato-ſpinoſa.

TETRAGYNIA.

PARNASSIA. Calyx quinquepartitus. Petala quinque. Nectaria quinque, cordata, ciliata apicibus globoſis. Capſula quadrivalvis.

paluſtris. 1. Parnaſſia. *Sp. pl. p.* 391. *Fl. dan. t.* 584. *Scop. carn. tom.* 1. *p.* 226. *Hall. hiſt. n.* 832.

Gramen Parnaſſi. *Lob. ic. p.* 603.

In pratis udis demiſſis, nec non in alpinis, & ſubalpinis. Fl. æſtate. Per.

Folia integerrima: radicalia petiolata; caulina ſeſſilia, amplexicaulia.

PENTAGYNIA.

STATICE. Calyx monophyllus, integer, plicatus, ſcarioſus. Petala quinque. Semen unicum, ſuperum.

Armeria. 1. S. ſcapo ſimplici capitato, foliis linearibus. *Sp. pl. p.* 394. *Scop. carn. tom.* 1. *p.* 227. *Hall. hiſt. n.* 835.

Caryophyllus nonus. *Clus. hiſt. p.* 286.

In alpium herbidis. Fl. æſtate. Per.

Limonium. 2. S. ſcapo paniculato tereti, foliis lævibus enerviis, ſubtus mucronatis. *Syſt. veget. p.* 300. *Fl. dan. t.* 315. *Scop. carn. tom.* 1. *p.* 227.

Limonium maritimum majus. *Bauh. pin.* 192. *Zann. iſt. p.* 164. *t.* 31.

In Litorali in paluſtribus agri aquilejenſis, duinenſisque. Fl. Iulio, Auguſto, Septembri. Per.

oleifolia. 3. S. foliis lanceolatis ſubariſtatis; margine cartilagineo; caulibus dichotomis, flexuoſis, ſubnudis. *Scop. inſubr.* I. *p.* 24. *t.* 10.

Limonium maritimum minus oleæ folio. *Magn. bot.* 155. *hort.* 119. *Zann. iſt. p.* 165. *t.* 2. 32.

In agro fluminenſi in rupeſtribus ad mare. Fl. Auguſto, Septembri. Per.

LINUM. Calyx pentaphyllus. Petala quinque. Capſula decemvalvis, apice dehiſcens. Diſſepimenta membranacea, tenuiſſima, valvulas connectentia. Semina ſolitaria.

uſitatiſſimum. I. L. calycibus capſulisque mucronatis, petalis crenatis, foliis lanceolatis alternis, caule ſubſolitario. *Sp. pl. p.* 397. *Curtis loud. Scop. carn. tom.* I. *p.* 228. *Hall. hiſt. n.* 836.

Le Lin. Regn. bot.

Inter ſegetes, ad pagos. Fl. æſtate.

viſcoſum. 2. L. foliis lanceolatis piloſis quinquenerviis. *Sp. pl. p.* 398.

Linum ſylveſtre. *Scop. carn. tom.* I. *p.* 229. *t.* 11.

Linum ſylveſtre latifolium, caule viſcoſo, flore rubro. *Bauh. pin.* 214.

In montanis Carinthiæ, Forijulii, Carnioliæ, Croatiæ. Fl. Iulio, Auguſto. Per.

hirſutum. 3. L. calycibus hirſutis, acuminatis, ſeſſilibus, alternis, foliis caulinis alternis: rameis oppoſitis. *Syſt. veget. p.* 302. *Iacqu. auſtr. t.* 31.

Linum ſylveſtre latifolium primum. *Clus. hiſt. p.* 316.

In Auſtriæ, Pannoniæ collibus apricis aſperis, pratisque ſiccioribus. Fl. Iunio, Iulio. Per.

narbonenſe. 4. L. calycibus acuminatis, foliis lanceolatis ſparſis ſtrictis ſcabris acuminatis, caule tereti baſi ramoſo. *Sp. pl. p.* 398.

Linum ſylveſtre cœruleum, folio acuto. *Bauh. pin.* 214. *prodr.* 107.

In Croatia in pratis montium, *mala Planina* dictorum. Fl. Iunio, Iulio. Per.

tenuifolium. 5. L. calycibus acuminatis, foliis ſparſis lineari-ſetaceis retrorſum ſcabris. *Sp. pl. p.* 398. *Iacqu. auſtr. t.* 215. *Scop. carn. tom.* 1. *p.* 230.

Linum foliis linearibus, calycibus longe ariſtatis. *Hall. hiſt. n.* 838.

Linum ſylveſtre quintum, anguſtifolium ſecundum. *Clus. hiſt. p.* 318.

In collibus apricis. Fl. æſtate. Per.

Folia ſunt ſpinulis albidis ſurſum arcuatis ciliata. Calycis foliola ovata, ariſtato-acuminata, pilis glanduliferis ciliata. Iacqu. l. c.

gallicum. 6. L. calycibus acutis, foliis lineari-lanceolatis alternis, paniculæ pedunculis bifloris. *Sp. pl. p.* 401. *Ger. prov. p.* 421. *t.* 15. *f.* 1.

Linum ſylveſtre minus, flore luteo. *Bauh. pin.* 214.

In Carniolia in monte *Nanas. I. Hoſt.* Fl. Iunio.

maritimum. 7. L. calycibus ovatis acutis muticis, foliis lanceolatis: inferioribus oppoſitis. *Sp. pl. p.* 400. *Iacqu. hort. tom.* 2. *p.* 72. *t.* 154.

Linum liburnicum. *Scop. carn. tom.* 1. *p.* 230.

Linum maritimum luteum. *Bauh. pin.* 214. *Zann. iſt. p.* 168. *t.* 78.

In arenoſis Litoralis ad mare. Fl. Auguſto, Septembri. Per.

alpinum. 8. L. calycibus rotundatis obtuſis, foliis linearibus acutiuſculis, caulibus declinatis. *Sp. pl. app. p.* 1672. *Iacqu. auſtr. t.* 321.

Lini ſylveſtris quinti anguſtifolii aliud genus. *Cluſ. hiſt. p.* 318.

In alpibus. Fl. æſtate. Per.

auſtriacum. 9. L. calycibus rotundatis obtuſis, foliis linearibus acutis rectiuſculis. *Syſt. veget. p.* 302. *Iacqu. auſtr. t.* 418.

Linum ſylveſtre quartum, anguſtifolium primum. *Cluſ. hiſt. p.* 318.

In Auſtria, Pannonia ad vias, agrorum margines, in collibus apricis. Fl. æſtate. Per.

Caules procumbentes. Pedunculi filiformes, uniflori. Calycis foliola obtuſa. Petala cærulea donantur ungue flavo. Iacqu. l. c.

flavum. 10. L. calycibus ſubſerrato-ſcabris lanceolatis ſubſeſſilibus, panicula ramis dichotomis. *Sp. pl. p.* 399. *Iacqu. auſtr. t.* 214. *Scop. carn. tom.* 1. *p.* 230.

Linum ſylveſtre tertium latifolium. *Cluſ. hiſt. p.* 317.

In pratis ſiccioribus, collibus, montoſis. Fl. æſtate. Per.

Folia ad caulem utrinque inſtructa ſunt glandula ſeſſili & badia. Calycis foliola acuminata, ſerrato-ſpinoſa. Petala flava. Stamina quinque alterna calyce duplo longiora, fertilia; alia quinque breviſſima & ſterilia. Iacqu. l. c.

ſtrictum. 11. L. calycibus ſubulatis, foliis lanceolatis ſtrictis mucronatis margine ſcabris. *Sp. pl. p.* 400.

Linum foliis aſperis umbellatum luteum. *Magn. monſp.* 164.

In Carnia circa *Artègnan. I. Hoſt.* Fl. Iunio, Iulio.

Caulis ſtriatus, ſuperne ramoſus. Folia lanceolata, ſtricta, margine aculeis ſurſum verſis ſcabra, mucronata. Rami alterni. Flores breviter pedunculati. Calycis foliola baſi dilatata, margine apicem verſus ſerrato-ſpinoſa, capſula duplo longiora.

cathar- ticum. 12. L. foliis oppoſitis ovato-lanceolatis, caule dichotomo corollis acutis. *Sp. pl. p.* 401. *Fl. dan. t.* 851. *Curtis lond. Scop. carn. tom.* I. *p.* 231.

Linum foliis conjugatis, ovatis, calycibus ariſtatis, patulis, lanceolatis. *Hall. hiſt. n.* 839.

Linum ſylveſtre octavum, Chamælinum. *Clus. hiſt. p.* 319.

In pratis. Fl. Iunio, Iulio.

DROSERA. Calyx quinquefidus. Petala quinque. Capſula unilocularis, apice tri- vel quinquevalvis. Semina plurima.

rotundi- folia. 1. D. ſcapis radicatis, foliis orbiculatis. *Sp. pl. p.* 402. *Fl. dan. t.* 1028.

Rorella caule nudo, paucifloro, foliis petiolatis ſubrotundis. *Hall. hiſt. n.* 834.

Salſirora ſeu Ros ſolis. *Thal. herc. t.* 9. *f.* 1.

In paluſtribus Auſtriæ prope *Annaberg*, Carnioliæ, Forijulii, Carniæ. Fl. Iunio, Iulio.

longifo- lia. 2. D. ſcapis radicatis, foliis oblongis. *Sp. pl. p.* 403.

Rorella foliis ellipticis, caule nudo paucifloro. *Hall. hiſt. n.* 833.

Salſirora ſeu ſponſa ſolis. *Thal. herc. t.* 9. *f.* 2.

Habitat & floret cum præcedente.

SIBBALDIA. Calyx decemfidus. Petala quinque, calyci inſerta. Styli e latere germinis. Semina quinque.

procum- bens. 1. S. foliis tridentatis. *Sp. pl. p.* 406. *Fl. dan. t.* 32.

Fragaria foliis ternatis, retuſis, tridentatis, flore calyci æquali, pentaſtemone. *Hall. hiſt. n.* 1116.

Fragariæ ſylveſtri affinis planta, flore luteo. *Sibb. ſcott.* 2. *p.* 25. *t.* 6. *f.* 1.

In alpibus Tyrolis, Carinthiæ, Styriæ, Carnioliæ, præcipue locis editiſſimis humidis. Fl. æſtate. Per.

POLYGYNIA.

MYOSURUS. Calyx pentaphyllus. Petala quinque, baſi tubulata, oblique introrſum aperta. Stamina quinque vel plura. Germina numeroſa. Receptaculum longiſſimum, ſeminibus imbricatim diſpoſitis tectum.

minimus. 1. Myoſurus. *Sp. pl. p.* 407. *Fl. dan. t.* 406. *Hall. hiſt. n.* 1159.

Cauda muris. *Lob. ic. p.* 40.

In Pannonia ad litus lacus Peiſonis. In Auſtria in herbidis agri vindobonenſis inter *Simeninger-* & *Laacherwald*, in paluſtribus prope *Erla*, alibique. Fl. Majo.

Stamina in noſtra planta nunquam quinque, ſed plura, numeravi.

HEXANDRIA.

MONOGYNIA.

GALANTHUS. Petala tria, concava. Nectarium ex petalis tribus, parvis, emarginatis. Stigma ſimplex.

nivalis. 1. Galanthus. *Sp. pl. p.* 413. *Iacqu. auſtr. t.* 313. *Scop. carn. tom.* 1. *p.* 233.

Galanthus uniflorus petalis alternis cordatis brevioribus. *Hall. hiſt. n.* 1254.

Leucojum bulboſum præcox minus. *Clus. hiſt. p.* 169.

In ſylvis, montium declivibus, pratis umbroſis. Fl. a Februario ad Aprilim. Per.

LEUCOJUM. Corolla campaniformis, ſexpartita, apicibus incraſſata. Stigma ſimplex.

vernum. 1. L. Spatha uniflora, ſtylo clavato. *Sp. pl. p.* 414. *Iacqu. auſtr. t.* 312. *Scop. carn. tom.* 1. *p.* 234.

Galanthus uniflorus, petalis ſubæqualibus. *Hall. hiſt. n.* 1253.

Leucojum bulboſum. *Clus. hiſt. p.* 168.

In Carnioliæ pratis circa Labacum, & *Ober-laybach.* In Auſtria circa *Krems;* in agro vindobonenſi in ſylva prope *Himberg.* Fl. primo vere. Per.

æſtivum. 2. L. ſpatha multiflora, ſtylo clavato. *Sp. pl. p.* 414. *Iacqu. auſtr. t.* 203. *Scop. carn. tom.* 1. *p.* 234.

Leucojum bulboſum ſerotinum majus primum. *Clus. hiſt. p.* 170.

In Auſtria in pratis prope *Himberg, Lachſenburg.* In Carniolia circa Labacum. Copioſiſſime in Croatia circa Zagrabiam, alibique. Fl. Majo. Per.

NARCISSUS. Petala ſex, æqualia: Nectario infundibuliformi, monophyllo. Stamina intra nectarium.

poëticus. 1. N. ſpatha uniflora, nectario rotato breviſſimo ſcarioſo crenulato. *Sp. pl. p.* 414. *Scop. carn. tom.* 1. *p.* 235.

Narciſſus uniflorus, foliis enſiformibus, ſcypho breviſſimo. *Hall. hiſt. n.* 1250.

Narciſſus albus, circulo purpureo. *Bauh. pin.* 48.

Le Narciſſe des Poetes. Bulliard herb. t. 306.

In Carnioliæ calidioris herbidis ſiccis. In Litorali in pratis montanis. Fl. primo vere. Per.

ALLIUM. Corolla ſexpartita, patens. Spatha multiflora. Umbella congeſta. Capſula ſupera.

Victorialis. 1. A. caule planifolio umbellifero, umbella rotundata, ſtaminibus lanceolatis corolla longioribus, foliis ellipticis. *Sp. pl. p.* 424. *Iacqu. auſtr. t.* 216.

Victorialis longa. *Clus. hiſt. p.* 189.

In alpium herbidis. Fl. Iulio, Auguſto. Per.

roſeum. 2. A. caule planifolio umbellifero, umbella faſtigiata, petalis emarginatis, ſtaminibus breviſſimis ſimplicibus. *Syſt. veget. p.* 321.

Allium caule planifolio umbellifero, umbella faſtigiata. filamentis ſimplicibus baſi dilatatis. *Ger. prov. p.* 150. *Gouan. monſp. p.* 166.

Allium illyricum. *Iacqu. collect. tom.* 3. *p.* 273. *ic. t.* 365.

Allium foliis enſiformibus, radice bulbis ſtipata, floribus amplis umbellatis. *Sauv. monſp. p.* 42.

Allium ſylveſtre, ſeu molli minus, roſeo flore. *Magn. monſp.* 2. *t.* 10.

In vineis, olivetis Literalis. Fl. Majo. Per.

Bulbus ſubrotundus, bulbillis cinctus. Folia inferna parte canaliculata caulem ſic amplexantia, ut minime radicalia, ſed caulina appareant. Superne plana cum dorſo leviter carinato. Scapus unus alterve, teres. Umbella faſtigiata, nec bulbifera. Spatha in duas tresve ſecedens valvulas acuminatas: acumine ſæpe deciduo. Pedunculi primum breves, dein elongantur. Petala lanceolata, vel ovata, apice acuta, vel obtuſa, integra, vel emarginata. Filamenta ſubulata, baſi dilatata. Capſula ovata.

carinatum. 3. A. caule planifolio bulbifero, ſtaminibus ſubulatis. *Sp. pl. p.* 426.

Allium ſeu Moly montanum ſecundum. *Cluſ. hiſt. p.* 193.

In Auſtria, Pannonia ad agrorum, vinearumque margines, in fruticoſis collibus. Fl. Iunio. Per.

ſphærocephalon. 4. A. caule teretifolio umbellifero, foliis ſemiteretibus, ſtaminibus tricuſpidatis corolla longioribus. *Sp. pl. p.* 426.

Allium ſeu Moly montanum quintum. *Cluſ. hiſt. p.* 195.

Cum præcedente. Fl. Majo, Iunio, Per.

flavum. 5. A. caule teretifolio umbellifero, floribus pendulis, petalis ovatis, ſtaminibus corolla longioribus. *Sp. pl. p.* 428. *Iacqu. auſtr. t.* 141.

Allii montani quarti ſpecies prima. *Cluſ. hiſt. p.* 194.

In Auſtriæ collibus aſperis, ad margines vinearum. Fl. Iulio, Auguſto. Per.

paniculatum. 6. A. caule ſubteretifolio umbellifero, pedunculis capillaribus effuſis, ſtaminibus ſubulatis, ſpatha longiſſima. *Sp. pl. p.* 428.

Habitat & floret cum præcedente. Per.

ſuaveolens. 7. A. ſcapo nudo ſubtereti, foliis linearibus canaliculatis, umbella ſubrotunda. *Iacqu. collect. tom.* 2. *p.* 305. *ic. t.* 364.

Allium maritimum venetum, autumnale, anguſtioribus foliis, flore purpuraſcente. *Zann. iſt. t. 11. 191.*

In pratis paludoſis Auſtriæ prope *Erla, Himberg, Lachſenburg.* In Litorali in paluſtribus aquilejenſibus ad fluvium *Anfora.* Fl. a Iulio ad Septembrim. Per.

Bulbus oblongus. Scapus teres cum uno alalterove angulo obſoleto. Folia linearia acuta, integerrima, canaliculata, baſibus vaginantibus connexa. Spatha bivalvis. Flores ſuaveolentes. Petala alba cum ſuffuſa purpura, ovata, acutiuſcula, tria alterna exteriora breviora. Germen obſcure triquetrum. Capſula ovata, obtuſa. Iacqu. l. c.

angulo-ſum. 8. A. ſcapo nudo ancipiti, foliis linearibus canaliculatis ſubtus ſubangulatis, umbella faſtigiata. *Sp. pl. p. 430. Iacqu. auſtr. t. 423.*

Allium pratenſe, radice iridis, flore purpuraſcente. *Zann. iſt. p. 7. t. 273. n. 1.*

Allium ſive Moly narciſſinis foliis primum & ſecundum. *Cluſ. hiſt. p. 195.*

In montanis, rupeſtribus Auſtriæ. In Litorali & in ſimilibus locis, & in pratis paluſtribus Duini, Aquilejæ. Fl. cum præc. Per.

nigrum. 9. A. ſcapo nudo tereti, foliis linearibus, umbella hemiſphærica, petalis erectis, ſpatha mucronata bifida. *Sp. pl. p. 430. Retzii obſ. faſc. 1. p. 15.*

Allium multibulboſum. *Iacqu. auſtr. t. 10.*

Allium monſpeſſulanum. *Gouan. illuſtr. p. 24. t. 16.*

In agro vindobonenſi in monte cætio. Fl. Iunio. Per.

Folia ſunt plana, ſtriata, acuta, ex lato enſiformia. Scapus bipedalis, teres. Umbella denſa, leviter convexa. Petala albeſcunt cum linea dorſali virente, ſuntque ovalia, obtuſa. Filamenta baſibus ſuis in unum corpus coalita. Germen ex atro viret. Iacqu. l. c.

urſinum. 10. A. ſcapo nudo ſubtriquetro, foliis lanceolatis petiolatis, umbella faſtigiata. *Retzii ſcand. p.* 76. *Sp. pl. p.* 431. *Fl. dan. t.* 757.

In umbroſis. Fl. primo vere. Per.

ſchœnoprasum. 11. A. radice multiplici, foliis radicalibus teretibus ſcapi longitudine; floribus umbellatis, cylindricis. *Hall. hiſt. n.* 1226. *Sp. pl. p.* 432. *Murr. in comm. gott.* 1775. *p.* 33. *t.* 4.

In alpibus ſtyriacis, carniolicis, carinthiacis, tyrolenſibus præcipue locis humidis. Fl. Iulio, Auguſto. Per.

LILIUM. Corolla hexapetala, campanulata: linea longitudinali nectarifera. Capſula valvulis pilo cancellato connexis.

bulbiferum. 1. A. foliis ſparſis, corollis campanulatis erectis: intus ſcabris. *Sp. pl. p.* 433. *Iacqu. auſtr. t.* 226. *Scop. carn. tom.* 1. *p.* 240.

Lilium caule folioſo, foliis ſulcatis, floribus campanulatis, extus floccoſis. *Hall. hiſt. n.* 1232.

Martagon bulbiferum tertium. *Clus. hiſt. p.* 136.

In Auſtriæ, Styriæ pratis ſubalpinis. In Carnioliæ, & agri fluminenſis montoſis, herbidis. Fl. Iunio, Iulio. Per.

chalcedonicum. 2. L. foliis ſparſis lanceolatis, floribus reflexis, corollis revolutis. *Sp. pl. p.* 434. *Iacqu. auſtr. app. t.* 20. *Scop. carn. tom.* 1. *p.* 240.

Lilium rubrum ſive miniatum byzantinum. *Clus. hiſt. p.* 131.

In Carniolia in herbidis alpium wochinenſium, tolmienſium, montis *Nanas*, & totius Planinæ. In Litorali in monte majore. Fl. Iunio, Iulio. Per.

Martagon. 3. L. foliis verticillatis, floribus reflexis, corollis revolutis. *Sp. pl. p.* 435. *Iacqu. auſtr. t.* 351. *Scop. carn. tom.* 1. *p.* 239.

Lilium foliis verticillatis, floribus pendulis revolutis. *Hall. hiſt. n.* 1233.

Lilium montanum ſive ſylveſtre primum majus, & ſecundum minus. *Clus. hiſt. p.* 134.

In ſylvis montanis, & ſubalpinis. Fl. æſtate. Per.

FRITILLARIA. Corolla hexapetala, campanulata, ſupra ungues cavitate nectarifera. Stamina longitudine corollæ.

Meleagris. 1. F. caule ſubunifloro, foliis omnibus alternis. *Sp. pl. p.* 436. *Iacqu. auſtr. app. t.* 32. *Curtis lond. Fl. dan. t.* 972. *Scop. carn. tom.* 1. *p.* 241.

Fritillaria caule paucifloro, foliis caulinis gramineis alternis. *Hall. hiſt. n.* 1235.

In Carniolia in pratis humidis circa Labacum, & *Oberlaibach*. In Litorali in umbroſis montis majoris. Fl. Martio, Aprili. Per.

UVULARIA. Corolla hexapetala, erecta. Fovea oblonga baſi ſinguli petali inſculpta. Filamenta breviſſima.

amplexifolia. 1. U. foliis amplexicaulibus mollibus, caule glabro. *Syſt. veget. p.* 325. *Hall. hiſt. n.* 1237.

Polygonum latifolium quartum ramoſum. *Clus. hiſt. p.* 276.

In Auſtria in umbroſis montis *Wechſel* ſylvis, & *Dürrenſtein* inter ſaxa & frutices. *Clus.* In umbroſis Carnioliæ ſylvis ſubalpinis. *Zois.* Fl. Iunio, Iulio. Per.

ERYTHRONIUM. Corolla hexapetala, campanulata. Tubercula duo calloſa ſingulo petalo alterno & interiori juxta baſin adnata.

Dens canis. 1. Erythronium. *Sp. pl. p.* 437. *Iacqu. auſtr. app. t.* 9. *Scop. carn. tom.* 1. *p.* 242. *Hall. hiſt. n.* 1234.

Dens caninus. *Clus. hiſt p.* 266.

In ſylvaticis Styriæ, Croatiæ, Carnioliæ, Litoralis. Fl. primo vere. Per.

TULIPA. Corolla hexapetala, campanulata. Stylus nullus.

ſylveſtris. 1. T. florê ſubnutante, foliis lanceolàtis. *Sp. pl. p.* 437. *Fl. dan. t.* 375. *Scop. carn. tom.* 1. *p.* 242. *Hall. hiſt. n.* 1236.

In Litorali in monte majôre. In agro vindobonenſi in hortorum ambulacris. Fl. primo vere. Per.

ORNITHOGALUM. Corolla hexapetala, erecta, perſiſtens, ſupra medium patens. Filamenta alterna baſi dilatata.

luteum. 1. O. ſcapo anguloſo diphyllo, pedunculis umbellatis. *Fl. dan. t.* 378. *Sp. pl. p.* 439. *Allion. pedem. tom.* 2. *p.* 163. *Scop. carn. tom.* 1. *p.* 244.

Ornithogalum minimum. *Sp. pl. p.* 440. *Schreb. ſpic. p.* 35.

Ornithogalum pallido flore. *Clus. hiſt. p.* 188.

Ornithogalum pannonicum luteo flore. *Clus. hiſt. p.* 189.

In ſylvis, apricis campeſtribus, cultis, alibique. Fl. primo vere. Per.

Variat, prout varia eſt ſoli natura, ac expoſitio, foliis planis, canaliculato-triquetris, glabris, hirſutis; pedunculis ſimplicibus, & ramoſis; petalis obtuſis, acutisque. Linnæus ſub Ornithogalo minimo citat Ornithogalum minus Bauhini, quod eſt O. pannonicum luteo flore Cluſii. Plantam cluſianam jam per plures annos in apricis agri vindobonenſis inter Simeninger- & Laacherwald *attente examinavi. Inveni individua ſcapo unifloro, bifloro, multifloro; pedunculis ſimplicibus, & ramoſis.*

pyrenaicum. 2. O. racemo longiſſimo, filamentis lanceolatis, pedunculis floriferis, patentibus æqualibus: fructiferis ſcapo approximatis. *Sp. pl. p.* 440. *Iacqu. auſtr. t.* 103. *Scop. carn. tom.* 1. *p.* 244. *Gouan. illuſtr. p.* 26.

Phalangium longiſſime ſpicatum, filamentis latis, lanceolatis. *Hall. hiſt. n.* 1210.

Ornithogalum majus primum. *Clus. hiſt. p.* 187.

In Auſtriæ, Carnioliæ calidioris pratis infrequens. In vineis agri fluminenſis copioſiſſime. Fl. Majo, Iunio. Per.

comoſum. 3. O. racemo breviſſimo, bracteis lanceolatis longitudine florum, petalis obtuſis, filamentis ſubulatis. *Sp. pl. p.* 440. *Iacqu. collect. tom.* 2. *p.* 313. *ic. t.* 426.

Ornithogalum pannonicum albo flore. *Clus. hiſt. p.* 187.

In Auſtriæ apricis ad Thermas badenſes; inter *Simeninger & Laacherwald*, alibique. Fl. Iunio. Per.

umbellatum. 4. O. floribus corymboſis, pedunculis ſcapo altioribus, filamentis emarginatis. *Sp. pl. p.* 441. *Iacqu. auſtr. t.* 343. *Scop. carn. tom.* 1. *p.* 243.

Ornithogalum ſtipulis maximis, petiolis lateralibus longiſſimis. *Hall. hiſt. n.* 1215.

Ornithogalum vulgare. *Clus. hiſt. p.* 188.

In cultis, pratis, apricis aſperis. Fl. Majo, Iunio. Per.

nutans. 5. O. floribus ſecundis pendulis, nectario ſtamineo campaniformi. *Sp. pl. p.* 441. *Iacqu. auſtr. t.* 301.

Ornithogalum firoibus ſpicatis, ſenescentibus pendulis, filamentis ſtaminum alternis majoribus, bicornibus. *Hall. hiſt. n.* 1216.

In agro vindobonenſi ad ſepes, in hortorum ambulacris. Fl. Aprili, Majo. Per.

SCILLA. Corolla hexapetala, patens decidua. Filamenta filiformia.

amœna. 1. S. floribus lateralibus alternis ſubnutantibus, ſcapo angulato. *Syſt. veget. p.* 328. *Iacqu. auſtr. t.* 218.

Hyacinthus ſtellaris cœruleus amœnus. *Bauh. pin.* 46.

In pratis agri vindobonenſis danubialibus. Fl. Aprili, Majo. Per.

bifolia. 2. S. floribus erectiusculis unilateralibus. *Retzii ſcand. p.* 77. *obs. b. faſc.* 4. *p.* 27. *Sp. pl. p.* 443. *Iacqu. auſtr. t.* 117.

Anthericum bifolium. *Scop. carn. tom.* 1. *p.* 247.

Phalangium radice bulboſa, foliis latiſſimis obtuſis, ſpica pauciflora, ſtipulis minimis. *Hall. hiſt. n.* 1211.

Hyacinthus ſtellatus vulgaris. *Clus. hiſt. p.* 184.

In Litoralis, Carnioliæ, Styriæ, Auſtriæ ſylvis, montium declivibus. Fl. Martio, Aprili. Per.

autumnalis. 3. S. foliis filiformibus linearibus, floribus corymboſis, pedunculis nudis adſcendentibus longitudine floris. *Sp. pl. p.* 443.

Anthericum autumnale. *Scop. carn. tom.* 1. *p.* 247.

Ornithogalum autumnale minus floribus cœruleis. *Tourn. inſt.* 381. *Garid. aix. t.* 73.

Hyacinthus autumnalis. *Clus. hiſt. p.* 185.

In agri fluminenſis vineis, locisque aſperis incultis. Fl. autumno. Per.

Filamenta ſunt petaliformia, plana & ut petala, linea elevata a baſi ad apicem decurrente, notata.

ASPHODELUS. Corolla ſexpartita. Nectarium conſtat valvulis ſex germen tegentibus.

ramoſus. 1. A. caule nudo, foliis enſiformibus carinatis lævibus. *Sp. pl. p.* 444. *Murr. in comm. gott.* 1776. *p.* 37. *t.* 7. *Scop. carn. tom.* 1. *p.* 246.

Aſphodelus primus. *Cluſ. hiſt. p.* 196.

Aſphodelus ſecundus. *Cluſ. hiſt. p.* 197.

In Carnioliæ montibus idrienſibus, & in monte *Nanas. Scop.* Ego in agro fluminenſi inter *Bergudi* & Præluca locis herbidis ad mare. In Croatia in *mala Planina*, alibique. Fl. Aprili, Majo, Iunio. Per.

Individua, quæ in montanis vidi, habent pedunculos ſimplices, caulem cubitalem. Iis contra, quæ ad litus maris legi, eſt caulis bicubitalis, altiorque pedunculi ſunt ramoſi.

Croatæ legionis lickenſis, hisque contermini, in annonæ caritate & inopia, tuberibus, quibus radix conſtat, coctis, veſci ſolent. At minime impune: ob tenaciſſimum enim, ventriculoque inimiciſſimum gluten, quo abundant, poſt diuturniorem uſum contrahunt viſcerum abdominalium infarctus obſtructiones, ac cum his febres, aliaque pertinaciſſima mala.

ANTHERICUM. Corolla hexapetala, patens. Capſula ovata.

ſerotinum. 1. A. foliis planiuſculis, ſcapo unifloro. *Sp. pl. p.* 444. *Iacqu. auſt. app. t.* 38.

Phalangium foliis craſſiuſculis, ſcapo unifloro. *Hall. hiſt. n.* 1209.

Bulbocodium alpinum juncifolium flore unico, intus albo, extus ſordide rubente. *Raj. ſyn.* 3. *p.* 374. *t.* 17. *f.* 2.

In herbidis, petroſisque alpium carinthiacarum, tyrolenſiumque locis. Fl. Iulio, Auguſto. Per.

ramoſum. 2. A. foliis planis, ſcapo ramoſo, corollis planis, piſtillo recto. *Sp. pl. p.* 445. *Iacqu. auſt. t.* 161. *Scop. carn. tom.* 1. *p.* 246.

Phalangium radicibus teretibus, foliis gramineis, caule ramoſo, laxe paniculato. *Hall. hiſt. n.* 1208.

In collibus, locisque montoſis ſylvaticis. Fl. Iunio, Iulio. Per.

Liliago. 3. A. foliis planis, ſcapo ſimpliciſſimo, corollis planis, piſtillo declinato. *Sp. p. p.* 445. *Iacqu. hort. tom.* 1. *p.* 36. *t.* 83. *Fl. dan. t.* 616.

Phalangium radicibus teretibus, foliis radicalibus carinatis, enſiformibus, petiolis unifloris. *Hall. hiſt. n.* 1207.

In Tyroli in rupeſtribus ſubalpinis leontinis. *Wulfen.* Fl. Iunio, Iulio. Per.

Liliaſtrum. 4. A. foliis planis, ſcapo ſimpliciſſimo, corollis campanulatis, ſtaminibus declinatis. *Sp. pl. p.* 445.

Hemerocallis floribus ſpicatis ſecundis. *Hall. hiſt. n.* 1230.

Allobrogie Hemerocallis. *Garden of Eden. t.* 57.

In Carnia in herbidis montis *Croſtis. I. Hoſt.* Fl. Iulio. Per.

calyculatum. 5. A. foliis enſiformibus, perianthiis trilobis, filamentis glabris, floribus trigynis. *Sp. pl. p.* 447. *Fl. dan. t.* 36.

Anthericum filamentis glabris. *Hall. hiſt. n.* 1205.

Anthericum Pſeudoaſphodelus. *Iacqu. en. p.* 233.

Scheuchzeria Pſeudoaſphodelus. *Scop. carn. tom.* 1. *p.* 263.

Pſeudoaſphodelus ſecundus. *Cluſ. hiſt. p.* 198.

In alpinis, & ſubalpinis. Fl. æſtate. Per.

ASPARAGUS. Corolla ſexpartita, erecta: petalis tribus interioribus apice reflexis. Bacca trilocularis, diſperma.

officinalis. 1. A. caule herbaceo tereti erecto, foliis ſetaceis, ſtipulis paribus. *Syſt. veget. p.* 332. *Fl. dan. t.* 805. *Scop. carn. tom.* 1. *p.* 248.

Aſparagus foliis filiformibus ſtipulatis. *Hall. hiſt. n.* 1239.

Aſparagus marinus. *Cluſ. hiſt. p.* CLXXIX.

L' Aſperge. Regn. bot.

In arenoſis, aſperiſque locis Litoralis. In ſylvis demiſſis Auſtriæ, Pannoniæ, Styriæ, Carnioliæ. Fl. Majo, Iunio. Per.

acutifolius. 2. A. caule inermi fruticoſo, foliis aciformibus rigidulis perennantibus mucronatis æqualibus. *Sp. pl. p.* 449.

Aſparagus Corruda. *Scop. carn. tom.* 1. *p.* 248.

Aſparagus foliis acutis. *Bauh. pin.* 490. *Zann. iſt. p.* 24. *t.* 179.

In Litorali, Forojulio ad ſepes, in dumetis, alibique. Fl. Auguſto.

Fluminenſes, conterminique Liburniæ incolæ, turiones coctos, ſale, aceto, oleo, pipereque conditos primovere comedunt, &, cum hisce vis amara, diureticaque inſignis, plerique turionibus Aſparagi officinalis ſapienter præferunt.

CONVALLARIA. Corolla ſex- vel quadrifida. Bacca tri- vel bilocularis, ante maturitatem maculoſa.

majalis. 1. C. ſcapo nudo. *Sp. pl. p.* 451. *Fl. dan. t.* 854. *Curtis lond. Scop. carn. tom.* 1. *p.* 250.

Polygonatum ſcapo diphyllo, floribus ſpicatis nutantibus, campaniformibus. *Hall. hiſt. n.* 1241.

Lilium convallium. *Camerar. epit. p.* 618.

Le Muguet. Regn. bot.

In ſylvaticis, nemorenſibus. Fl. Majo. Per.

verticillata. 2. C. foliis verticillatis. *Sp. pl. p.* 451. *Fl. dan. t.* 86. *Scop. carn. tom.* 1. *p.* 250.

Polygonatum caule ſimplici erecto, foliis ellipticis & verticillatis. *Hall. hiſt. n.* 1244.

Polygonatum quintum ſive anguſtifolium primum. *Clus. hiſt. p.* 277.

In ſubalpinis, & alpinis. Fl. a Iunio ad Auguſtum. Per.

Polygonatum. 3. C. foliis alternis amplexicaulibus, caule ancipiti, pedunculis axillaribus ſubunifloris. *Sp. pl. p.* 451. *Fl. dan. t.* 377. *Scop. carn. tom.* 1. *p.* 251.

Polygonatum caule ſimplici, anguloſo, cernuo, foliis ovato-lanceolatis, rigidis, alis unifloris. *Hall. hiſt. n.* 1242.

Le Sceau de Salamon. Regn. bot.

In ſylvis. Fl. Majo. Per.

latifolia. 4. C. foliis alternis amplexicaulibus acuminatis, caule angulato, pedunculis axillaribus, multifloris. *Syſt. veget. p.* 334. *Iacqu. auſtr. t.* 232.

Polygonatum tertium latiore folio. *Clus. hiſt. p.* 276.

In ſylvis Auſtriæ, Pannoniæ. Fl. Majo, Iunio. Per.

Habitum habet Convallariæ multifloræ; flores C. Polygonati; & folia C. majalis. Iacqu. l. c.

multiflora. 5. C. foliis alternis amplexicaulibus, caule tereti, pedunculis axillaribus multifloris. *Sp. pl. p.* 452. *Fl. dan. t.* 152. *Scop. carn. tom.* 1. *p.* 251.

Polygonatum caule ſimplici cernuo, foliis ovato-lanceolatis, petiolis multifloris. *Hall. hiſt.* n. 1243.

Polygonatum latifolium primum. *Clus. hiſt.* p. 275.

Le Muguet multiflóre. Bulliard herb. t. 307.

In ſylvis. Fl. Majo, Iunio. Per.

ia. 6. C. foliis cordatis, floribus tetrandris. *Syſt. veget.* p. 335. *Fl. dan. t.* 291. *Scop. carn. tom.* 1. *p.* 251.

Unifolium. *Hall. hiſt. n.* 1240.

Lilium convallium minus. *Bauh. pin.* 304.

In ſylvis ſubalpinis. Fl. Iunio, Iulio. Per.

Ut unicam tertiam partem in partitione corollæ ſtaminumque numero excludit, in piſtillo quoque idem obſervat; ſtigma enim eſt bifidum, bacca bilocularis. Retzii obs. fasc. 1. p. 16.

ACINTHUS. Corolla campanulata: pori tres melliferi ad apicem germinis.

ſus. 1. H. corollis angulato-cylindricis: ſummis ſterilibus longius pedicellatis. *Sp. pl. p.* 455. *Iacqu. auſtr. t.* 126. *Scop. carn. tom.* 1. *p.* 252.

Hyacinthus ſpica longiſſima, floribus ſupremis ſterilibus erectis, inferioribus fœcundis patulis. *Hall. hiſt. n.* 1247.

In arvis, collibus herbidis, pratis. Fl. Majo, Iunio. Per.

i- 2. H. corollis globoſis uniformibus, foliis canaliculato-cylindricis ſtrictis. *Sp. pl. p.* 455. *Curtis magaz.* 157.

Hyacinthus botryoides purpureus tertius. *Clus. hiſt. p.* 181.

Hyacinthus racemoſus cœruleus major. *Bauh. pin.* 42.

In vineis agri fluminenſis. Fl. Marti Aprili. Per.

racemoſus. 3. H. corollis ovatis: ſummis ſeſſilibus, foliis xis. *Sp. pl. p.* 455. *Iacqu. auſtr. t.* 187.

Hyacinthus botryoides. *Scop. carn. tom.* 1. *p.* 2

Hyacinthus botyoides primus. *Clus hiſt. p.* 1

Hyacinthus racemoſus cœruleus minor junc lius. *Bauh. pin.* 43.

In cultis, collibus, campis, & alibi. Martio, Aprili. Per.

Hyacintho botryoidi ſunt folia latiora, e ta; flores dilute cœrulei, inodori. Hyacint racemoſus contra habet folia anguſtiora, ad ram reflexa; flores ſaturate cœruleos, odora

HEMEROCALLIS. Corolla campanulata: tubo c lindrico. Stamina declinata.

flava. 1. H. foliis lineari-ſubulatis carinatis, corollæ ciniis margine planis. *Ehrh. beytr. tom.* 4 166. *Sp. pl. p.* 462. *Iacqu. hort. tom.* 2. *p. t.* 139.

Hemerocallis Lilio-Aſphodelus. *Scop. c tom.* 1. *p.* 253.

Lilio-Aſphodelus luteo flore. *Clus. hiſt. p.* 1

In uliginoſis, humidis in Iſtria prope *pichi. I. Hoſt.* In Carniolia circa *Oſeltza*, & Udalricum. *Scop.* In Pannoniæ comitatu c ferrei in pratis uliginoſis non procul ab op *Németh-Uivár* (*Güſing*) *Clus.* Fl. Iunio. Pe

ACORUS. Spadix cylindricus, tectus floſculis. lus nullus. Capſula trilocularis.

Calamus. 1. A. ſcapi mucrone longiſſimo foliaceo. *Ait.* *tom.* 1. *p.* 474. *Sp. pl. p.* 462. *Leers. p.* 87. *t.* 13. *f.* 12. *Scop. carn. tom.* 1. *p. Hall. hiſt. n.* 1307.

Le Ione odorant. Regn. bot.

In aquis ſtagnantibus, ad fluviorum ripas. Fl. Iunio. Per.

UNCUS. Calyx hexaphyllus. Corolla nulla. Capſula unilocularis.

* *Culmis nudis.*

onglomeratus. 1. I. culmo nudo ſtricto, capitulo laterali. *Sp. pl. p.* 464. *Leers herb. p.* 87. *t.* 13. *f.* 1.

Iuncus caule nudo, foliis teretibus mollibus, panicula conglomerata. *Hall. hiſt. n.* 1312.

Iuncus lævis, panicula conglomerata. *Scheuch. agr.* 343.

In uliginoſis, paluſtribus. Fl. Iunio. Per.

effuſus. 2. I. culmo nudo ſtricto: panicula laterali. *Sp. pl. p.* 464. *Leers herb. p.* 88. *t.* 13. *f.* 2.

Iuncus lævis, panicula ſparſâ major. *Bauh. pin.* 12. *Scheuch. agr.* 341.

In foſſis, aliisque udis locis. Fl. Iunio, Iulio. Per.

Huic, & præcedenti ſunt flores triandri, qua nota a junco inflexo, cui flores hexandri, ſufficienter diſtinguuntur. Ehrhart.

inflexus. 3. I. culmo nudo: apice membranaceo incurvo, panicula laterali. *Sp. pl. p.* 464. *Leers herb. p.* 88. *t.* 13. *f.* 3.

Iuncus acumine reflexo alter. *Bauh. pin.* 12. *Scheuch. agr.* 345.

Habitat & floret cum præcedente. Per.

filiformis. 4. I. culmo nudo filiformi nutante, panicula laterali. *Sp. pl. p.* 465.

Iuncus caule nudo, foliis teretibus mollibus, panicula laterali pauciflora. *Hall. hiſt. n.* 1313.

Iuncus lævis, panicula ſparſa minor. *Bauh pin.* 12. *Scheuch. agr.* 347. *t.* 7. *f.* 11.

In Styria in turfoſis ad lacus alpium judenburgenſium. Fl. Iulio, Auguſto. Per.

trifidus. 5. I. culmo nudo, foliis floribusque tribus terminalibus. *Sp. pl. p.* 465. *Fl. dan. t.* 107.

Iuncus monanthos. *Iacqu. en. p.* 236.

Iuncus acumine reflexo minor, & trifidus. *Bauh. pin.* 12. *Hall. hiſt. n.* 1315.

Iuncoides alpinum trifidum. *Scheuch. agr.* 325.

In alpibus. Fl. Iunio, Iulio. Per.

Variat culmo uni- bi- & trifloro.

** *Culmis folioſis.*

articulatus. 6. I. foliis nodoſo-articulatis, petalis obtuſis. *Sp. pl. p.* 465. *Leers herb. p.* 89. *t.* 13. *f.* 6.

In aquoſis. Fl. æſtate. Per.

bulboſus. 7. I. foliis linearibus canaliculatis, capſulis obtuſis. *Sp. pl. p.* 466. *Leers herb. p.* 90. *t.* 13. *f.* 7.

Iuncus foliis carinatis, panicula ſparſa, fructu globoſo. *Hall. hiſt. n.* 1318.

Iuncus compreſſus. *Iacqu. en. p.* 235.

Gramen junceum, foliis & ſpica junci. *Bauh. pin.* 5. *Scheuch. agr.* 322.

In udis. Fl. æſtate. Per.

bufonius. 8. I. culmo dichotomo, foliis angulatis, floribus ſolitariis ſeſſilibus. *Sp. pl. p.* 466. *Leers herb. p.* 90. *t.* 13. *f.* 8.

Iuncus caule brachiato, foliis ſetaceis, floribus ſolitariis, ad ramum ſeſſilibus. *Hall. hiſt. n.* 1319.

Gramen nemoroſum, calyculis paleaceis, repens. *Bauh. pin.* 7. *Scheuch. agr.* 229.

In udis. Fl. æſtate.

Iacquini. 9. I. folio ſubulato, capitulo ſubquadrifloro terminali. *Syſt. veget. p.* 341 *Iacqu. auſtr. t.* 221.

Iuncus caule unifolio, petiolis bifloris, floribus umbellatis. *Hall. hiſt. n.* 1316.

Iuncoides alpinum, floſculis junci glomeratis, atro-fuſcis. *Scheuch. agr.* 323.

In alpibus. Fl. Iunio, Iulio. Per.

Radix horizontalis, nodoſa. Folium radicale unicum, ſolitarium; in culmo etiam unicum. Varius in variis individuis florum numerus. Altitudo culmorum, & foliorum radicalium in diverſis alpibus varia. In alpibus auſtriacis culmos pollicares, & bipollicares; in aliis alpibus pedales, & pede longiores legi.

triglumis. 10. I. foliis planis, gluma triflora terminali. *Sp. pl. p.* 467. *Fl. dan. t.* 132.

Iuncus gluma triflora culmum terminante. *Fl. lapp. n.* 115. *t.* 10. *f.* 5.

Iuncus caule bifolio, foliis teretibus, floribus terminantibus ternis. *Hall. hiſt. n.* 1314.

Iuncus exiguus montanus mucrone carens. *Bauh. pin.* 12. *prodr.* 22.

In udis alpium ſtyriacarum, carinthiacarum, tyrolenſium. Fl. Iulio. Per.

piloſus. 11. I. foliis planis piloſis, corymbo ramoſo. *Sp. pl. p.* 468. *Leers herb. p.* 91. *t.* 13. *f.* 10.

Iuncus foliis gramineis hirſutis, floribus umbellatis, ſolitariis, petiolatis, ariſtatis. *Hall. hiſt. n.* 1325.

Gramen hirſutum, latifolium, minus. *Bauh. pin.* 7. *Scheuch. agr.* 315.

In montanis ſylvaticis. Fl. Majo, Iunio. Per.

ſylvaticus. 12. I. foliis planis piloſis, corymbo decompoſito, floribus faſciculatis ſeſſilibus. *Huds. angl. ed.* 2. 157. *Sibthorp. oxon. p.* 116. *Curtis lond.*

Iuncus *maximus*, foliis planis piloſis, paniculæ ramoſæ floribus ſubternis ſubſeſſilibus. *Retzii ſcand. p.* 82. *Ehrh. beytr. tom.* 6. *p.* 139.

Iuncus foliis gramineis hirſutis, floribus paniculatis, faſciculatis. *Hall. hiſt. n.* 1324.

Iuncus latifolius. *Wulfen in Iacqu. collect. tom.* 3. *p.* 59.

Gramen nemoroſum, hirſutum, latifolium majus. *Bauh. pin.* 7. *Scheuch. agr.* 317.

In ſylvis montoſis, ſubalpinis, & in ipſis alpibus. Fl. a Majo ad Iulium. Per.

niveus, 13. I. foliis planis ſubpiloſis, corymbis folio brevioribus, floribus faſciculatis. *Sp. pl. p.* 468. *Leers herb. p.* 91. *t.* 13. *f.* 9.

Iuncus foliis gramineis, paniculis umbellatis, petalis interioribus duplo longioribus. *Hall. hiſt. n.* 1328.

Iuncoides montanum, nemoroſum, flore niveo, *Scheuch. agr.* 320.

In montoſis ſylvaticis Styriæ, Carnioliæ. Fl. Iunio, Iulio. Per.

campeſtris. 14. I. foliis planis ſubpiloſis, ſpicis ſeſſilibus, pedunculatisque. *Sp. pl. p.* 468. *Curtis lond. Leers herb. p.* 92. *t.* 13. *f.* 5.

Iuncus foliis gramineis hirſutis, ſpicis petiolatis, nutantibus, petalis ariſtatis. *Hall. hiſt. n.* 1330.

Iuncoides villoſum capitulis Pſyllii. *Scheuch. agr.* 310.

In ſylvis, & pratis ſiccioribus. Fl. Martio, Aprili. Per.

BERBERIS. Calyx hexaphyllus. Petala ſex. Glandulæ duæ baſi ſinguli petali adnatæ. Stylus nullus. Bacca apice perforata, diſperma.

vulgaris. 1. B. pedunculis racemoſis. *Sp. pl. p.* 471. *Fl. dan. t.* 904. *Scop. carn. tom.* 1. *p.* 259.

Berberis floribus racemoſis, foliis ciliatis. *Hall. hiſt. n.* 828.

L' Epine Vinette. Regn. bot.

In ſylvis, dumetis, ad ſepes. Fl. Majo, Iunio. Fr.

LORANTHUS. Germen inferum. Calyx nullus. Corolla ſexfida, revoluta. Stamina adnata baſibus petalorum, longitudine corollæ. Bacca monoſperma.

europæus. 1. L. racemis omnibus terminalibus ſimplicibus, floribus dioicis. *Iacqu. en. p.* 230. *Fl. auſtr. t.* 30. *Sp. pl. app. p.* 1672.

Ramis quercuum innaſcitur in ſylvis Auſtriæ, Pannoniæ, Iſtriæ. Fl. Aprili, Majo. Fr.

Flores in aliis individuis fertiles, in aliis ſteriles. Sterilium perianthium eſt margo parvus, obſolete ſexdentatus, coronans germen; quod etiam ſquamulæ inſidet propriæ. Stamina ſex, raro pauciora aut plura. Antheræ ſubgeminæ. Germen turbinatum. Stylus ſine ſtigmatæ. Fertilibus calyx, corolla ut ſterilibus. Filamenta antheris deſtituta. Piſtillum habet ſtigma obtuſum. Iacqu. l. c.

PEPLIS. Perianthium campanulatum: ore decemfido. Petala ſex, calyci inſerta. Capſula bilocularis.

Portula. 1. P. floribus hexandris, foliis cuneiformibus obtuſis. *Wild. ber. p.* 128. *Sp. pl. p.* 474. *Fl. dan. t.* 64. *Curtis lond.*

Peplis caule repente, floribus axillaribus, petalis minutiſſimis fugacibus. *Sibthorp. oxon. p.* 108.

In inundatis. Fl. Iulio, Auguſto.

TRIGYNIA.

RUMEX. Calyx triphyllus. Petala tria, conniventia. Semen unicum, triquetrum.

* *Hermaphroditi: valvulis granulo notatis.*

criſpus. 1. R. floribus hermaphroditis: valvulis integris graniferis, foliis lanceolatis undulatis acutis. *Sp. pl. p.* 476. *Curtis lond.*

Lapathum foliis criſpis, imis ovatis, ſupremis lanceolatis, calycibus verrucoſis. *Hall. hiſt. n.* 1589.

Lapathum criſpum. *Scop. carn. tom.* 1. *p.* 261.

Ad vias, foſſas, & alibi. Fl. Iunio, Iulio. Per.

Hydrolapathum. 2. R. floribus hermaphroditis: valvulis integris graniferis, foliis lanceolatis acutis. *Huds. angl. ed.* 2. 154. *Sibthorp. oxon. p.* 118.

Rumex Nemolapathum. *Linn. ſuppl. p.* 212.

Lapathum maximum aquaticum, ſeu Hydrolapathum. *Raj. ſyn.* 140.

In Auſtriæ, Pannoniæ arundinetis, foſſis, ad fluviorum, lacuumque ripas. Fl. Iunio, Iulio. Per.

maritimus. 3. R. floribus hermaphroditis: valvulis dentatis graniferis, foliis linearibus. *Sp. pl. p.* 478. *Curtis lond.*

Lapathum aquaticum luteolæ folio. *Tourn. inſt.* 504.

In Auſtriæ paluſtribus prope *Himberg, Bruck an der Leitha, Rohrau*, alibique. Fl. Iunio, Iulio.

acutus. 4. R. floribus hermaphroditis: valvulis dentatis graniferis, foliis cordato-oblongis acuminatis. *Sp. pl. p.* 478. *Curtis lond.*

Lapathum petiolo lateſcente, foliis longe lanceolatis, calyce ſerrato. *Hall. hiſt. n.* 1591.

Lapathum acutum. *Scop. carn. tom.* 1. *p.* 262.

Ad foſſas, fluvios, vias, in pratis udis, alibique. Fl. Iunio. Per.

obtuſifolius. 5. R. floribus hermaphroditis: valvulis dentatis graniferis, foliis cordato-oblongis obtuſiuſculis crenatis. *Sp. pl. p.* 478. *Curtis lond.*

Lapathum foliis ovatis, circa petiolum emarginatis, floribus denſe paniculatis, ſerratis, verrucoſis. *Hall. hiſt. n.* 1592.

In pomariis, pratis, ad vias. Fl. Iunio, Iulio. Per.

Folia radicalia cordato-ovata, petiolata, obtuſiuſcula, ſubtus venoſa. Caulina acuta, ſubundulata. Curtis l. c.

pulcher. 6. R. floribus hermaphroditis: valvulis dentatis: ſubunica granifera, foliis radicalibus panduriformibus. *Sp. pl. p.* 477.

Lapathum foliis radicalibus utrinque emarginatis, calycibus reticulatis, ciliatis, verrucoſis. *Hall. hiſt. n.* 1593.

Lapathum pulchrum bononienſe ſinuatum. *Bauh. hiſt.* 2. *p.* 988.

Lapathum ramis procumbentibus, ſeminis involucro dentato, foliis inferioribus inſtar fidium. *Moris hiſt.* 2. *p.* 580. *ſect.* 5. *t.* 27. *f.* 13.

In agro vindobonenſi ad margines agrorum, ad vias, in ruderatis. Fl. Iunio, Iulio. Per.

*** Hermaphroditi : valvulis granulo deſtitutis.*

ſcutatus. 7. R. floribus hermaphroditis, foliis cordato-haſtatis, caule tereti. *Syſt. veget. p.* 347.

Lapathum foliis ſagittatis, cis mucronem latiſſimis, hamis reflexis divergentibus. *Hall. hiſt. n.* 1594.

Acetoſa ſcutata repens. *Bauh. pin.* 114. *prodr.* 55. *Triumphett. obs. p.* 26. *ic.* 68.

In Litorali inter Veloscam & Caſtuam in maceriis, aliisque difformibus lapidum acervis infrequentem vidi. Fl. Iunio. Per.

digynus. 8. R. floribus hermaphroditis digynis. *Sp. pl. p.* 480. *Fl. dan. t.* 14.

Lapathum foliis antice emarginatis. *Hall. hiſt. n.* 1595.

Locis ſaxoſis ſubalpinis, & alpinis. Fl. Iulio, Auguſto. Per.

**** Floribus diclinis.*

alpinus. 9. R. floribus hermaphroditis ſterilibus femineisque, valvulis integerrimis nudis, foliis cordatis obtuſis rugoſis. *Sp. pl. p.* 480.

Lapathum foliis obtuſis, circa petiolum emarginatis, paniculis denſe ſpicatis, vaginis maximis. *Hall. hiſt. n.* 1587.

Rumex latifolius vulgo Rha æſtimatus. *Clus. hiſt. p. LXIX.*

In alpinis, & ſubalpinis, præprimis ad paſtorum caſas. Fl. æſtate. Per.

Acetoſa. 10. R. floribus dioicis, foliis oblongis ſagittatis. *Sp. pl. p.* 481.

Lapathum ſexubus ſeparatis, foliis ſagittatis, hamis retrorſum porrectis. *Hall. hiſt. n.* 1597.

Lapathum Acetoſa. *Scop. carn. tom.* 1. *p.* 260.

Acetoſa pratenſis. *Bauh. pin.* 114. *Zann. iſt. p.* 4. *t.* 233.

In pratis, paſcuis demiſſis, & alpinis. Fl. a Majo ad Iulium. Per.

Aceto-ſella. 11. R. floribus dioicis, foliis lanceolato-haſtatis. *Sp. pl. p.* 481. *Curtis lond.*

Lapathum ſexubus ſeparatis, foliis ſagittatis, hamis acutis, recurvis. *Hall. hiſt. n.* 1596.

Lapathum Acetoſella. *Scop. carn. tom.* 1. *p.* 261.

In ruderatis, collibus, montoſis ſylvaticis, & alibi. Fl. Majo, Iunio. Per.

TRIGLOCHIN. Calyx triphyllus. Petala tria, calyciformia. Stylus nullus. Stigmata tria vel ſex, plumoſa. Capſula tri- vel ſexlocularis, baſi dehiscens. Semina ſolitaria.

paluſtre. 1. T. floribus trigynis, capſulis ſublinearibus, trigonis, trilocularibus. *Roth. germ. tom.* 1. *p.* 158. *Sp. pl. p.* 482. *Fl. dan. t.* 490. *Scop. carn. tom.* 1. *p.* 265. *Leers herb. p.* 93. *t.* 12. *f.* 5.

In udis, paluſtribus. Fl. Iulio. Per.

maritimum. 2. T. floribus hexagynis, capſulis ovatis, hexagonis, ſexlocularibus. *Roth germ. tom.* 1. *p.* 158. *Sp. pl. p.* 483. *Fl. dan. t.* 306. *Scop. carn. tom.* 1. *p.* 265.

Iuncago paluſtris, & maritima, perennis, fructu breviore, quinquecapſulari. *Mich. gen. p.* 44. *Zann. iſt. p.* 149. *t.* 273. *n.* 2.

In paluſtribus Auſtriæ prrope *Himberg.* In Pannonia ad lacum Peiſonis. In Litorali in paludibus duinenſibus, aquilejenſibus. Fl. Iunio, Iulio. Per.

In vineis agri fluminenſis. Fl. Martio, Aprili. Per.

racemoſus. 3. H. corollis ovatis: ſummis ſeſſilibus, foliis laxis. *Sp. pl. p.* 455. *Iacqu. auſtr. t.* 187.

Hyacinthus botryoides. *Scop. carn. tom.* 1. *p.* 252.

Hyacinthus botyoides primus. *Clus hiſt. p.* 181.

Hyacinthus racemoſus cœruleus minor juncifolius. *Bauh. pin.* 43.

In cultis, collibus, campis, & alibi. Fl. Martio, Aprili. Per.

Hyacintho botryoidi ſunt folia latiora, erecta; flores dilute cœrulei, inodori. Hyacinthus racemoſus contra habet folia anguſtiora, ad terram reflexa; flores ſaturate cœruleos, odoros.

HEMEROCALLIS. Corolla campanulata: tubo cylindrico. Stamina declinata.

flava. 1. H. foliis lineari-ſubulatis carinatis, corollæ laciniis margine planis. *Ehrh. beytr. tom.* 4. *p.* 166. *Sp. pl. p.* 462. *Iacqu. hort. tom.* 2. *p.* 6[illegible] *t.* 139.

Hemerocallis Lilio-Aſphodelus. *Scop. carn. tom.* 1. *p.* 253.

Lilio-Aſphodelus luteo flore. *Clus. hiſt. p.* 13[illegible]

In uliginoſis, humidis in Iſtria prope C[illegible]pichi. *I. Hoſt.* In Carniolia circa *Oſeltza*, & [illegible] Udalricum. *Scop.* In Pannoniæ comitatu caſ[illegible] ferrei in pratis uliginoſis non procul ab oppi[illegible] *Németh-Uivár* (*Güſing*) *Clus.* Fl. Iunio. Per.

ACORUS. Spadix cylindricus, tectus floſculis. Stylus nullus. Capſula trilocularis.

Calamus. 1. A. ſcapi mucrone longiſſimo foliaceo. *Ait.* [illegible] *tom.* 1. *p.* 474. *Sp. pl. p.* 462. *Leers.* [illegible] *p.* 87. *t.* 13. *f.* 12. *Scop. carn. tom.* 1. *p.* [illegible] *Hall. hiſt. n.* 1307.

Le Ione odorant. Regn. bot.

In aquis ſtagnantibus, ad fluviorum ripas. Fl. Iunio. Per.

IUNCUS. Calyx hexaphyllus. Corolla nulla. Capſula unilocularis.

* *Culmis nudis.*

conglomeratus. 1. I. culmo nudo ſtricto, capitulo laterali. *Sp. pl. p.* 464. *Leers herb. p.* 87. *t.* 13. *f.* 1.

Iuncus caule nudo, foliis teretibus mollibus, panicula conglomerata. *Hall. hiſt. n.* 1312.

Iuncus lævis, panicula conglomerata. *Scheuch. agr.* 343.

In uliginoſis, paluſtribus. Fl. Iunio. Per.

effuſus. 2. I. culmo nudo ſtricto: panicula laterali. *Sp. pl. p.* 464. *Leers herb. p.* 88. *t.* 13. *f.* 2.

Iuncus lævis, panicula ſparſâ major. *Bauh. pin.* 12. *Scheuch. agr.* 341.

In foſſis, aliisque udis locis. Fl. Iunio, Iulio. Per.

Huic, & præcedenti ſunt flores triandri, qua nota a junco inflexo, cui flores hexandri, ſufficienter diſtinguuntur. Ehrhart.

inflexus. 3. I. culmo nudo: apice membranaceo incurvo, panicula laterali. *Sp. pl. p.* 464. *Leers herb. p.* 88. *t.* 13. *f.* 3.

Iuncus acumine reflexo alter. *Bauh. pin.* 12. *Scheuch. agr.* 345.

Habitat & floret cum præcedente. Per.

filiformis. 4. I. culmo nudo filiformi nutante, panicula laterali. *Sp. pl. p.* 465.

Iuncus caule nudo, foliis teretibus mollibus, panicula laterali pauciflora. *Hall. hiſt. n.* 1313.

Iuncus lævis, panicula ſparſa minor. *Bauh. pin.* 12. *Scheuch. agr.* 347. *t.* 7. *f.* 11.

In Styria in turfoſis ad lacus alpium judenburgenſium. Fl. Iulio, Auguſto. Per.

trifidus. 5. I. culmo nudo, foliis floribusque tribus terminalibus. *Sp. pl. p.* 465. *Fl. dan. t.* 107.

Iuncus monanthos. *Iacqu. en. p.* 236.

Iuncus acumine reflexo minor, & trifidus. *Bauh. pin.* 12. *Hall. hiſt. n.* 1315.

Iuncoides alpinum trifidum. *Scheuch. agr.* 325.

In alpibus. Fl. Iunio, Iulio. Per.

Variat culmo uni- bi- & trifloro.

** *Culmis folioſis.*

articulatus. 6. I. foliis nodoſo-articulatis, petalis obtuſis. *Sp. pl. p.* 465. *Leers herb. p.* 89. *t.* 13. *f.* 6.

In aquoſis. Fl. æſtate. Per.

bulboſus. 7. I. foliis linearibus canaliculatis, capſulis obtuſis. *Sp. pl. p.* 466. *Leers herb. p.* 90. *t.* 13. *f.* 7.

Iuncus foliis carinatis, panicula ſparſa, fructu globoſo. *Hall. hiſt. n.* 1318.

Iuncus compreſſus. *Iacqu. en. p.* 235.

Gramen junceum, foliis & ſpica junci. *Bauh. pin.* 5. *Scheuch. agr.* 322.

In udis. Fl. æſtate. Per.

bufonius. 8. I. culmo dichotomo, foliis angulatis, floribus ſolitariis ſeſſilibus. *Sp. pl. p.* 466. *Leers herb. p.* 90. *t.* 13. *f.* 8.

Iuncus caule brachiato, foliis ſetaceis, floribus ſolitariis, ad ramum ſeſſilibus. *Hall. hiſt. n.* 1319.

Gramen nemoroſum, calyculis paleaceis, repens. *Bauh. pin.* 7. *Scheuch. agr.* 229.

In udis. Fl. æſtate.

Iacquini. 9. I. folio ſubulato, capitulo ſubquadrifloro terminali. *Syſt. veget. p.* 341 *Iacqu. auſtr. t.* 221.

Iuncus caule unifolio, petiolis bifloris, floribus umbellatis. *Hall. hiſt. n.* 1316.

Iuncoides alpinum, flosculis junci glomeratis, atro-fuscis. *Scheuch. agr.* 323.

In alpibus. Fl. Iunio, Iulio. Per.

Radix horizontalis, nodoſa. Folium radicale unicum, ſolitarium; in culmo etiam unicum. Varius in variis individuis florum numerus. Altitudo culmorum, & foliorum radicalium in diverſis alpibus varia. In alpibus auſtriacis culmos pollicares, & bipollicares; in aliis alpibus pedales, & pede longiores legi.

triglumis. 10. I. foliis planis, gluma triflora terminali. *Sp. pl. p.* 467. *Fl. dan. t.* 132.

Iuncus gluma triflora culmum terminante. *Fl. lapp. n.* 115. *t.* 10. *f.* 5.

Iuncus caule bifolio, foliis teretibus, floribus terminantibus ternis. *Hall. hiſt. n.* 1314.

Iuncus exiguus montanus mucrone carens. *Bauh. pin.* 12. *prodr.* 22.

In udis alpium ſtyriacarum, carinthiacarum, tyrolenſium. Fl. Iulio. Per.

piloſus. 11. I. foliis planis piloſis, corymbo ramoſo. *Sp. pl. p.* 468. *Leers herb. p.* 91. *t.* 13. *f.* 10.

Iuncus foliis gramineis hirſutis, floribus umbellatis, ſolitariis, petiolatis, ariſtatis. *Hall. hiſt. n.* 1325.

Gramen hirſutum, latifolium, minus. *Bauh. pin.* 7. *Scheuch. agr.* 315.

In montanis ſylvaticis. Fl. Majo, Iunio. Per.

ſylvaticus. 12. I. foliis planis piloſis, corymbo decompoſito, floribus fasciculatis ſeſſilibus. *Huds. angl. ed.* 2. 157. *Sibthorp. oxon. p.* 116. *Curtis lond.*

Iuncus *maximus*, foliis planis piloſis, paniculæ ramoſæ floribus ſubternis ſubſeſſilibus. *Retzii ſcand. p.* 82. *Ehrh. beytr. tom.* 6. *p.* 139.

Iuncus foliis gramineis hirſutis, floribus paniculatis, faſciculatis. *Hall. hiſt. n.* 1324.

Iuncus latifolius. *Wulfen in Iacqu. collect. tom.* 3. *p.* 59.

Gramen nemoroſum, hirſutum, latifolium majus. *Bauh. pin.* 7. *Scheuch. agr.* 317.

In ſylvis montoſis, ſubalpinis, & in ipſis alpibus. Fl. a Majo ad Iulium. Per.

niveus. 13. I. foliis planis ſubpiloſis, corymbis folio brevioribus, floribus faſciculatis. *Sp. pl. p.* 468. *Leers herb. p.* 91. *t.* 13. *f.* 9.

Iuncus foliis gramineis, paniculis umbellatis, petalis interioribus duplo longioribus. *Hall. hiſt. n.* 1328.

Iuncoides montanum, nemoroſum, flore niveo, *Scheuch. agr.* 320.

In montoſis ſylvaticis Styriæ, Carnioliæ. Fl. Iunio, Iulio. Per.

campeſtris. 14. I. foliis planis ſubpiloſis, ſpicis ſeſſilibus, pedunculatisque. *Sp. pl. p.* 468. *Curtis lond. Leers herb. p.* 92. *t.* 13. *f.* 5.

Iuncus foliis gramineis hirſutis, ſpicis petiolatis, nutantibus, petalis ariſtatis. *Hall. hiſt. n.* 1330.

Iuncoides villoſum capitulis Pſyllii. *Scheuch. agr.* 310.

In ſylvis, & pratis ſiccioribus. Fl. Martio, Aprili. Per.

BERBERIS. Calyx hexaphyllus. Petala ſex. Glandulæ duæ baſi ſinguli petali adnatæ. Stylus nullus. Bacca apice perforata, diſperma.

vulgaris. 1. B. pedunculis racemoſis. *Sp. pl. p.* 471. *Fl. dan. t.* 904. *Scop. carn. tom.* 1. *p.* 259.

Berberis floribus racemoſis, foliis ciliatis. *Hall. hiſt. n.* 828.

L' Epine Vinette. Regn. bot.

In ſylvis, dumetis, ad ſepes. Fl. Majo, Iunio. Fr.

LORANTHUS. Germen inferum. Calyx nullus. Corolla ſexfida, revoluta. Stamina adnata baſibus petalorum, longitudine corollæ. Bacca monoſperma.

europæus. 1. L. racemis omnibus terminalibus ſimplicibus, floribus dioicis. *Iacqu. en. p.* 230. *Fl. auſtr. t.* 30. *Sp. pl. app. p.* 1672.

Ramis quercuum innaſcitur in ſylvis Auſtriæ, Pannoniæ, Iſtriæ. Fl. Aprili, Majo. Fr.

Flores in aliis individuis fertiles, in aliis ſteriles. Sterilium perianthium eſt margo parvus, obſolete ſexdentatus, coronans germen; quod etiam ſquamulæ inſidet propriæ. Stamina ſex, raro pauciora aut plura. Antheræ ſubgeminæ. Germen turbinatum. Stylus ſine ſtigmatæ. Fertilibus calyx, corolla ut ſterilibus. Filamenta antheris deſtituta. Piſtillum habet ſtigma obtuſum. Iacqu. l. c.

PEPLIS. Perianthium campanulatum: ore decemfido. Petala ſex, calyci inſerta. Capſula bilocularis.

Portula. 1. P. floribus hexandris, foliis cuneiformibus obtuſis. *Wild. ber. p.* 128. *Sp. pl. p.* 474. *Fl. dan. t.* 64. *Curtis lond.*

Peplis caule repente, floribus axillaribus, petalis minutiſſimis fugacibus. *Sibthorp. oxon. p.* 108.

In inundatis. Fl. Iulio, Auguſto.

TRIGYNIA.

RUMEX. Calyx triphyllus. Petala tria, conniventia. Semen unicum, triquetrum.

* *Hermaphroditi: valvulis granulo notatis.*

criſpus. 1. R. floribus hermaphroditis: valvulis integris graniferis, foliis lanceolatis undulatis acutis. *Sp. pl. p.* 476. *Curtis lond.*

Lapathum foliis criſpis, imis ovatis, ſupremis lanceolatis, calycibus verrucoſis. *Hall. hiſt. n.* 1589.

Lapathum criſpum. *Scop. carn. tom.* 1. *p.* 261.

Ad vias, foſſas, & alibi. Fl. Iunio, Iulio. Per.

Hydrolapathum. 2. R. floribus hermaphroditis: valvulis integris graniferis, foliis lanceolatis acutis. *Huds. angl. ed.* 2. 154. *Sibthorp. oxon. p.* 118.

Rumex Nemolapathum. *Linn. ſuppl. p.* 212.

Lapathum maximum aquaticum, ſeu Hydrolapathum. *Raj. ſyn.* 140.

In Auſtriæ, Pannoniæ arundinetis, foſſis, ad fluviorum, lacuumque ripas. Fl. Iunio, Iulio. Per.

maritimus. 3. R. floribus hermaphroditis: valvulis dentatis graniferis, foliis linearibus. *Sp. pl. p.* 478. *Curtis lond.*

Lapathum aquaticum luteolæ folio. *Tourn. inſt.* 504.

In Auſtriæ paluſtribus prope *Himberg, Bruck an der Leitha*, *Rohrau*, alibique. Fl. Iunio, Iulio.

acutus. 4. R. floribus hermaphroditis: valvulis dentatis graniferis, foliis cordato-oblongis acuminatis. *Sp. pl. p.* 478. *Curtis lond.*

Lapathum petiolo lateſcente, foliis longe lanceolatis, calyce ſerrato. *Hall. hiſt. n.* 1591.

Lapathum acutum. *Scop. carn. tom.* 1, *p.* 262.

Ad foſſas, fluvios, vias, in pratis udis, alibique. Fl. Iunio. Per.

obtuſifolius. 5. R. floribus hermaphroditis: valvulis dentatis graniferis, foliis cordato-oblongis obtuſiuſculis crenatis. *Sp. pl. p.* 478. *Curtis lond.*

Lapathum foliis ovatis, circa petiolum emarginatis, floribus denſe paniculatis, ſerratis, verrucoſis. *Hall. hiſt. n.* 1592.

In pomariis, pratis, ad vias. Fl. Iunio, Iulio. Per.

Folia radicalia cordato-ovata, petiolata, obtuſiuſcula, ſubtus venoſa. Caulina acuta, ſubundulata. Curtis l. c.

pulcher. 6. R. floribus hermaphroditis: valvulis dentatis: ſubunica granifera, foliis radicalibus panduriformibus. *Sp. pl. p.* 477.

Lapathum foliis radicalibus utrinque emarginatis, calycibus reticulatis, ciliatis, verrucoſis. *Hall. hiſt. n.* 1593.

Lapathum pulchrum bononienſe ſinuatum. *Bauh. hiſt.* 2. *p.* 988.

Lapathum ramis procumbentibus, ſeminis involucro dentato, foliis inferioribus inſtar fidium. *Moris hiſt.* 2. *p.* 580. *ſect.* 5. *t.* 27. *f.* 13.

In agro vindobonenſi ad margines agrorum, ad vias, in ruderatis. Fl. Iunio, Iulio. Per.

*** Hermaphroditi: valvulis granulo deſtitutis.*

ſcutatus. 7. R. floribus hermaphroditis, foliis cordato-haſtatis, caule tereti. *Syſt. veget. p.* 347.

Lapathum foliis ſagittatis, cis mucronem latiſſimis, hamis reflexis divergentibus. *Hall. hiſt. n.* 1594.

Acetoſa ſcutata repens. *Bauh. pin.* 114. *prodr.* 55. *Triumphett. obs. p.* 26. *ic.* 68.

In Litorali inter Veloscam & Caſtuam in maceriis, aliisque difformibus lapidum acervis infrequentem vidi. Fl. Iunio. Per.

digynus. 8. R. floribus hermaphroditis digynis. *Sp. pl. p.* 480. *Fl. dan. t.* 14.

Lapathum foliis antice emarginatis. *Hall. hiſt. n.* 1595.

Locis ſaxoſis ſubalpinis, & alpinis. Fl. Iulio, Auguſto. Per.

**** Floribus diclinis.*

alpinus. 9. R. floribus hermaphroditis ſterilibus femineisque, valvulis integerrimis nudis, foliis cordatis obtuſis rugoſis. *Sp. pl. p.* 480.

Lapathum foliis obtuſis, circa petiolum emarginatis, paniculis denſe ſpicatis, vaginis maximis. *Hall. hiſt. n.* 1587.

Rumex latifolius vulgo Rha æſtimatus. *Clus. hiſt. p. LXIX.*

In alpinis, & ſubalpinis, præprimis ad paſtorum caſas. Fl. æſtate. Per.

Acetoſa. 10. R. floribus dioicis, foliis oblongis ſagittatis. *Sp. pl. p.* 481.

Lapathum ſexubus ſeparatis, foliis ſagittatis, hamis retrorſum porrectis. *Hall. hiſt. n.* 1597.

Lapathum Acetoſa. *Scop. carn. tom.* 1. *p.* 260.

Acetoſa pratenſis. *Bauh. pin.* 114. *Zann. iſt. p.* 4. *t.* 233.

In pratis, paſcuis demiſſis, & alpinis. Fl. a Majo ad Iulium. Per.

Aceto-ſella. 11. R. floribus dioicis, foliis lanceolato-haſtatis. *Sp. pl. p.* 481. *Curtis lond.*

Lapathum ſexubus ſeparatis, foliis ſagittatis, hamis acutis, recurvis. *Hall. hiſt. n.* 1596.

Lapathum Acetoſella. *Scop. carn. tom.* 1. *p.* 261.

In ruderatis, collibus, montoſis ſylvaticis, & alibi. Fl. Majo, Iunio. Per.

TRIGLOCHIN. Calyx triphyllus. Petala tria, calyciformia. Stylus nullus. Stigmata tria vel ſex, plumoſa. Capſula tri- vel ſexlocularis, baſi dehiscens. Semina ſolitaria.

paluſtre. 1. T. floribus trigynis, capſulis ſublinearibus, trigonis, trilocularibus. *Roth. germ. tom.* 1. *p.* 158. *Sp. pl. p.* 482. *Fl. dan. t.* 490. *Scop. carn. tom.* 1. *p.* 265. *Leers herb. p.* 93. *t.* 12. *f.* 5.

In udis, paluſtribus. Fl. Iulio. Per.

maritimum. 2. T. floribus hexagynis, capſulis ovatis, hexagonis, ſexlocularibus. *Roth germ. tom.* 1. *p.* 158. *Sp. pl. p.* 483. *Fl. dan. t.* 306. *Scop. carn. tom.* 1. *p.* 265.

Iuncago paluſtris, & maritima, perennis, fructu breviore, quinquecapſulari. *Mich. gen. p.* 44. *Zann. iſt. p.* 149. *t.* 273. *n.* 2.

In paluſtribus Auſtriæ prrope *Himberg.* In Pannonia ad lacum Peiſonis. In Litorali in paludibus duinenſibus, aquilejenſibus. Fl. Iunio, Iulio. Per.

COLCHICUM. Calycis loco ſpatha. Corolla ſexpartita: tubo radicato. Germen intra radicem ſepultum. Capſulæ tres, connæxæ, inflatæ.

autumnale. 1. C. foliis planis lanceolatis erectis. *Sp. pl. p.* 485. *Scop. carn. tom.* 1. *p.* 265.

Colchicum flore folia longe præcedente, petalis ovatis. *Hall. hiſt. n.* 1255.

Colchicum pannonicum. *Cluſ. hiſt. p.* 199.

Le Colchique, ou Tue-chien. Regn. bot. Bulliard herb. t. 18.

In pratis. Fl. ab Auguſto ad ſerum autumnum. Per.

Fructus & foliorum rudimenta hyemem ſub terra tranſigunt, ſequenti vere e terra proditura.

POLYGYNIA.

ALISMA. Calyx triphyllus. Petala tria. Semina plura.

Plantago. 1. A. foliis ovatis acutis, fructibus obtuſe trigonis. *Sp. pl. p.* 486. *Curtis lond. Fl. dan. t.* 561. *Scop. carn. tom.* 1. *p.* 266.

Damaſonium foliis ellipticis, lanceolatis, capitulo rotunde triquetro. *Hall. hiſt. n.* 1184.

In aquoſis, paluſtribus. Fl. æſtate. Per.

HEPTANDRIA.

MONOGYNIA.

TRIENTALIS. Calyx heptaphyllus. Corolla ſeptempartita, æqualis, plana. Bacca exſucca, ſuturis variis dehiſcens.

europæa. 1. T. foliis lanceolatis integerrimis, *Sp. pl. p.* 488. *Fl. dan. t.* 86.

Pyrola, alſines flore, europæa. *Bauh. pin.* 191. *prodr.* 101.

In Carnia in ſylva prope pagum *Comeglians. I. Hoſt.* Fl. Iulio, Auguſto. Per.

ÆSCULUS. Calyx monophyllus, quinquedentatus. Petala quinque, calyci inſerta. Capſula trilocularis.

Hippo-Caſtanum. 1. Æ. floribus heptandris. *Sp. pl. p.* 488. *Miller illuſtr. ic. Hall. hiſt. n.* 1029.

Caſtanea equina. *Clus. hiſt. p.* 8.

Caſtanea folio multifido. *Bauh. pin.* 419.

Le Marronier d' Inde. Regn. bot.

Occurrit ad latera viarum, ad pagos, hortos, frequentior in Auſtria, quam in reliquis Provinciis. Fl. primo vere. Arb.

OCTANDRIA.

MONOGYNIA.

OENOTHERA. Calyx quadrifidus. Petala quatuor. Capſula cylindrica. infera. Semina nuda.

biennis. 1. Oe. foliis ovato-lanceolatis, planis, caule muricato-villoſo. *Syſt. veget. p.* 358. *Fl. dan. t.* 446.

Oenothera foliis ovato-lanceolatis, planis, caule hirſuto. *Hall. hiſt. n.* 994.

Onagra biennis. *Scop. carn. tom.* 1. *p.* 269.

Ad rivulos', fluvios, foſſas. Fl. æſtate.

EPILOBIUM. Calyx quadripartitus. Petala quatuor. Capſula oblonga, infera. Semina pappoſa.

anguſtifolium. 1. E. foliis ſparſis lineari-lanceolatis integerrimis venoſis, floribus inæqualibus. *Ait. Kew. tom.* 2. *p.* 4. *Curtis lond. Sp. pl. p.* 493.

Epilobium flore difformi, foliis lanceolatis, transverſim nervoſis. *Hall. hiſt. n.* 1000.

Epilobium Gesneri. *Allion pedem. tom.* 1. *p.* 279.

Chamænerion anguſtifolium. *Scop. carn. tom.* 1. *p.* 271.

Ad pedes montium, in ſylvis montanis, & ſubalpinis. Fl. æſtate. Per.

anguſtiſſimum. 2. E. foliis ſparſis linearibus obſolete denticulatis aveniis, petalis æqualibus integerrimis. *Ait. Kew. tom.* 2. *p.* 5. *Curtis magaz.* 76. *Weber dec. p.* 3.

Epilobium flore difformi, foliis linearibus. *Hall. hiſt. n.* 1001.

Epilobium Dodonæi. *Allion. pedem. tom.* 1. *p.* 279.

Epilobium roſmarinifolium. *Hæncke in Iacqu. collect. tom. 2. p.* 50.

Chamænerion paluſtre. *Scop. carn. tom.* 1. *p.* 271.

Lyſimachia Chamænerion dicta alpina. *Bauh. pin.* 116.

In Carniolia in udis agri labacenſis, ad pedes alpium wochinenſium, & montis *Nanas.* In Auſtria in declivibus, arenoſisque locis prope *Tulln, Kloſterneuburg, St. Pölten,* alibique ad Danubii ripas. In Tyroli circa Leontium. Fl. Iulio, Auguſto. Per.

Cave, ne novellos ramulos pro foliorum faſciculis ſumas.

hirſutum. 3. E. foliis ovato-lanceolatis, ſemiamplexicaulibus hirſutis, caule ramoſiſſimo, radice repente. *Curtis lond. Ait. Kew. tom.* 2. *p.* 5. *Sp. pl. p.* 494.

Epilobium grandiflorum. *Allion. pedem. tom.* 1. *p.* 279.

Epilobium ramoſum. *Huds. angl.* 162.

Lyſimachia filiquoſa major. *Cluſ. hiſt. p. LI.*

Lyſimachia ſiliquoſa hirſuta magno flore. *Bauh. pin.* 245.

Ad fluvios, foſſas, rivulos, ſepes humidas. Fl. æſtate. Per.

villoſum. 4. E. foliis oblongo-lanceolatis, dentatis, pubeſcentibus, caule tereti villoſo. *Curtis lond. Ait. Kew. tom.* 2. *p.* 5.

Chamænerion *parviflorum,* foliis oppoſitis lanceolatis ſerratis ſeſſilibus cauleque villoſis. *Schreb. ſpic. p.* 146.

Epilobium hirſutum. *Huds. angl.* 161. *Allion pedem. tom.* 1. *p.* 279.

Lyſimachia ſiliquoſa hirſuta, parvo flore. *Bauh. pin.* 245. *prodr.* 116.

Habitat & floret cum præcedente. Per.

monta-num. 5. E. foliis oppoſitis ovatis dentatis; ſtigmate quadrifido. *Sp. pl. p.* 494. *Curtis lond.*

Epilobium foliis ovato-lanceolatis, glabris, dentatis. *Hall. hiſt. n.* 996.

Chamænerion montanum. *Scop. carn. tom.* 1. *p.* 270.

In ſylvis, ad ſepes. Fl. æſtate.

roſeum. 6. E. foliis oppoſitis alternisque ovato-lanceolatis petiolatis ſerratis glabris; ſtigmate indiviſo. *Schreb. ſpic. p.* 147.

Epilobium minus flore albo. *Schmidel. op. Gesn.* 2. *faſc.* 2. *p.* 20. *t.* 20. *f.* 73.

In udis, & umbroſis. Fl. æſtate.

tetrago-num. 7. E. foliis lanceolatis denticulatis: imis oppoſitis, caule tetragono. *Sp. p. p.* 494. *Curtis lond.*

Epilobium foliis lanceolatis glabris dentatis. *Hall. hiſt. n.* 997.

Chamænerion tetragonum. *Scop. carn. tom.* 1. *p.* 271.

In rivulis, aliisque udis locis. Fl. æſtate. Per.

alpinum. 8. E. foliis oppoſitis ovato-lanceolatis integerrimis, ſiliquis ſeſſilibus, caule repente. *Sp. pl. p.* 495. *Fl. dan. t.* 322.

In alpibus ſtyriacis, carinthiacis, & tyrolenſibus. Fl. Iulio, Auguſto. Per.

CHLORA. Calyx octophyllus. Corolla monopetala, octofida. Capſula unilocularis, bivalvis, polyſperma.

perfoliata. 1. C. foliis perfoliatis. *Syſt. veget. p.* 361.

Gentiana foliis radicalibus ovatis, caulinis triangularibus perfoliatis, floribus octofidis. *Hall. hiſt. n.* 649.

Gentiana perfoliata. *Sp. pl. p.* 335. *Scop. carn. tom.* 1. *p.* 183.

Centaurium parvum flavo flore. *Cluſ. hiſt. p. CLXXX.*

Centaurium luteum perfoliatum. *Bauh. pin.* 278. *Zann. iſt. p.* 56. *t.* 20. 181. *n.* 1.

In udis Carniæ, Forijulii. Fl. Majo, Iunio.

VACCINIUM. Calyx ſuperus. Corolla monopetala. Filamenta receptaculo inſerta. Bacca quadrilocularis, polyſperma.

Myrtillus. 1. V. pedunculis unifloris, foliis ſerratis ovatis deciduis, caule angulato. *Sp. pl. p.* 498. *Fl. dan. t.* 974. *Scop. carn. tom.* 1. *p.* 273.

Vaccinium foliis rugoſis, ovato-lanceolatis, ſerratis, caule anguloſo. *Hall. hiſt. n.* 1020.

L' Airelle, ou Myrtille. Regn. bot.

In montanis, alpinis, & ſubalpinis. Fl. ab Aprili ad Iunium. Fr.

uliginoſum. 2. V. pedunculis unifloris, foliis integerrimis obovatis obtuſis lævibus. *Syſt. veget. p.* 363. *Fl. dan. t.* 231.

Vaccinium foliis integerrimis, ovatis, venoſis. *Hall. hiſt. n.* 1021.

Vitis idæa ſecunda. *Cluſ. hiſt. p.* 61.

In alpibus. Fl. poſt ſolutas nives. Fr.

Vitis idæa. 3. V. racemis terminalibus nutantibus, foliis obovatis revolutis integerrimis ſubtus punctatis. *Sp. pl. p.* 500. *Fl. dan. t.* 40. *Scop. carn. tom.* 1. *p.* 273.

Vaccinium foliis perennantibus ſubtus punctatis; oris contractis. *Hall. hiſt. n.* 1022.

In ſubalpinis, & alpinis. Fl. Majo, Iunio. Fr.

Leobenſes, conterminique Styriæ ſuperioris incolæ, quibus res bene domi, baccarum ſesqui-

menſuram cum libra una aceti in vaſe clauſo per quatuordecim dies macerant, ex maceratis ſuccum exprimunt, ex quo, addita ſufficienti quantitate ſachari albi, ſyrupum parant gratiſſimum. Levioris ſortis homines fructus, vaſi fictili ſine addito alio liquore commiſſos, in furno poſt extractum panem tamdiu detinent, donec pars aquoſa evaporaverit, ſaporque acidulus exaltatus fuerit. Tali methodo incorruptibiles factas, in recipiente, ad hoc negotium proprio, ſub communi nomine Krankenbeeren *conſervant. Quotiescunque in morbis æſtus nimius moderandus, ſitis mitiganda eſt, hi baccas ægris comedendas dant, ii ſyrupum, aqua dilutum, bibendum propinant. Hæc benevole mecum communicavit Cl. Kotnig M. D., & civitatis leobenſis Phyſicus.*

Oxycoccos. 4. V. foliis integerrimis revolutis ovatis, caulibus repentibus filiformibus nudis. *Sp. pl. p.* 500. *Fl. dan. t.* 80. *Scop. carn. tom.* 1. *p.* 272.

Vaccinium caule proſtrato, foliis ovato-lanceolatis, acutis, flore quadripartito. *Hall. hiſt. n.* 1023.

In paluſtribus: in Auſtria prope *Annaberg, Weitra;* in Carniolia prope Labacum. Fl. primo vere. Per.

Corolla quadripartita: laciniis revolutis.

ERICA. Calyx tetraphyllus. Corolla quadrifida. Filamenta receptaculo inſerta. Antheræ bifidæ. Capſula quadrilocularis.

vulgaris. 1. E. antheris ariſtatis, corollis campanulatis ſubæqualibus, calycibus duplicatis, foliis oppoſitis ſagittatis. *Syſt. veget. p.* 363. *Curtis lond. Fl. dan. t.* 677.

Erica foliis imis ſimplicibus adpreſſis, floralibus calcaratis. *Hall. hiſt. n.* 1012.

Erica Coris folio prima. *Clus. hiſt. p.* 41.

In montoſis ſylvaticis, ſubalpinis, & alpinis. Fl. Auguſto, Septembri. Fr.

herbacea. 2. E. antheris muticis exertis, corollis oblongis, ſtylo exerto, foliis quaternis, floribus ſecundis. *Syſt. veget. p.* 370. *Iacqu. auſtr. t.* 32. *Scop. carn. tom.* 1. *p.* 275.

Erica foliis patentibus, quaternis, acutis, caducis, flore calyce longiori, antheris eminentibus. *Hall. hiſt. n.* 1013.

Erica carnea. *Sp. pl. p.* 504.

Erica Coris folio octava. *Clus. hiſt. p.* 44.

Erica Coris folio nona. *Clus. hiſt. p.* 44.

In collibus, ſylvis montanis, & ſubalpinis. Fl. primo vere, quandoque & autumno. Fr.

Flores primum virent, clauſi ſunt; dein carnei fiunt, antherasque exertas habent.

DAPHNE. Calyx nullus. Corolla monopetala, infundibuliſormis, tubuloſa, marceſcens. Stamina corollæ tubo inſerta. Bacca monoſperma.

Mezereum. 1. D. floribus ſeſſilibus ternis caulinis, foliis lanceolatis deciduis. *Sp. pl. p.* 509. *Fl. dan. t.* 268.

Thymelæa ſpica cylindrica, ſuperne folioſa. *Hall. hiſt. n.* 1024.

Thymelæa Mezereum. *Scop. carn. tom.* 1. *p.* 276.

La Laureole femelle. Regn. bot.

In ſylvis montanis, & ſubalpinis. Fl. primo vere. Per.

Laureola. 2. D. racemis axillaribus, foliis lanceolatis glabris. *Sp. pl. p.* 510. *Iacqu. auſtr. t.* 183.

Thymelæa foliis perennantibus, ellipticis, floribus ex alis nutantibus. *Hall. hiſt. n.* 1025.

Thymelæa Laureola. *Scop. carn. tom.* 1. *p.* 276.

La Laureole mâle. Regn. bot.

In ſylvis montanis. Fl. primo vere. Fr.

Cneorum. 3. D. floribus faſciculatis terminalibus ſeſſilibus, foliis lanceolatis nudis mucronatis. *Syſt. veget. p.* 371. *Iacqu. auſtr. t.* 426.

Thymelæa foliis glabris linearibus, ariſtatis, floribus ſeſſilibus umbellatis. *Hall. hiſt. n.* 1027.

Thymelæa Cneorum. *Scop. carn. tom.* 1. *p.* 276.

Cneorum Matthioli. *Clus. hiſt. p.* 89.

La Laureole odorante, Bulliard. herb. t. 121.

In collibus herbidis, in alpinis, & ſubalpinis. Fl. a Majo ad Iulium. Fr.

alpina. 4. D. floribus ſeſſilibus aggregatis terminalibus lateralibusque, foliis lanceolatis tomentoſis. *Sp. pl. p.* 510. *Wulfen in Iacqu. collect. tom.* 3. *p.* 6.

Thymelæa foliis ellipticis, calveſcentibus, floribus aggregatis ſeſſilibus, hirſutis. *Hall. hiſt. n.* 1026.

Thymelæa candida. *Scop. carn. tom.* 1. *p.* 277.

Chamelæa pumila ſaxatilis flore pallido italica. *Barr. t.* 234.

Chamelæa alpina, folio inferne incano. *Bauh. pin.* 462.

In Forojulio in ſaxoſis circa Goritiam. In Carniolia ad pedem montis *Nanas*; prope Poſtoinam (*Adlersberg*). In Croatia in *mala Plaſtina*. In agro fluminenſi in *Rezzine*, *Grohovo*. Fl. Aprili, Majo, Iunio. Fr.

Caulis pedalis, ſæpe ſeſquipedalis. Rami diffuſi. Folia alia lanceolata, alia obovata, pauca emarginata. Iuniora hirſuta, dein calveſcentia. Flores veſpere ſuavi odore pollentes, candidi, pubeſcentes. Bacca oblonga, rubella, hinc linea impreſſa, a baſi ad apicem per medium excurrente, notata.

STELLERA. Calyx nullus. Corolla quadrifida. Stamina breviſſima in ſumma parte corollæ. Semen unicum, obtuſe mucronatum, corolla perſiſtente incluſum.

Paſſerina. 1. S. foliis linearibus, floribus quadrifidis. *Sp. pl. p.* 512. *Iacqu. collect. tom.* 1. *p.* 65. *ic. t.* 68. *Hall. hiſt. n.* 1028.

In Auſtriæ arvis. Fl. æſtate.

D I G Y N I A.

MOEHRINGIA. Calyx tetraphyllus. Petala quatuor. Capſula unilocularis, quadrivalvis.

muscoſa. 1. Mœhringia. *Sp. pl. p.* 515. *Iacqu. auſtr. t.* 449. *Scop. carn. tom.* 1. *p.* 277.

Alſine octoſtemon foliis linearibus connatis. *Hall. hiſt. n.* 860.

In humidis, umbroſis ſubalpinis, & alpinis. Fl. æſtate. Per.

T R I G Y N I A.

POLYGONUM. Calyx turbinatus, quinquepartitus. Corolla nulla. Semen unicum, angulatum.

Biſtorta. 1. P. caule ſimpliciſſimo monoſtachyo, foliis ovatis in petiolum decurrentibus. *Sp. pl. p.* 516. *Fl. dan. t.* 421. *Curtis lond.*

Polygonum radice lignoſa intorta, ſpica ovata, folioſis petiolis alatis. *Hall. hiſt. n.* 1559.

Biſtorta major. *Clus. hiſt. p.* LXIX.

La Biſtorte. Regn. bot.

In Tyrolis, Carinthiæ, Styriæ, Auſtriæ alpinis, ſubalpinis, alpium convallibus, pratis humidis. Fl. æſtate. Per.

viviparum. 2. P. caule ſimpliciſſimo monoſtachyo, foliis lanceolatis. *Sp. pl. p.* 516. *Fl. dan. t.* 13. *Scop. carn. tom.* 1. *p.* 279.

Polygonum radice lignoſa, intorta, foliis nervoſis, radicalibus ovato-lanceolatis, caulinis linearibus. *Hall. hiſt. n.* 1558.

Biſtorta minor. *Clus. hiſt. p. LXIX.*

In alpinis, & ſubalpinis. Fl. æſtate. Per.

amphibium. 3. P. floribus pentandris ſemidigynis, ſpica ovata. *Syſt. veget. p.* 377. *Curtis lond. Fl. dan. t.* 282.

Polygonum foliis ovato-lanceolatis, ciliatis, ſpicis ovatis. *Hall. hiſt. n.* 1565.

In foſſis, inundatis. Fl. æſtate. Per.

Perſicaria. 4. P. floribus hexandris ſemidigynis, pedunculis lævibus, ſtipulis ciliatis, ſpicis ovato-oblongis erectis. *Curtis lond. Sp. pl. p.* 518. *Fl. dan. t.* 702. *Scop. carn. tom.* 1. *p.* 279.

Polygonum foliis ovato-lanceolatis, ſubhirſutis, ſpicis ovatis, vaginis ciliatis. *Hall. hiſt. n.* 1557.

La Perſicaire. Regn. bot.

In cultis, humidis, & alibi. Fl. æſtate.

lapathifolium. 5. P. floribus hexandris digynis, ſtipulis muticis, pedunculis ſcabris, feminibus utrinque depreſſis. *Sibthorp. oxon. p.* 129. *Sp. pl. p.* 517.

Polygonum penſylvanicum. *Curtis lond.*

In Carniolia circa *Eck* inter ſegetes. *Zois.* Fl. Iulio, Auguſto.

Hydropiper. 6. P. floribus hexandris ſemidigynis, foliis lanceolatis undulatis, ſpicis filiformibus nutantibus. *Sibthorp. oxon. p.* 129. *Curtis lond. Sp. pl. p.* 517.

La Perſicaire brulante. Bulliard herb. t. 127.

In udis. Fl. æſtate.

mariti-mum. 7. P. floribus octandris trigynis axillaribus, foliis ovali-lanceolatis ſempervirentibus, caule ſuffrutescente. *Sp. pl. p.* 519.

Polygonum marinum. *Bauh. hiſt.* 3. *p.* 376. *Zann. iſt. p.* 216. *t.* 229.

In Litorali ad mare. Fl. Iulio. Per.

Folia tuberculis, in lineas digeſtis, aſpera, pleraque ad apicem margine revoluta, exſiccatione cœrulescentia.

avicula-re. 8. P. floribus octandris trigynis axillaribus, foliis lanceolatis, caule procumbente herbaceo. *Sp. pl. p.* 519. *Fl. dan. t.* 803. *Curtis lond. Scop. carn. tom.* I. *p.* 280.

Polygonum procumbens, foliis linearibus acutis, floribus ſolitariis. *Hall. hiſt. n.* 1560.

La Renouée, ou la Traînaſſe. Regn. bot.

In cultis, ruderatis, ad vias, & alibi. Fl. æſtate.

Fagopy-rum. 9. P. foliis cordato-ſagittatis, caule erectiusculo inermi, ſeminum angulis æqualibus. *Sp. pl. p.* 522.

Polygonum caule erecto, foliis ſagittatis, valvulis integerrimis. *Hall. hiſt. n.* 1563.

Le Bled noir, ou Sarraſin. Regn. bot.

In arvis, ad margines agrorum, pagos, & alibi. Fl. æſtate.

Convol-vulus. 10. P. foliis cordatis, caule volubili angulato, floribus obtuſatis. *Syſt. veget. p.* 379. *Curtis lond. Fl. dan. t.* 744. *Scop. carn. tom.* I. *p.* 279.

Polygonum caule volubili, foliis ſagittatis. *Hall. hiſt. n.* 1561.

In agris, ad vias, ſepes. Fl. Iunio, Iulio.

dumeto-rum. 11. P. foliis cordatis, caule volubili lævi, floribus carinato-alatis. *Sp. pl. p.* 522. *Fl. dan. t.* 756.

Polygonum caule volubili, foliis ſagittatis, valvulis ſeminalibus alatis. *Hall. hiſt. n.* 1562.

In dumetis, ad ſepes. Fl. Auguſto, Septembri.

TETRAGYNIA.

PARIS. Calyx tetraphyllus. Petala quatuor. Bacca globoſo-tetragona, quadrilocularis. Semina plura, gemino ordine incumbentia.

quadrifolia. 1. Paris. *Sp. pl. p.* 526. *Fl. dan. t.* 139. *Scop. carn. tom.* 1. *p.* 281. *Hall. hiſt. n.* 1006.

Le Raiſin de Renard. Regn. bot.

La Pariſette a quatre feuilles. Bulliard herb. t. 119.

In ſylvis umbroſis. Fl. Martio, Aprili. Per.

ADOXA. In flore capitulum terminante eſt corolla ſupera, quadrifida. Stamina octo. Styli quatuor. Bacca quadrilocularis inferne cum calyce bifido coalita. In floribus lateralibus corolla quinquefida. Stamina decem. Styli quinque. Bacca quinquelocularis.

Moſchatellina. 1. Adoxa. *Sp. pl. p.* 527. *Curtis lond. Fl. dan. t.* 94.

Moſchatella Adoxa. *Scop. carn. tom.* 1. *p.* 281.

Moſchatellina. *Hall. hiſt n.* 1005.

In ſylvis. Fl. Aprili. Per.

ELATINE. Calyx tetraphyllus. Petala quatuor. Capſula quadrilocularis, quadrivalvis, depreſſa.

Alſinaſtrum. 1. E. foliis verticillatis. *Sp. pl. p.* 527. *Hall. hiſt. n.* 857.

Alſinaſtrum galii folio. *Vaill. paris.* 6. *t.* 1. *f.* 6.

In Auſtria in foſſis circa *Hütteldorf. Weſthofen.* Fl. æſtate. Per.

ENNEANDRIA.

MONOGYNIA.

LAURUS. Calyx nullus. Corolla quadripartita, nuda. Nectarium glandulis tribus, biſetis, germen cingentibus. Filamenta interiora glandulifera. Drupa monoſperma.

nobilis. 1. L. foliis venoſis lanceolatis perennantibus, floribus quadrifidis dioicis. *Sp. pl. p.* 529. *Scop. carn. tom.* 1. *p.* 282.

Laurus foliis ovato-lanceolatis, ramis florigeris folio brevioribus. *Hall. hiſt. n.* 1602.

Le Laurier. Regn. bot.

In Tyroli calidiore, Forojulio, Litorali, copioſiſſime in agro fluminenſi, prope Voloscam, Lauranam, in *Abbazia*, viciniſque locis. Fl. primo vere. Arb.

HEXAGYNIA.

BUTOMUS. Involucrum univerſale ſimplex, triphyllum. Petala ſex. Capſulæ ſex polyſpermæ.

umbellatus. 1. Butomus. *Sp. pl. p.* 532. *Curtis lond. Fl. dan. t.* 604. *Scop. carn. tom.* 1. *p.* 283. *Hall. hiſt. n.* 1186.

In aquoſis. Fl. Iunio, Iulio. Per.

DECANDRIA.

MONOGYNIA.

DICTAMNUS. Calyx pentaphyllus. Petala quinque, patula. Filamenta punctis glanduloſis adſperſa. Capſulæ quinque, margine introrſum coalitæ.

albus. 1. D. foliis pinnatis, caule ſimplici. *Syſt. veget.* p. 397. *Iacqu. auſtr. t.* 428. *Scop. carn. tom.* 1. *p.* 284.

Fraxinella. *Clus. hiſt. p.* 99. *Hall. hiſt. n.* 1029.

La Fraxinelle. Regn. bot.

In aſperis, fruticoſis, ſylvaticis. Fl. Majo, Iunio. Per.

RUTA. Calyx quinquepartitus. Petala concava. Receptaculum punctis melliferis decem cinctum. Capſula lobata.

graveolens. 1. R. foliis decompoſitis, floribus lateralibus quadrifidis. *Syſt. veget. p.* 397. *Scop. carn. tom.* 1. *p.* 285. *Hall. hiſt. n.* 1003.

La Rue des Iardins. Regn. bot.

In aſperis, montoſis Croatiæ, Carnioliæ calidioris, Forijulii, Litoralis. Fl. æſtate. Per.

TRIBULUS. Calyx quinquepartitus. Petala quinque, patentia. Stylus nullus. Capſulæ quinque, gibbæ, ſpinoſæ, polyſpermæ.

terreſtris. 1. T. foliis ſubſexjugis ſubæqualibus, ſeminibus quadricornibus. *Syſt. veget. p.* 402. *Clus. hiſt. p.* CCXLI.

Tribulus foliis æqualibus duodenis, conjugatis, fructu quadricorni. *Hall. hiſt. n.* 947.

Tribulus terreſtris, ciceris folio, fructu aculeato. *Bauh. pin.* 350. *Zann. iſt. p.* 263. *t.* 222.

Flumine in vineis, ad vias, copioſiſſime circa *il Lazzaretto.* Fl. Majo, Iunio.

MONOTROPA. Calyx nullus. Petala decem; horum quinque exteriora baſi excavato-mellifera. Capſula pentagona., quinquevalvis.

Hypopithys. 1. M. floribus lateralibus octandris: terminali decandro. *Sp. pl. p.* 555. *Fl. dan. t.* 232.

Hypopithys ſpica florida nutante. *Hall. hiſt. n.* 1002.

Hypopithys multiflora. *Scop. carn. tom.* 1. *p.* 285.

In ſylvis umbroſis montanis, & ſubalpinis. Fl. Iunio, Iulio.

LEDUM. Calyx quinquefidus. Corolla plana, quinquepartita. Capſula quinquelocularis, baſi dehiscens.

paluſtre. 1. L. foliis linearibus, ſtaminibus corolla longioribus. *Retzii ſcand. p.* 100. *obs. faſc.* 4. *p.* 26. *Sp. pl. p.* 561. *Hæncke in Iacqu. collect. tom.* 2. *p.* 42. *Burgsdorf abb. t.* 83.

Ledum ſileſiacum. *Clus. hiſt. p.* 83.

Ciſtus Ledon foliis rosmarini ferrugineis. *Bauh. pin.* 467.

In Auſtriæ paluſtribus prope *Weitra. R. Liborius Miller.* Fl. Majo, Iunio. Fr.

RHODODENDRON. Calyx quinquepartitus. Corolla ſubinfundibuliformis. Stamina declinata. Capſula quinquelocularis.

ferrugineum. 1. R. foliis glabris ſubtus leproſis, corollis infundibuliformibus. *Sp. pl. p.* 405. *Iacqu. auſtr. t.* 255. *Scop. carn. tom.* 1. *p.* 286.

Rhododendron foliis ellipticis, glabris, ſubtus rubiginoſis. *Hall. hiſt. n.* 1015.

Ledi alpini genus alterum. *Clus. hiſt. p.* 82.

In alpibus. Fl. Iulio. Fr.

hirſutum. 2. R. foliis ciliatis nudis, corollis infundibuliformibus. *Sp. pl. p.* 562. *Iacqu. auſtr. t.* 98. *Scop. carn. tom.* 1. *p.* 286.

Rhododendron foliis ovatis, ciliatis, ſubtus punctatis. *Hall. hiſt. n.* 1016.

Ledum alpinum. *Clus. hiſt. p.* 81.

In alpinis, & ſubalpinis. Fl. Iulio. Fr.

Chamæciſtus. 3. R. foliis ciliatis, corollis rotatis. *Sp. pl. p.* 562. *Iacqu. auſtr. t.* 217. *Scop. carn. tom.* 1. *p.* 287.

Ledum foliis ſerpylli, ad margines cilii inſtar piloſis; flore purpureo. *Mich. gen. p.* 225. *t.* 106.

Chamæciſtus octavus. *Clus. hiſt. p.* 76.

In alpibus auſtriacis, ſtyriacis, carniolicis, tyrolenſibus, locis rupeſtribus calcareis. Fl. Iunio, Iulio. Fr.

ANDROMEDA. Calyx quinquepartitus. Corolla ovata: ore quinquefido. Capſula quinquelocularis.

cærulea. 1. A. pedunculis aggregatis, corollis ovatis, foliis ſparſis linearibus obtuſis planis. *Sp. pl. p.* 563. *Fl. dan. t.* 57.

In Carniæ umbroſis non procul a monte *Forca. I. Hoſt.* Fl. Iulio, Auguſto. Fr.

Polifolia. 2. A. pedunculis aggregatis, corollis ovatis, foliis alternis lanceolatis revolutis. *Sp. pl. p.* 564. *Fl. dan. t.* 54. *Pallas roſſic. t.* 71. *Hall. hiſt n.* 1017.

Rhododendron Polifolium. *Scop. carn. tom.* 1. *p.* 287.

Rosmarinus ſylveſtris aosmos. *Clus. hiſt. p.* 84.

In Carniolia in paluſtribus labacenſibus. *Scop.* In Carinthia in turfoſis alpium reichenavienſium ad S. Laurentium. *Wulfen.* Fl. ab Aprili ad Iunium. Fr.

PYROLA. Calyx quinquepartitus. Petala quinque. Capſula quinquelocularis, angulis dehiscens.

rotundifolia. 1. P. ſtaminibus adſcendentibus, piſtillo declinato. *Sp. pl. p.* 567. *Fl. dan. t.* 110.

Pyrola foliis ſubrotundis, tubis recurvis. *Hall. hiſt. n.* 1010.

La Pyrole. Regn. bot.

In ſylvis ſubalpinis. Fl. Iunio, Iulio. Per.

minor. 2 P. floribus racemoſis diſperſis, ſtaminibus piſtillisque rectis. *Sp. pl. p.* 567. *Fl. dan. t.* 55. *Scop. carn. tom.* 1. *p.* 288.

Pyrola foliis ſubrotundis, tubis rectis. *Hall. hiſt. n.* 1009.

Pyrola prima vulgatior. *Clus hiſt. p. CXVI.*

In ſylvis montanis, & ſubalpinis. Fl. Iunio, Iulio. Per.

ſecunda. 3. P. racemo unilaterali. *Sp. pl. p.* 567. *Fl. dan. t.* 402. *Scop. carn. tom.* 1. *p.* 289. *Hall. hiſt. n.* 1008.

Pyrola folio mucronato. *Riv. pent. t.* 136.

Pyrola ſecunda tenerior. *Clus. hiſt. p. CXVII.*

In ſylvis ſubalpinis, montanis. Fl. Iulio. Per.

uniflora. 4. P. ſcapo unifloro. *Sp. pl. p.* 568. *Fl. dan. t.* 8. *Scop. carn. tom.* 1. *p.* 289. *Hall. hiſt. n.* 1011.

Pyrola quarta minima. *Clus. hiſt. p.* CXVIII.

In ſylvis ſubalpinis, & in ipſis alpibus. Fl. Iunio, Iulio. Per.

Pyrola rotundifolia habet ſtigma incraſſatum, quinquedentatum: dentibus erectis, acutis. Simile habet P. uniflora, niſi quod majus ſit & peltatum. P. minori eſt peltatum, convexum, quinquelobum. Pyrolæ ſecundæ eſt ſtigma ſimile, & poris quinque pertuſum. Retzii obs. fasc. 1. p. 17.

ARBUTUS. Calyx quinquepartitus. Corolla ovata: ore quinqueſido: baſi diaphana. Bacca quinquelocularis.

Unedo. 1. A. caule arboreo, foliis glabris obtuſe ſerratis, panicula terminali, baccis polyſpermis. *Syſt. veget. p.* 407.

Arbutus. *Clus. hiſt. p.* 47.

Arbutus folio ſerrato. *Bauh. pin.* 460. *Miller. dict. t.* 48. *f.* 1. 2.

In Carnia in montibus *Forca, Chiadin. I. Hoſt.* Fl. Iunio. Fr.

Fructus frago ſimilis, orbiculatus, initio viridis, deinde flaveſcens, plene maturus rubro colore elegantiſſimo conſpicuus. Clus. l. c.

alpina. 2. A. caulibus procumbentibus, foliis rugoſis ſerratis. *Sp. pl. p.* 566. *Fl. dan. t.* 73. *Hall. hiſt. n.* 1019.

Vitis idæa prima. *Clus. hiſt. p.* 61.

In alpibus. Fl. poſt ſolutas nives. Fr.

Baccæ ſunt primo virides, dein rubræ, ultimo nigræ. Clus. l. c.

Uva urſi. 3. A. caulibus procumbentibus, foliis integerrimis. *Sp. pl. p.* 566. *Fl. dan. t.* 33. *Scop. carn. tom.* 1. *p.* 288. *Hall. hiſt. n.* 1018.

Uva urſi. *Cluſ. hiſt. p. 63.*

In ſubalpinis, & alpinis. Fl. Majo, Iunio. Fr.

Fructus ſunt primo virides, rubri dum maturi.

DIGYNIA.

CHRYSOSPLENIUM. Corolla nulla. Capſula biroſtris, unilocularis, polyſperma. Calyx in flore primario quinquefidus, ſtamina decem. In floribus ſerioribus calyx quadrifidus, ſtamina octo.

alternifolium. 1. C. foliis alternis. *Sp. pl. p. 569. Fl. dan. t. 366. Scop. carn. tom. 1. p. 290. Hall. hiſt. n. 1548.*

Saxifraga aurea. *Cluſ. hiſt. p. 308.*

In umbroſis, humidis montanis, ſubalpinis, & alpinis. Fl. a Majo ad Iulium. Per.

SAXIFRAGA. Calyx quinquepartitus. Corolla pentapetala. Capſula biroſtris, unilocularis, polyſperma.

Cotyledon. 1. S. foliis radicatis aggregatis lingulatis cartilagineo-ſerratis, caule paniculato. *Sp. pl. p. 570. Fl. dan. t. 241. Wulfen in Iacqu. collect. tom. 4. p. 291. Scop. carn. tom. 1. p. 291.*

Saxifraga Aizoon. *Iacqu. auſtr. t. 438.*

In ſaxoſis, herbidisque alpinis, & ſubalpinis. Fl. Iulio, Auguſto. Per.

Variat foliis brevibus ſubrotundis; oblongis latioribus, & anguſtiſſimis; caule multifloro, & paucifloro; pedunculis ſimplicibus, ac ramoſis; petalis punctatis, atque immaculatis.

mutata. 2. S. foliis radicalibus aggregatis lingulatis cartilagineo-ſerratis, caule racemoſo. *Sp. pl. p.* 570. *Wulfen in Iacqu. collect. tom.* 1. *p.* 284. *ic. t.* 466.

Saxifraga foliorum ora cartilaginea, rariſſime dentata, petalis maculoſis. *Hall. hiſt. n.* 979. *t.* 16.

Geum alpinum majus viscoſum, foliis oblongo-rotundis, atro-rubentibus & croceis floribus. *Scheuch. it.* 2. *p.* 124.

In Auſtria *in der Brein* in alpium præcipitiis. *Iacqu.* In Carinthiæ alpibus reichenavienſibus. *Wulfen.* Ego in Styriæ alpibus judenburgenſibus non procul a lacu, *Bürgerſee* dicto. Fl. Auguſto. Per.

Caule magis viscido, floribus quadruplo majoribus, petalis lineari-lanceolatis diſtincta a præcedente.

androſacea. 3. S. foliis lanceolatis obtuſis piloſis, caule nudo bifloro. *Sp. pl. p.* 571. *Iacqu. auſtr. t.* 389.

Saxifraga foliis hirſutis, ellipticis & tridentatis, caule paucifloro. *Hall. hiſt. n.* 984.

Saxifraga pyrenaica. *Scop. carn. tom.* 1. *p.* 296.

In alpium humidiusculis. Fl. Iulio, Auguſto. Per.

Folia ſunt lanceolata, baſi anguſtata, intetegerrima, vel ſubtridentata.

cæſia. 4. S. foliis linearibus perforato-punctatis aggregatis recurvatis, caule multifloro. *Syſt. veget. p.* 411. *Iacqu. auſtr. t.* 374. *Scop. carn. tom.* 1. *p.* 294.

Saxifraga foliis craſſis, duris, recurvis, ſubtus ſulcatis, baſi ciliatis. *Hall. hiſt. n.* 982.

Saxifraga alpina, minima, foliis cæſiis, deorſum incurvis. *Tourn. inſt.* 253.

Sedum minus nonum. *Clus. hiſt. p. LXI.*

In alpium ſaxoſis, rupeſtribus. Fl. Iulio, Auguſto. Per.

burſeriana. 5. S. foliis aggregatis imbricatis ſubulatis lævibus, caule ſubnudo unifloro. *Sp. pl. p.* 572. *Wulfen in Iacqu. miscell. tom.* 1, *p.* 152. *t.* 17. *f.* 3.

Saxifraga foliis campactis, plicatis, pungentibus, caule viscido, paucifloro. *Hall. hiſt. n.* 983.

Saxifraga alpina foliis glaucis acutis, monanthos, caule folioſo. *Segu. ver.* 3. *p.* 201. *t.* 5. *f.* 2.

In Carniolia in alpibus javerburgenſibus. In Tyroli circa Leontium. In Carinthia in alpibus ortenburgenſibus prope *Spital.* In Auſtria in præcipitiis montis *Schneeberg*, *die Saugrube* dictis, ubique ſolo rupeſtri calcareo. Fl. Iunio. Per.

ſedoides. 6. S. foliis aggregatis alternis oppoſitisque ſublanceolatis, flore pedunculato. *Sp. pl. p.* 572. *Wulfen in Iacqu. miscell. tom.* 2. *p.* 134. *t.* 21. *f.* 2.

Saxifraga foliis mollibus, ſubhirſutis, ellipticis, caule paucifloro. *Hall. hiſt. n.* 985.

Saxifraga Trichodes. *Scop. carn. tom.* 1. *p.* 295. *t.* 15.

In alpium humidis. Fl. Iulio, Auguſto. Per.

Folia ſunt lineari-lanceolata, pilis albis, uti tota planta, petalis exceptis, adſperſa, ad caulium baſin aggregata, reliqua ſparſa. Caulem terminat pedunculus plerumque uniflorus, raro bi- vel triflorus.

bryoides. 7. S. foliis ciliatis inflexis imbricatis, caule nudiusculo unifloro. *Sp. pl. p.* 572. *Wulfen in Iacqu. miscell. tom.* 2. *p.* 49. *t.* 5. *f.* 1. *Scop. carn. tom.* 1. *p.* 295. *t.* 15.

Saxifraga foliis lanceolatis, ciliatis, compactis, caule unifloro. *Hall. hiſt. n.* 969.

In ſummis alpium jugis. Fl. Iulio, Auguſto. Per.

Caules inſperſi glandulis pedicellatis. Folia linearia, inflexa, ciliata, pungentia. Petala *oblonga, pallide lutea. Quo humidiora loca, in qui-*

bus creſcit, eo caules magis elongantur, & folia, quæ cæteroquin imbricata, per elongatos caules ſparguntur.

aſpera. 8. S. foliis caulinis lanceolatis alternis ciliatis, caulibus procumbentibus. *Syſt. veget. p.* 413. *Iacqu. auſtr. app. t.* 31.

Saxifraga foliis duris, ciliatis, caule ramoſo, paucifloro. *Hall. hiſt. n.* 970.

Saxifraga ſedi foliis crenatis, aſperis. *Tourn. inſt.* 252. *Scheuch. it.* 2. *p.* 140. *t.* 20. *f.* 3.

In Carniæ, Carinthiæ, Tyrolis alpibus locis rupeſtribus, ſaxoſis humidis. Fl. Iulio, Auguſto. Per.

tenella. 9. S. foliis lineari-ſubulatis ciliatis mucronatis, caulibus floriferis erectis ſubnudis paucifloris. *Wulfen in Iacqu. collect. tom.* 3. *p.* 144. *t.* 17.

In Carniolia in rupeſtribus umbroſis alpium wochinenſium, in aſcenſu montis *czerna Perſt, Sucha.* Fl. Iunio. Per.

Folia viridia pilis brevibus, deciduis ciliata, apice in ſpinulam deſinentia. Corollæ petala immaculata. Differt a Saxifraga bronchiali, cui folia ſpinoſo-ciliata, apice valde pungentia, ſi non omnia, inferiora ſaltem rubentia; petala interne, ad baſin punctis luteſcentibus, apicem verſus ſanguineis inſperſa.

aizoides. 10. S. foliis caulinis lanceolatis ſparſis ciliato-aculeatis, caulibus procumbentibus. *Retzii ſcand. p.* 102. *obs. faſc.* 1. *p.* 18.

Saxifraga foliis ellipticis, ciliatis, caule ſimplici, folioſo, paucifloro. *Hall. hiſt. n.* 971.

Saxifraga autumnalis. *Sp. pl. p.* 575. *Scop. carn. tom.* 1. *p.* 293. *t.* 14. *Fl. dan. t.* 72. *Hæncke in Iacqu. collect. tom.* 2. *p.* 18.

Sedum minus ſextum ſive alpinum primum. *Cluſ. hiſt. p. LX.*

In alpibus ad rivulos, & in aliis humidis locis. Fl. Iulio, Auguſto, Septembri. Per.

Folia modo ad radicem aggregata, modo ſparſa, alia aculeis ciliata, alia margine nuda. Petala in aliquibus individuis immaculata ſunt, in aliis punctis aurantiacis, vel purpureis notata; dum & ipſa vel pallide lutea, vel aurantiaca ſunt.

oppoſitifolia. 11. S. foliis caulinis ovatis oppoſitis imbricatis: ſummis ciliatis. *Sp. pl. p.* 575. *Fl. dan. t.* 34.

Saxifraga foliis ovatis quadrangulo - imbricatis, ramis procumbentibus. *Fl. lapp. n.* 179. *t.* 2. *f.* 1.

Saxifraga caule repente, foliis quadrifariam imbricatis, cartilagineis, ciliatis. *Hall. hiſt. n.* 980.

In Styria in ſummis jugis alpium judenburgenſium. In Tyroli in alpibus leontinis, In alpibus carniolicis. Fl. Iulio. Per.

ſtellaris. 12. S. foliis ſerratis, caule nudo ramoſo, petalis acuminatis. *Sp. pl. p.* 572. *Fl. dan. t.* 23. *Scop. carn. tom.* 1. *p.* 292. *t.* 13. *Wulfen in Iacqu. collect. tom.* 1. *p.* 202. *t.* 13.

Saxifraga foliis rhomboideis, acutè ſerratis, caule nudo ramoſo. *Hall. hiſt. n.* 973.

Saniculæ montanæ alterius ſpecies ſecunda. *Cluſ. hiſt. p.* 308.

In alpium riguis, humidis, ad rivulos. Fl. Iulio, Auguſto. Per.

cuneifolia. 13. S. foliis cuneiformibus obtuſiſſimis repandis, caule nudo paniculato. *Sp. pl. p.* 574. *Scop. carn. tom.* 1. *p.* 291. *t.* 13.

Saxifraga foliis petiolatis, obtuſis, caule fragili, nudo, ramoſo. *Hall. hiſt. n.* 974.

In rupeſtribus Carnioliæ. Fl. Iunio. Per.

Geum. 14. S. folis reniformibus, dentatis, caule nudo paniculato. *Sp. pl. p.* 574. *Scop. carn. tom.* 1. *p.* 292.

In Litorali in monte majore. Fl. Majo, Iunio. Per.

Folia erecta: ima ex rotundo ovata; ſuperiora cordata. Petioli pilis inſperſi. Petala punctata.

rotundifolia. 15. S. foliis caulinis reniformibus dentatis petiolatis, caule paniculato. *Sp. pl. p.* 576. *Scop. carn. tom.* 1. *p.* 290.

Saxifraga foliis reniformibus, acute ſerratis, hirſutis, petiolatis, caule ramoſo. *Hall. hiſt. n.* 975.

Sanicula montana ſecunda. *Clus. hiſt. p.* 307.

La Saxifrage a feuilles rondes. Bulliard herb. t. 327.

In alpinis, & ſubalpinis. Fl. Iulio, Auguſto. Per.

granulata. 16. S. foliis caulinis reniformibus lobatis, caule ramoſo, radice granulata. *Sp. pl. p.* 576. *Fl. dan. t.* 514.

Saxifraga foliis radicalibus reniformibus, obtuſe dentatis, caulinis palmatis. *Hall. hiſt. n.* 976.

Saxifraga alba. *Clus. hiſt. p.* 308.

La Saxifrage, ou Percepierre. Regn. bot.

In pratis, collibus herbidis. Fl. Majo. Per.

bulbifera. 17. S. foliis palmato - lobatis: caulinis ſeſſilibus, caule ramoſo bulbifero. *Sp. pl. p.* 577. *Fl. dan. t.* 390.

Saxifraga ad folia bulbos gerens. *Bauh. pin.* 309.

In pratis. Fl. Majo. Per.

tridactylites. 18. S. foliis caulinis cuneiformibus trifidis alternis, caule erecto ramoſo. *Sp. pl. p.* 578. *Curtis lond. Scop. carn. tom.* 1. *p.* 297.

Saxifraga foliis petiolatis, trilobatis, caule erecto, ramoſo & folioſo. *Hall. hiſt. n.* 986.

Sedum tridactylites tectorum. *Bauh. pin.* 285.

In muris vetuſtis, collibus apricis ſaxoſis, pratis ſicciorибus. Fl. primo vere.

In agro fluminenſi, & in Auſtria prope Bertholdsdorf, Medling *inveni varietatem foliis omnibus integris.*

petræa. 19. S. foliis cuneiformibus: radicalibus integris tridentatisque: caulinis quinquedentatis, ſuperioribus trifidis, pedunculis ſubtrifloris. *Vahl. skr. nat. ſelsk.* 2. *p.* 10. *Retzii ſcand. p.* 103.

Saxifraga foliis radicalibus congeſtis ſubquinquedentatis. *Fl. dan. t.* 68.

Saxifraga tridactylites. *Fl. N.* 544. *t.* 9. *f.* 1. 3.

Saxifraga adſcendens. *Wulfen in Iacqu. collect. tom.* 1. *p.* 197. *t.* 11. 12. *f.* 1. 2.

Saxifraga hypnoides. *Scop. carn. tom.* 1. *p.* 296. *t.* 16.

In alpibus. Fl. Iulio, Auguſto. Per.

geranioides. 20. S. foliis radicalibus reniformibus quinquelobis multifidis, caulinis linearibus, caule ſubnudo ramoſo. *Syſt. veget. p.* 414. *Gouan illuſtr. p.* 28. *t.* 18. *f.* 2.

Saxifraga petræa. *Wulfen in Iacqu. collect. tom.* 1. *p.* 200. *ic. t.* 81.

Saxifraga alba petræa. *Pon. in Cluſ. hiſt. p.* CCCXXXVII.

In Carniolia in ſpecu Timavi ſub *Krainburg;* in collibus *Pezh, Skerli,* alibique in rupibus *in der Wochein.* Fl. Iunio. Per.

Tota planta villoſa, viſcida. Folia radicalia caulinaque inferiora longe petiolata, lobata. Quæ quinque lobis conſtant, ſubreniformia ſunt. Caulina ſuperiora ſunt vel linearia, vel lanceolata, integra, vel bifida, vel trifida. Nec rarum invenire eſt individua, in quibus quaſi omnia folia integra ſunt.

ceſpitoſa. 21. S. foliis radicalibus aggregatis linearibus integris trifidisque, caule erecto ſubnudo ſubbifloro. *Sp. pl. p.* 578. *Scop. carn. tom.* 1. *p.* 293. *t.* 14. *Fl. dan. t.* 71. *Wulfen in Iacqu. collect. tom.* 1. *p.* 290.

Saxifraga groenlandica. *Fl. N.* 689. *t.* 7. *f.* 1. *Syſt. veget. p.* 414.

Saxifraga muscoides. *Wulfen in Iacqu. miscell. tom.* 2. *p.* 123. *Syſt. veget. p.* 413.

Saxifraga foliis partim integris, partim trifidis. *Segu. ver.* 1. *p.* 451. *t.* 9. *f.* 4.

In ſummis alpium jugis. Fl. Iulio, Auguſto. Per.

moſchata. 22. S. foliis radicalibus aggregatis integris trifidisque una cum caule glutinoſis hirſutis, floribus longe pedunculatis. *Syſt. veget. p.* 414. *Wulfen in Iacqu. miscell. tom.* 2. *p.* 128. *t.* 21. *f.* 21.

Saxifraga foliis palmato - trifidis, ſuperne lineatis hirſutis; caule aſcendente multifloro; petalis trinerviis. *Vill. proſp. p.* 47.

Saxifraga exarata. *Allion. pedem. tom.* 2. *p.* 73. *t.* 88. *f.* 2.

Sedulum, quod moſchatella alpina lutea vocari poteſt. *Gesn. fasc. p.* 25. *t.* 6. *f.* 31.

In Tyroli in excelſis ſchleinitzenſibus alpibus prope Leontium. *Wulfen*; ego in agro œnopontino in montibus non procul a *Lüſenſerferner.* Fl. Iunio, Iulio. Auguſto. Per.

Chryſomela oleracea devaſtat in horto ſaxifragam Aizoidem, androſaceam, ceſpitoſam; hanc ſpeciem intactam relinquit.

SCLERANTHUS. Calyx monophyllus. Corolla nulla. Semina duo, calyce incluſa.

annuus. 1. S. calycibus fructus patulis. *Sp. pl. p.* 580. *Fl. dan. t.* 504.

Scleranthus floribus patulis. *Hall. hiſt. n.* 1551.

Knavel annuum. *Scop. carn. tom.* 1. *p.* 298.

In arvis. Fl. æſtate.

GYPSOPHILA. Calyx campanulatus, monophyllus, angulatus. Petala quinque, ovata, ſeſſilia. Capſula globoſa, unilocularis.

repens. 1. G. foliis lanceolatis, ſtaminibus corolla emarginata brevioribus. *Syſt. veget. p.* 415. *Iacqu. auſtr. t.* 407.

Saponaria radice lignoſa, maxima, foliis pulpoſis, petalis emarginatis. *Hall. hiſt. n.* 905.

In alpium declivibus. Fl. Iulio, Auguſto. Per.

panicula-ta. 2. G. foliis lineari-lanceolatis: inferioribus ſcabris, ſtaminibus minutis, ſtylis corolla longioribus. *Ait. Kew. tom.* 2. *p.* 84. *Sp. pl. p.* 583. *Iacqu. auſtr. app. t.* 1.

In Pannonia in campis aridis circa Budam. In Litorali in aridis circa Tergeſtum. Fl. æſtate. Per.

muralis. 3. G. foliis linearibus planis, calycibus aphyllis, caule dichotomo, petalis crenatis. *Sp. pl. p.* 583.

Saponaria foliis filiformibus, caule diffuſo, brachiato. *Hall. hiſt. n.* 903.

Lychnis parva paluſtris, foliis acutis lanceolatis, floſculis purpureis. *Mentz. pug. t.* 7. *f.* 4.

In ſiccis. Fl. æſtate.

ſaxifra-ga. 4. G. foliis linearibus, calycibus angulatis: ſquamis quatuor, corollis emarginatis. *Syſt. veget. p.* 416.

Tunica calycibus pelviformibus. *Hall. hiſt. n.* 902.

Tunica ſaxifraga. *Scop. carn. tom.* 1. *p.* 300.

Habitat & floret cum præc. Per.

SAPONARIA. Calyx monophyllus, nudus. Petala quinque, unguiculata. Capſula oblonga, unilocularis.

officinalis. 1. S. calycibus cylindricis, foliis ovato-lanceolatis. *Sp. pl. p.* 584. *Fl. dan. t.* 543. *Curtis lond.*

Saponaria foliis ovato-lanceolatis, trinerviis, floribus tubuloſis umbellatis. *Hall. hiſt. n.* 908.

Lychnis officinalis. *Scop. carn. tom.* 1. *p.* 303.

La Saponaire. Regn. bot.

Ad ſepes, vias, fluvios, in pratis humidis. Fl. Iulio, Auguſto. Per.

Vaccaria. 2. S. calycibus pyramidatis quinquangularibus, foliis ovatis acuminatis ſeſſilibus. *Sp. pl. p.* 585.

Saponaria calycibus pyramidatis, quinquangularibus, foliis ovatis connatis. *Hall. hiſt. n.* 907.

Lychnis Vaccaria. *Scop. carn. tom.* 1. *p.* 303.

Inter ſegetes. Fl. Iunio, Iulio.

Ocymoides. 3. S. calycibus cylindricis villoſis, caulibus dichotomis procumbentibus. *Sp. pl. p.* 585. *Iacqu. auſtr. app. t.* 23.

Saponaria caulibus decumbentibus, nodoſis, foliis ovato-lanceolatis, calycibus tubuloſis hirſutis. *Hall. hiſt. n.* 909.

In Carinthia, Tyroli in montium ſaxoſis, arenoſis. Fl. æſtate. Per.

DIANTHUS. Calyx cylindricus, monophyllus: baſi ſquamis quatuor. Petala quinque, unguiculata. Capſula cylindrica, unilocularis.

* *Flores aggregati.*

barbatus. 1. D. floribus aggregatis fasciculatis: ſquamis calycinis ovato-ſubulatis tubum æquantibus, foliis lanceolatis. *Sp. pl. p.* 586.

Tunica barbata. *Scop. carn. tom.* I. *p.* 298.

Armerius ſimplici flore pannonicus. *Clus. hiſt. p.* 287.

In Carnioliæ, Croatiæ, Forijulii herbidis montanis, & alpinis. Fl. Iunio, Iulio.

carthuſianorum. 2. D. floribus ſubaggregatis, ſquamis calycinis ovatis ariſtatis tubo brevioribus, foliis linearibus trinerviis. *Smith in Linn. Trans. tom.* 2. *on. Dianthus, Sp. pl. p.* 586.

Tunica fasciculis trifloris ramos terminantibus, ſtipulis ariſtatis maximis. *Hall. hiſt. n.* 899.

Tunica carthuſianorum. *Scop. carn. tom.* I. *p.* 299.

Caryophyllus arvenſis, calyculo florum numeroſo. *Loes. pruſſ.* 37. *f.* 7.

In pratis, collibus, & alibi. Fl. æſtate. Per.

Armeria. 3. D. floribus aggregatis fasciculatis: ſquamis calycinis lanceolatis villoſis tubum æquantibus. *Sp. pl. p.* 586. *Fl. dan. t.* 230.

Tunica floribus umbellatis, ſquamis calycinis hirſutis, mucronatis, tubum æquantibus. *Hall. hiſt. n.* 900.

In ſteribus, ſylvaticis. Fl. Iunio, Iulio.

prolifer. 4. D. floribus aggregatis capitatis: ſquamis calycinis ovatis, obtuſis muticis tubum ſuperantibus. *Sp. pl. p.* 587. *Fl. dan. t.* 221.

Tunica capitulo ovali, compacto, calyce univerſali quadrifolio. *Hall. hiſt. n.* 901.

Tunica prolifera. *Scop. carn. tom.* 1. *p.* 299.

In collibus, arvis agri vindobonenſis, goritienſis. Fl. æſtate.

Non rarum eſt invenire ſpecimina calyce unicum florem includente. Hanc varietatem D. diminutum Linn. ſuſpicatur eſſe Cl. Smith, *aſſerens, nullum ſpecimen Dianthi diminuti in herbario Linnæano proſtare; & quotquot in aliorum botanicorum herbariis ſub tali nomine vidit, nil aliud eſſe, quam ſupradictam D. proliferi varietatem.*

** *Flores ſolitarii, plures in eodem caule.*

deltoides. 5. D. floribus ſolitariis, ſquamis calycinis ovato-lanceolatis acutis ſubbinis, foliis obtuſiuſculis ſubpubescentibus, petalis crenatis. *Smith. l. c. Sp. pl. p.* 588. *Fl. dan. t.* 577.

Caryophyllus ſylveſtris ſeptimus. *Clus. hiſt. p.* 285.

In Auſtriæ, Pannoniæ ſylvaticis. Fl. æſtate. Per.

Folia ciliata. Squamæ calycinæ binæ, quandoque quatuor. Petala purpurea. Limbus facie punctis argenteis, ad baſin linea transverſali ſaturate purpurea notatus. At in montis hainburgenſis apricis, aridis legi varietatem, in qua petala plerumque alba, punctis argenteis deſtituta. Hæc varietas eſt Caryophyllus ſylveſtris ſeptimus Cluſii, &, ſecundum Cl. Smith, *D. glaucus Linnæi.*

plumarius. 6. D. floribus ſolitariis, ſquamis calycinis ſubovatis breviſſimis obtuſiſſimis muticis, corollis multifidis. *Smith. l. c.*

Tunica foliis glaucis, patentibus, floribus ſerratis, faucibus lanuginoſis. *Hall. hiſt. n.* 897.

Caryophyllus flos ſylveſtris quintus. *Clus. hiſt. p.* 283.

In Auſtriæ, Pannoniæ apricis rupeſtribus. Fl. Iunio, Iulio. Per.

ſuperbus. 7. D. floribus ſolitariis paniculatis, ſquamis calycinis breviſſimis acuminatis, petalis multifido-capillaribus, caule erecto. *Smith. l. c. Sp. pl. p.* 589. *Fl. dan. t.* 578. *Iacqu. obs.* 1. *p.* 40. *t.* 25.

Tunica petalis minutiſſime laciniatis. *Hall. hiſt. n.* 898.

Caryophyllus ſylveſtris ſextus. *Clus. hiſt. p.* 284.

In Auſtriæ, Pannoniæ paluſtribus. In reliquarum provinciarum ſylvaticis, ſubalpinis, & alpinis. Fl. Iulio, Auguſto. Per.

virgineus. 8. D. floribus ſolitariis, caulibus paucifloris, ſquamis calycinis breviſſimis obtuſiſſimis binis, petalis crenatis. *Smith. l. c. Iacqu. auſtr. app. t.* 15. *Sp. pl. p.* 590.

Dianthus rupeſtris. *Linn. ſuppl. p.* 240.

Tunica virginea. *Scop. carn. tom.* 1. *p.* 302.

Caryophyllus ſylveſtris repens multiflorus. *Bauh. pin.* 209. *prodr.* 104. *Herb. burs. vol.* 11. *p.* 99.

In aſperis, herbidis Litoralis, Forijulii, Carnioliæ, Carinthiæ, & Tyrolis calidioris. Fl. æſtate. Per.

*** *Caule unifloro herbaceo.*

alpinus. 9. D. caule unifloro, petalis crenatis, ſquamis calycinis exterioribus foliaceis tubum ſubæquantibus. *Smith l. c. Iacqu. auſtr. t.* 52.

Caryophyllus ſylveſtris ſecundus. *Clus. hiſt. p.* 282.

In alpinis, & ſubalpinis. Fl. æſtate. Per.

TRIGYNIA.

CUCUBALUS. Calyx inflatus. Petala quinque, unguiculata, abſque corona ad faucem. Capſula trilocularis.

baccife-rus. 1. C. calycibus campanulatis, petalis diſtantibus, pericarpiis coloratis, ramis divaricatis. *Sp. pl. p.* 591.

Viscago baccifera petalis ſerratis. *Hall. hiſt. n.* 912.

Lychnis baccifera. *Scop. carn. tom.* 1. *p.* 306.

Alſine repens major. *Clus. hiſt. p.* CLXXXIII.

In nemoribus, ad ſepes Auſtriæ, Pannoniæ, Styriæ, Carnioliæ. Fl. æſtate. Per.

Behen. 2. C. calycibus ſubgloboſis glabris reticulato-venoſis, capſulis trilocularibus, corollis ſubnudis. *Sp. pl. p.* 591. *Fl. dan. t.* 857.

Viscago caule brachiato, nodoſo, calycibus inflatis, venoſis. *Hall. hiſt. n.* 913.

Lychnis Behen. *Scop. carn. tom.* 1. *p.* 305.

Le Carnillet Behen. Bulliard herb. t. 321.

In pratis, montanis ſylvaticis, ad vias. Fl. æſtate. Per.

viscoſus. 3. C. floribus lateralibus undique decumbentibus, caule indiviſo, foliis baſi reflexis. *Sp. pl. p.* 592.

Lychnis orientalis maxima, bugloſſi folio undulato. *Tourn. cor.* 24. *itin.* 2. *p.* 361.

In pratis ſiccioribus Pannoniæ inter *Bruck an der Leitha* & *Bahrendorf.* Fl. Iunio. Per.

Tota planta viscoſa. Folia radicalia in petiolum decurrentia. Caulina amplexicaulia: omnia margine undulata, & ad baſin reflexa.

Otites. 4. C. floribus dioicis, petalis linearibus indiviſis. *Sp. pl. p.* 594. *Fl. dan. t.* 518.

Viſcago floribus verticillatis, ſpicatis, ſexu diſtinctis, petalis linearibus. *Hall. hiſt. n.* 920.

Lychnis Otites. *Scop. carn. tom.* 1. *p.* 305.

Saſamoides magnum ſalmanticum. *Clus. hiſt. p.* 295.

In ſterilibus, aſperis. Fl. æſtate. Per.

pumilio. 5. C. caulibus unifloris flore brevioribus. *Syſt. veget. p.* 420.

Silene pumilio. *Iacqu. auſtr. app. t.* 2. *Wulfen in Iacqu. collect. tom.* 2. *p.* 126. *t.* 10.

Lychnis pumilio. *Scop. carn. tom.* 1. *p.* 304.

Caryophyllus ſylveſtris octavus. *Clus. hiſt. p.* 285.

In alpibus Styriæ, Carinthiæ, Tyrolis. Fl. Iulio, Auguſto, Septembri. Per.

SILENE. Calyx ventricoſus. Petala quinque, unguiculata: coronata ad faucem. Capſula trilocularis.

anglica. 1. S. hirſuta, petalis emarginatis, floribus erectis, fructibus reflexis pedunculatis alternis. *Sp. pl. p.* 594. *Curtis lond.*

Viſcago ceraſtii foliis, vasculis pendulis anglica. *Dill. elth.* 417.

In Carniolia inter ſegetes circa *Eck. Zois.* Fl. Iunio.

Folia oppoſita, obovato-lanceolata, integerrima, ſubviscida, punctis prominulis aſpera, baſi pilis longis ciliata.

nutans. 2. S. petalis bifidis, floribus lateralibus ſecundis cernuis, panicula nutante. *Sp. pl. p.* 596. *Fl. dan. t.* 242.

Viscago foliis lanceolatis, hirſutis, floribus paniculatis nutantibus. *Hall. hiſt. n.* 915.

Lychnis ſylveſtris octava. *Clus. hiſt. p.* 291.

In pratis, collibus, montoſis ſylvaticis, in alpibus. Fl. a Majo ad Iulium. Per.

conica. 3. S. calycibus fructus conicis ſtriis triginta, foliis mollibus, petalis bifidis. *Sp. pl. p.* 598. *Iacqu. auſtr. t.* 253.

Lychnis conica. *Scop. carn. tom.* 1. *p.* 308.

In collibus, inter ſegetes agri vindobonenſis circa *Weinhaus.* Fl. Iunio, Iulio.

inaperta. 4. S. caule dichotomo paniculato, calycibus lævibus, petalis breviſſimis emarginatis, foliis glabris lanceolatis. *Sp. pl. p.* 600.

Lychnis inaperta. *Scop. carn. tom.* 1. *p.* 309.

Viscago lævis, inaperto flore. *Dill. elth.* 424. *t.* 315. *f.* 407.

Circa Tergeſtum. *Scop.* In Carnia prope pagum *Caneva. I. Hoſt.* Fl. Iunio, Iulio.

Calyx manet clauſus usque ad perfectam ſeminum maturitatem. Petala alia bifida, alia integra: omnia calyce, & ſtaminibus breviora. Ehrh. beytr. tom. 6. p. 107.

noctiflora. 5. S. calycibus decemangularibus: dentibus tubum æquantibus, caule dichotomo, petalis bifidis. *Syſt. veget. p.* 421.

Viscago foliis ovato-lanceolatis, hirſutis, caule brachiato, calycibus venoſis. *Hall. hiſt. n.* 911.

Ocymoides noctiflorum. *Cam. hort. p.* 109. *t.* 34.

In Auſtriæ ſylvis, ad ſepes. Fl. æſtate.

Armeria. 6. S. floribus fasciculatis faſtigiatis, foliis ſuperioribus cordatis glabris, petalis integris. *Sp. pl. p.* 422. *Fl. dan. t.* 559.

Viscago foliis glabris, ovato-lanceolatis, floribus umbellatis. *Hall. hiſt. n.* 916.

Lychnis ſylveſtris prima. *Cluſ. hiſt. p.* 288.

In Tyroli ad pedes montium circa *Sterzing, Mittewalde* & *Brunegen.* Fl. Iulio, Auguſto.

rupeſtris. 7. S. floribus erectis, petalis emarginatis, calycibus teretibus, foliis lanceolatis. *Sp. pl. p.* 602. *Fl. dan. t.* 4. *Hænke in Iacqu. collect. tom.* 2. *p.* 85.

Viscago foliis lanceolatis, caulibus brachiatis, calycibus infundibuliformibus, floribus laxe umbellatis, erectis. *Hall. hiſt. n.* 917.

[illegible] fa[illegible]ilis, alpina, glabra, pumila. *Tourn. inſt.* 338. *Sche[illegible]. it.* 2. *p.* 137.

Alſine alpina glabra. *Bauh. pin.* 251. *prodr.* 118.

In alpibus ſtyriacis, carinthiacis, tyrolenſibus. Fl. Iulio, Auguſto. Per.

alpeſtris. 8. S. petalis [illegible]dentatis, caule dichotomo, capſulis [illegible] oblongis, foliis lineari - lanceolatis glabris erectis, pedunculis viscidis. *Ait. Kew. tom.* 2. *p.* 98. *Iacqu. auſtr. t.* 96.

Lychnis alpeſtris. *Linn. ſuppl. p.* 244.

Lychnis ſylveſtris decima. *Clus. hiſt. p.* 291.

In ſubalpinis, & alpinis. Fl. æſtate.

Saxifraga. 9. S. caulibus ſubunifloris, pedunculis longitudine caulis, floribus hermaphroditis feminseisque, petalis bifidis. *Syſt. veget. p.* 422.

Lychnis Saxifraga. *Scop. carn. tom.* 1. *p.* 308.

Lychnis minor ſaxifraga. *Segu. ver.* 1. *p.* 431. *t.* 6. *f.* 1.

In declivibus alpium carniolicarum, carnicarum, tyrolenſium. Fl. æſtate. Per.

acaulis. 10. S. acaulis depreſſa, petalis emarginatis. *Syſt. veget. p.* 422. *Fl. dan. t.* 21.

Viscago foliis gramineis, caule breviſſimo, unifloro. *Hall. hiſt. n.* 919.

Ocymoides muscoſus. *Clus. hiſt. p.* CCCXL.

In alpibus auſtriacis, ſtyriacis. Fl. a Iunio ad Septembrim. Per.

Saxifraga groenlandica. *Fl. N.* 689. *t.* 7. *f.* 1. *Syſt. veget. p.* 414.

Saxifraga muscoides. *Wulfen in Iacqu. miscell. tom.* 2. *p.* 123. *Syſt. veget. p.* 413.

Saxifraga foliis partim integris, partim trifidis. *Segu. ver.* 1. *p.* 451. *t.* 9. *f.* 4.

In ſummis alpium jugis. Fl. Iulio, Auguſto. Per.

moſchata. 22. S. foliis radicalibus aggregatis integris trifidisque una cum caule glutinoſis hirſutis, floribus longe pedunculatis. *Syſt. veget. p.* 414. *Wulfen in Iacqu. miscell. tom.* 2. *p.* 128. *t.* 21. *f.* 21.

Saxifraga foliis palmato-trifidis, ſuperne lineatis hirſutis; caule aſcendente multifloro; petalis trinerviis. *Vill. proſp. p.* 47.

Saxifraga exarata. *Allion. pedem. tom.* 2. *p.* 73. *t.* 88. *f.* 2.

Sedulum, quod moſchatella alpina lutea vocari poteſt. *Gesn. fasc. p.* 25. *t.* 6. *f.* 31.

In Tyroli in excelſis ſchleinitzenſibus alpibus prope Leontium. *Wulfen*; ego in agro œnopontino in montibus non procul a *Lüſenſerferner.* Fl. Iunio, Iulio. Auguſto. Per.

Chryſomela oleracea devaſtat in horto ſaxifragam Aizoidem, androſaceam, ceſpitoſam; hanc ſpeciem intactam relinquit.

SCLERANTHUS. Calyx monophyllus. Corolla nulla. Semina duo, calyce incluſa.

annuus. 1. S. calycibus fructus patulis. *Sp. pl. p.* 580. *Fl. dan. t.* 504.

Scleranthus floribus patulis. *Hall. hiſt. n.* 1551.

Knavel annuum. *Scop. carn. tom.* 1. *p.* 298.

In arvis. Fl. æſtate.

GYPSOPHILA. Calyx campanulatus, monophyllus, angulatus. Petala quinque, ovata, ſeſſilia. Capſula globoſa, unilocularis.

repens. 1. G. foliis lanceolatis, ſtaminibus corolla emarginata brevioribus. *Syſt. veget. p.* 415. *Iacqu. auſtr. t.* 407.

Saponaria radice lignoſa, maxima, foliis pulpoſis, petalis emarginatis. *Hall. hiſt. n.* 905.

In alpium declivibus. Fl. Iulio, Auguſto. Per.

paniculata. 2. G. foliis lineari-lanceolatis: inferioribus ſcabris, ſtaminibus minutis, ſtylis corolla longioribus. *Ait. Kew. tom.* 2. *p.* 84. *Sp. pl. p.* 583. *Iacqu. auſtr. app. t.* 1.

In Pannonia in campis aridis circa Budam. In Litorali in aridis circa Tergeſtum. Fl. æſtate. Per.

muralis. 3. G. foliis linearibus planis, calycibus aphyllis, caule dichotomo, petalis crenatis. *Sp. pl. p.* 583.

Saponaria foliis filiformibus, caule diffuſo, brachiato. *Hall. hiſt. n.* 903.

Lychnis parva paluſtris, foliis acutis lanceolatis, floſculis purpureis. *Mentz. pug. t.* 7. *f.* 4.

In ſiccis. Fl. æſtate.

ſaxifraga. 4. G. foliis linearibus, calycibus angulatis: ſquamis quatuor, corollis emarginatis. *Syſt. veget. p.* 416.

Tunica calycibus pelviformibus. *Hall. hiſt. n.* 902.

Tunica ſaxifraga. *Scop. carn. tom.* 1. *p.* 300.

Habitat & floret cum præc. Per.

SAPONARIA. Calyx monophyllus, nudus. Petala quinque, unguiculata. Capſula oblonga, unilocularis.

officinalis. 1. S. calycibus cylindricis, foliis ovato-lanceolatis. *Sp. pl. p.* 584. *Fl. dan. t.* 543. *Curtis lond.*

Saponaria foliis ovato-lanceolatis, trinerviis, floribus tubuloſis umbellatis. *Hall. hiſt. n.* 908.

Lychnis officinalis. *Scop. carn. tom.* 1. *p.* 303.

La Saponaire. Regn. bot.

Ad ſepes, vias, fluvios, in pratis humidis. Fl. Iulio, Auguſto. Per.

Vaccaria. 2. S. calycibus pyramidatis quinquangularibus, foliis ovatis acuminatis ſeſſilibus. *Sp. pl. p.* 585.

Saponaria calycibus pyramidatis, quinquangularibus, foliis ovatis connatis. *Hall. hiſt. n.* 907.

Lychnis Vaccaria. *Scop. carn. tom.* 1. *p.* 303.

Inter ſegetes. Fl. Iunio, Iulio.

Ocymoides. 3. S. calycibus cylindricis villoſis, caulibus dichotomis procumbentibus. *Sp. pl. p.* 585. *Iacqu. auſtr. app. t.* 23.

Saponaria caulibus decumbentibus, nodoſis, foliis ovato-lanceolatis, calycibus tubuloſis hirſutis. *Hall. hiſt. n.* 909.

In Carinthia, Tyroli in montium ſaxoſis, arenoſis. Fl. æſtate. Per.

DIANTHUS. Calyx cylindricus, monophyllus: baſi ſquamis quatuor. Petala quinque, unguiculata. Capſula cylindrica, unilocularis.

* *Flores aggregati.*

barbatus. 1. D. floribus aggregatis faſciculatis: ſquamis calycinis ovato-ſubulatis tubum æquantibus, foliis lanceolatis. *Sp. pl. p.* 586.

Tunica barbata. *Scop. carn. tom.* 1. *p.* 298.

Armerius ſimplici flore pannonicus. *Cluſ. hiſt. p.* 287.

In Carnioliæ, Croatiæ, Forijulii herbidis montanis, & alpinis. Fl. Iunio, Iulio.

carthuſianorum. 2. D. floribus ſubaggregatis, ſquamis calycinis ovatis ariſtatis tubo brevioribus, foliis linearibus trinerviis. *Smith in Linn. Trans. tom.* 2. *on. Dianthus, Sp. pl. p.* 586.

Tunica faſciculis trifloris ramos terminantibus, ſtipulis ariſtatis maximis. *Hall. hiſt. n.* 899.

Tunica carthuſianorum. *Scop. carn. tom.* 1. *p.* 299.

Caryophyllus arvenſis, calyculo florum numeroſo. *Loes. pruſſ.* 37. *f.* 7.

In pratis, collibus, & alibi. Fl. æſtate. Per.

Armeria. 3. D. floribus aggregatis faſciculatis: ſquamis calycinis lanceolatis villoſis tubum æquantibus. *Sp. pl. p.* 586. *Fl. dan. t.* 230.

Tunica floribus umbellatis, ſquamis calycinis hirſutis, mucronatis, tubum æquantibus. *Hall. hiſt. n.* 900.

In ſteribus, ſylvaticis. Fl. Iunio, Iulio.

prolifer. 4. D. floribus aggregatis capitatis: ſquamis calycinis ovatis, obtuſis muticis tubum ſuperantibus. *Sp. pl. p.* 587. *Fl. dan. t.* 221.

Tunica capitulo ovali, compacto, calyce univerſali quadrifolio. *Hall. hiſt. n.* 901.

Tunica prolifera. *Scop. carn. tom.* 1. *p.* 299.

In collibus, arvis agri vindobonenſis, goritienſis. Fl. æſtate.

Non rarum eſt invenire ſpecimina calyce unicum florem includente. Hanc varietatem D. diminutum Linn. ſuſpicatur eſſe Cl. Smith, *aſſerens, nullum ſpecimen Dianthi diminuti in herbario Linnæano proſtare; & quotquot in aliorum botanicorum herbariis ſub tali nomine vidit, nil aliud eſſe, quam ſupradictam D. proliferi varietatem.*

** *Flores ſolitarii, plures in eodem caule.*

deltoides. 5. D. floribus ſolitariis, ſquamis calycinis ovato-lanceolatis acutis ſubbinis, foliis obtuſiuſculis ſubpubescentibus, petalis crenatis. *Smith. l. c. Sp. pl. p.* 588. *Fl. dan. t.* 577.

Caryophyllus ſylveſtris ſeptimus. *Clus. hiſt. p.* 285.

In Auſtriæ, Pannoniæ ſylvaticis. Fl. æſtate, Per.

Folia ciliata. Squamæ calycinæ binæ, quandoque quatuor. Petala purpurea. Limbus facie punctis argenteis, ad baſin linea transverſali ſaturate purpurea notatus. At in montis hainburgenſis apricis, aridis legi varietatem, in qua petala plerumque alba, punctis argenteis deſtituta. Hæc varietas eſt Caryophyllus ſylveſtris ſeptimus Cluſii, &, ſecundum Cl. Smith, *D. glaucus Linnæi.*

plumarius. 6. D. floribus ſolitariis, ſquamis calycinis ſubovatis breviſſimis obtuſiſſimis muticis, corollis multifidis. *Smith. l. c.*

Tunica foliis glaucis, patentibus, floribus ſerratis, faucibus lanuginoſis. *Hall. hiſt. n.* 897.

Caryophyllus flos ſylveſtris quintus. *Clus. hiſt. p.* 283.

In Auſtriæ, Pannoniæ apricis rupeſtribus. Fl. Iunio, Iulio. Per.

ſuperbus. 7. D. floribus ſolitariis paniculatis, ſquamis calycinis breviſſimis acuminatis, petalis multifido-capillaribus, caule erecto. *Smith. l. c. Sp. pl. p.* 589. *Fl. dan. t.* 578. *Iacqu. obs.* 1. *p.* 40. *t.* 25.

Tunica petalis minutiſſime laciniatis. *Hall. hiſt. n.* 898.

Caryophyllus ſylveſtris ſextus. *Clus. hiſt. p.* 284.

In Auſtriæ, Pannoniæ paluſtribus. In reliquarum provinciarum ſylvaticis, ſubalpinis, & alpinis. Fl. Iulio, Auguſto. Per.

virgineus. 8. D. floribus ſolitariis, caulibus paucifloris, ſquamis calycinis breviſſimis obtuſiſſimis binis, petalis crenatis. *Smith. l. c. Iacqu. auſtr. app. t.* 15. *Sp. pl. p.* 590.

Dianthus rupeſtris. *Linn. ſuppl. p.* 240.

Tunica virginea. *Scop. carn. tom.* 1. *p.* 302.

Caryophyllus ſylveſtris repens multiflorus. *Bauh. pin.* 209. *prodr.* 104. *Herb. burs. vol.* 11. *p.* 99.

In aſperis, herbidis Litoralis, Forijulii, Carnioliæ, Carinthiæ, & Tyrolis calidioris. Fl. æſtate. Per.

*** *Caule unifloro herbaceo.*

alpinus. 9. D. caule unifloro, petalis crenatis, ſquamis calycinis exterioribus foliaceis tubum ſubæquantibus. *Smith l. c. Iacqu. auſtr. t.* 52.

Caryophyllus ſylveſtris ſecundus. *Clus. hiſt. p.* 282.

In alpinis, & ſubalpinis. Fl. æſtate. Per.

TRIGYNIA.

CUCUBALUS. Calyx inflatus. Petala quinque, unguiculata, abſque corona ad faucem. Capſula trilocularis.

bacciferus. 1. C. calycibus campanulatis, petalis diſtantibus, pericarpiis coloratis, ramis divaricatis. *Sp. pl. p.* 591.

Viscago baccifera petalis ſerratis. *Hall. hiſt. n.* 912.

Lychnis baccifera. *Scop. carn. tom.* 1. *p.* 306.

Alſine repens major. *Clus. hiſt. p.* CLXXXIII.

In nemoribus, ad ſepes Auſtriæ, Pannoniæ, Styriæ, Carnioliæ. Fl. æſtate. Per.

Behen. 2. C. calycibus ſubgloboſis glabris reticulato-venoſis, capſulis trilocularibus, corollis ſubnudis. *Sp. pl. p.* 591. *Fl. dan. t.* 857.

Viscago caule brachiato, nodoſo, calycibus inflatis, venoſis. *Hall. hiſt. n.* 913.

Lychnis Behen. *Scop. carn. tom.* 1. *p.* 305.

Le Carnillet Behen. Bulliard herb. t. 321.

In pratis, montanis ſylvaticis, ad vias. Fl. æſtate. Per.

viscoſus. 3. C. floribus lateralibus undique decumbentibus, caule indiviſo, foliis baſi reflexis. *Sp. pl. p.* 592.

Lychnis orientalis maxima, bugloſſi folio undulato. *Tourn. cor.* 24. *itin.* 2. *p.* 361.

In pratis ſiccioribus Pannoniæ inter *Bruck an der Leitha* & *Bahrendorf*. Fl. Iunio. Per.

Tota planta viscoſa. Folia radicalia in petiolum decurrentia. Caulina amplexicaulia: omnia margine undulata, & ad baſin reflexa.

Otites. 4. C. floribus dioicis, petalis linearibus indiviſis. *Sp. pl. p.* 594. *Fl. dan. t.* 518.

Viſcago floribus verticillatis, ſpicatis, ſexu diſtinctis, petalis linearibus. *Hall. hiſt. n.* 920.

Lychnis Otites. *Scop. carn. tom.* 1. *p.* 305.

Saſamoides magnum ſalmanticum. *Clus. hiſt. p.* 295.

In ſterilibus, aſperis. Fl. æſtate. Per.

pumilio. 5. C. caulibus unifloris flore brevioribus. *Syſt. veget. p.* 420.

Silene pumilio. *Iacqu. auſtr. app. t.* 2. *Wulfen in Iacqu. collect. tom.* 2. *p.* 126. *t.* 10.

Lychnis pumilio. *Scop. carn. tom.* 1. *p.* 304.

Caryophyllus ſylveſtris octavus. *Clus. hiſt. p.* 285.

In alpibus Styriæ, Carinthiæ, Tyrolis. Fl. Iulio, Auguſto, Septembri. Per.

SILENE. Calyx ventricoſus. Petala quinque, unguiculata: coronata ad faucem. Capſula trilocularis.

anglica. 1. S. hirſuta, petalis emarginatis, floribus erectis, fructibus reflexis pedunculatis alternis. *Sp. pl. p.* 594. *Curtis lond.*

Viſcago ceraſtii foliis, vasculis pendulis anglica. *Dill. elth.* 417.

In Carniolia inter ſegetes circa *Eck. Zois.* Fl. Iunio.

Folia oppoſita, obovato-lanceolata, integerrima, ſubviscida, punctis prominulis aſpera, baſi pilis longis ciliata.

nutans. 2. S. petalis bifidis, floribus lateralibus ſecundis cernuis, panicula nutante. *Sp. pl. p.* 596. *Fl. dan. t.* 242.

Viſcago foliis lanceolatis, hirſutis, floribus paniculatis nutantibus. *Hall. hiſt. n.* 915.

Lychnis ſylveſtris octava. *Clus. hiſt. p.* 291.

In pratis, collibus, montoſis ſylvaticis, in alpibus. Fl. a Majo ad Iulium. Per.

conica. 3. S. calycibus fructus conicis ſtriis triginta, foliis mollibus, petalis bifidis. *Sp. pl. p.* 598. *Iacqu. auſtr. t.* 253.

Lychnis conica. *Scop. carn. tom.* I. *p.* 308.

In collibus, inter ſegetes agri vindobonenſis circa *Weinhaus.* Fl. Iunio, Iulio.

inaperta. 4. S. caule dichotomo paniculato, calycibus lævibus, petalis breviſſimis emarginatis, foliis glabris lanceolatis. *Sp. pl. p.* 600.

Lychnis inaperta. *Scop. carn. tom.* I. *p.* 309.

Viscago lævis, inaperto flore. *Dill. elth.* 424. *t.* 315. *f.* 407.

Circa Tergeſtum. *Scop.* In Carnia prope pagum *Caneva. I. Hoſt.* Fl. Iunio, Iulio.

Calyx manet clauſus usque ad perfectam ſeminum maturitatem. Petala alia bifida, alia integra : omnia calyce, & ſtaminibus breviora. Ehrh. beytr. tom. 6. p. 107.

noctiflora. 5. S. calycibus decemangularibus: dentibus tubum æquantibus, caule dichotomo, petalis bifidis. *Syſt. veget. p.* 421.

Viscago foliis ovato-lanceolatis, hirſutis, caule brachiato, calycibus venoſis. *Hall. hiſt. n.* 911.

Ocymoides noctiflorum. *Cam. hort. p.* 109. *t.* 34.

In Auſtriæ ſylvis, ad ſepes. Fl. æſtate.

Armeria. 6. S. floribus fasciculatis faſtigiatis, foliis ſuperioribus cordatis glabris, petalis integris. *Sp. pl. p.* 422. *Fl. dan. t.* 559.

Viscago foliis glabris, ovato-lanceolatis, floribus umbellatis. *Hall. hiſt. n.* 916.

Lychnis ſylveſtris prima. *Cluſ. hiſt. p.* 288.

In Tyroli ad pedes montium circa *Sterzing, Mittewalde* & *Brunegen.* Fl. Iulio, Auguſto.

rupeſtris. 7. S. floribus erectis, petalis emarginatis, calycibus teretibus, foliis lanceolatis. *Sp. pl. p.* 602. *Fl. dan. t.* 4. *Hænke in Iacqu. collect. tom.* 2. *p.* 85.

Viscago foliis lanceolatis, caulibus brachiatis, calycibus infundibuliformibus, floribus laxe umbellatis, erectis. *Hall. hiſt. n.* 917.

Lychnis ſaxatilis, alpina, glabra, pumila. *Tourn. inſt.* 338. *Scheuch. it.* 2. *p.* 137.

Alſine alpina glabra. *Bauh. pin.* 251. *prodr.* 118.

In alpibus ſtyriacis, carinthiacis, tyrolenſibus. Fl. Iulio, Auguſto. Per.

alpeſtris. 8. S. petalis quadridentatis, caule dichotomo, capſulis ovato - oblongis, foliis lineari - lanceolatis glabris erectis, pedunculis viscidis. *Ait. Kew. tom.* 2. *p.* 98. *Iacqu. auſtr. t.* 96.

Lychnis alpeſtris. *Linn. ſuppl. p.* 244.

Lychnis ſylveſtris decima. *Clus. hiſt. p.* 291.

In ſubalpinis, & alpinis. Fl. æſtate.

Saxifraga. 9. S. caulibus ſubunifloris, pedunculis longitudine caulis, floribus hermaphroditis femineisque, petalis bifidis. *Syſt. veget. p.* 422.

Lychnis Saxifraga. *Scop. carn. tom.* 1. *p.* 308.

Lychnis minor ſaxifraga. *Segu. ver.* 1. *p.* 431. *t.* 6. *f.* 1.

In declivibus alpium carniolicarum, carnicarum, tyrolenſium. Fl. æſtate. Per.

acaulis. 10. S. acaulis depreſſa, petalis emarginatis. *Syſt. veget. p.* 422. *Fl. dan. t.* 21.

Viscago foliis gramineis, caule breviſſimo, unifloro. *Hall. hiſt. n.* 919.

Ocymoides muscoſus. *Clus. hiſt. p.* CCCXL.

In alpibus auſtriacis, ſtyriacis. Fl. a Iunio ad Septembrim. Per.

STELLARIA. Calyx pentaphyllus, patens. Petala quinque, bipartita. Capſula unilocularis, polyſperma.

nemorum. 1. S. foliis cordatis petiolatis, panicula pedunculis ramoſis. *Sp. pl. p.* 603. *Fl. dan. t.* 271.

Myoſotis foliis petiolatis, cordatis, tubis ternis. *Hall. hiſt. n.* 886.

In nemoribus, ascenditque ad ipſas alpes. Fl. Iunio, Iulio. Per.

Holoſtea. 2. S. foliis lanceolatis, ſerrulatis, petalis bifidis. *Sp. pl. p.* 603. *Scop. carn. tom.* 1. *p.* 314. *Curtis lond. Fl. dan. t.* 698.

In nemoribus. Fl. Aprili, Majo. Per.

graminea. 3. S. foliis linearibus integerrimis, floribus paniculatis. *Sp. pl. p.* 604. *Miller illuſtr. ic.*

Alſine foliis lanceolatis lineatis, petiolis ramoſis. *Hall. hiſt. n.* 882.

In pratis, ad ſepes, ſylvarum margines. Fl. æſtate. Per.

hypericifolia. 4. S. foliis lanceolatis baſi ciliatis, caule procumbente. *Allion. pedem. tom.* 2. *p.* 115.

Stellaria *dilleniana*, foliis lanceolatis integerrimis, caule procumbente. *Leers herb. p.* 108.

Stellaria *media*, foliis lineari-lanceolatis integerrimis glaucis decuſſatis, petalis calyce dimidio majoribus. *Sibthorp. oxon. p.* 141.

Alſine foliis lanceolatis, petalis bipartitis, petiolis unifloris. *Hall. hiſt. n.* 881.

Stellaria uliginoſa. *Roth. germ. tom.* 1. *p.* 191.

Stellaria fontana. *Wulfen in Iacq. collect. tom.* 1. *p.* 327.

Alſine uliginoſa. *Vill. proſp. p.* 48.

Alſine aquatica folio gratiolæ, ſtellato flore. *Dill. giſſ. p.* 58.

In uliginoſis, udis alpinis, & ſubalpinis. Fl. æſtate. Per.

bulboſa. 5. S. foliis lanceolato-ovatis, caule ſubdichotomo, pedunculis elongatis unifloris, radice repente bulbifera. *Wulfen in Iacqu. collect. tom.* 3. *p.* 21. *ic. t.* 468.

In Carniolia in agro labacenſi *unterm Thurn. Wulfen;* prope arcem *Röttmansdorf auf der Gemeinde. Zois.* Fl. Aprili, Majo. Per.

Ceraſtoides. 6. S. foliis oblongis, pedunculis ſubbifloris. *Sp. pl. p.* 604. *Wulfen in Iacqu. collect. tom.* 1. *p.* 254. *t.* 19. *Fl. N.* 951. *t.* 6. *f.* 2. *Fl. dan. t.* 92.

Myoſotis foliis lanceolatis glabris, petiolis refractis. *Hall. hiſt. n.* 890.

In Carinthia in editioribus Taureri malnitzenſis apricis paſcuis. *Wulfen.*

biflora. 7. S. foliis ſubulatis, ſcapis ſubbifloris, petalis emarginatis, germinibus oblongis, calycibus ſtriatis. *Sp. pl. p.* 604. *Fl. dan. t.* 12.

Sagina ramis erectis bifloris. *Fl. lapp. n.* 158.

Alſine foliis linearibus, obtuſis, calycibus viſcidis. *Hall. hiſt. n.* 863.

Stellaria ciliata. *Scop. carn. tom.* 1. *p.* 315. *t.* 17.

Arenaria Polygonoides. *Wulfen in Iacqu. collect. tom.* 1. *p.* 241. *t.* 15.

Alſine polygonoides, foliis brevibus, flore albo. *Segu. ver.* 3. *p.* 177. *t.* 4. *f.* 1.

In ſummis jugis alpium carniolicarum, carinthiacum, tyrolenſium.

ARENARIA. Calyx pentaphyllus, patens. Petala quinque, integra. Capſula unilocularis, polyſperma.

biflora. 1. A. foliis obovatis obtuſis, caulibus procumbentibus, pedunculis lateralibus bifloris. *Syſt. veget. p.* 423. *Allion. pedem. tom.* 2. *p.* 111. *t.* 44. *f.* 1. *t.* 64. *f.* 3. *Wulfen in Iacqu. collect. tom.* 1. *p.* 250. *ic. t.* 83.

Alſine caule recto proſtrato, foliis ovatis. *Hall. hiſt. n.* 877.

In ſummis Styriæ, Carinthiæ, Tyrolis alpium jugis. Fl. Iulio, Auguſto. Per.

trinervia. 2. A. foliis ovatis acutis petiolatis nervoſis. *Sp. pl. p.* 605. *Curtis lond. Fl. dan. t.* 329.

Alſine foliis ovato-lanceolatis trinerviis. *Hall. hiſt. n.* 878.

Alſine plantaginis folio. *Bauh. hiſt.* 3. *p.* 364.

In cultis, ſylvaticis. Fl. Majo, Iunio.

multicaulis. 3. A. foliis ovatis enerviis ſeſſilibus acutis, corollis calyce majoribus. *Syſt. veget. p.* 423. *Iacqu. en. p.* 75.

Alſine foliis ovato-lanceolatis, ciliatis, petalis calyce majoribus. *Hall. hiſt. n.* 876. *t.* 17.

In ſummis alpium jugis ſaxoſis. Fl. æſtate. Per.

ſerpyllifolia. 4. A. foliis ſubovatis acutis ſeſſilibus, corollis calyce brevioribus. *Sp. pl. p.* 606. *Curtis lond.*

Alſine foliis ovato-lanceolatis, ſubhirſutis, petalis calyce brevioribus. *Hall. hiſt. n.* 875.

Stellaria ſerpyllifolia. *Scop. carn. tom.* 1. *p.* 319.

In cultis, ſylvaticis, alibique in aridis. Fl. æſtate.

rubra. 5. A. foliis filiformibus, ſtipulis membranaceis vaginantibus. *Sp. pl. p.* 606. *Fl. dan. t.* 740.

Alſine foliis linearibus, ſtipulis ovato-lanceolatis argenteis. *Hall. hiſt. n.* 872.

In arenoſis aridis, & humidis paluſtribus, & ad mare. Fl. æſtate.

verna. 6. A. foliis ſubulatis, caulibus paniculatis, calycibus acuminatis, ſtriatis. *Syſt. veget. p.* 424. *Iacqu. auſtr. t.* 404.

In Auſtriæ collibus apricis, ſaxoſis aſperis. Fl. æſtate. Per.

Caules inferne glabri, ſuperne, ut pedunculi, denſe hirſutuli. Folia radicalia & caulina ſunt ſetacea, acuta: ramea lanceolata, concava. Capſula conica. Iacqu. l. c.

laricifolia. 7. A. foliis ſetaceis, caule ſuperne nudiuſculo, calycibus ſubhirſutis. *Sp. pl. p.* 607. *Iacqu. auſtr. t.* 272.

Alſine foliis linearibus anguſtiſſimis, calycibus tubuloſis, villoſis. *Hall. hiſt. n.* 869.

In alpinis, & ſubalpinis. Fl. Iulio, Auguſto. Per.

Folia ima conferta, quandoque faſciculata: ſuperiora oppoſita. Omnia ſubulata, ſeſſilia, ad lentem ciliata, facie plana, dorſo convexa. Flos magnus, campanulatus. Foliola calycis villoſa, ſtriata, oblonga, acuminata. Petala plana, calyce longiora, oblonga, obtuſiſſima, lineis impreſſis ramoſis notata. Filamenta alterna breviora. Capſula ovata, trivalvis: valvulis bifidis. Iacqu. l. c.

ſtriata. 8. A. foliis linearibus erectis appreſſis, calycibus oblongis ſtriatis. *Sp. pl. p.* 608. *Allion. pedem. tom.* 2. *p.* 113. *t.* 26. *f.* 4.

Arenaria *Cherleroides*, caule baſi nodoſo, ſuffruticoſo, foliis lineari-lanceolatis, calycinis æqualibus; caulibus ſubunifloris. *Vill. proſp. p.* 48. *t.* 30. *f.* 1, 2.

In alpibus tyrolenſibus, ſtyriacis, carniolicis. Fl. Iulio, Auguſto. Per.

Radix emittit caules plures, diffuſos. Folia ima conferta, ſuperiora oppoſita. Omnia lineari-lanceolata, margine ciliata. Pedunculi hirſutuli, ſubbiflori. Calycis foliola dorſo ſtriata, margine albida.

recurva. 9. A. foliis radicalibus confertis recurvis ſubulatis, caule ſimplici ſubtrifloro. *Allion. pedem. tom.* 2. *p.* 113. *t.* 89. *f.* 3. *Wulfen in Iacqu. collect. tom.* 1. *p.* 244. *t.* 16. *f.* 1.

Alſine foliis ſulcatis, recurvis, radicalibus linearibus congeſtis, caulinis lanceolatis. *Hall. hiſt. n.* 868.

In ſummis alpium jugis. Fl. Iulio, Auguſto. Per.

Folia linearia, recurva, facie plana, dorſo convexa, glauca: radicalia conferta, caulina oppoſita. Foliola calycina purpura ſuffuſa, ſtriata, mucronata.

faſciculata. 10. A. foliis ſubulatis, caule erecto ſtricto, floribus faſciculatis, petalis breviſſimis. *Syſt. veget. p.* 424. *Iacqu. auſtr. t.* 182.

In collibus apricis, ſaxoſis Auſtriæ, Pannoniæ. Fl. æſtate.

auſtriaca. 11. A. foliis linearibus acutis, caulibus procumbentibus ſubdichotomis, petalis emarginatis. *Syſt. veget. p.* 425. *Iacqu. auſtr. t.* 270. *Allion. pedem. tom.* 2. *p.* 112. *t.* 64. *f.* 2.

Alſine caule flaccido dichotomo, foliis linearibus acutis. *Hall. hiſt. n.* 864.

Stellaria biflora. *Wulfen in Iacqu. collect. tom.* 1. *p.* 251. *t.* 18.

In alpibus. Fl. æſtate. Per.

Folia oppoſita, linearia, acuta, glabra. Pedunculi & calyces pubeſcunt. Flores ſemper erecti. Petala calyce duplo longiora, pleraque emarginata. Capſula oblonga, conica, unilocularis. Iacqu. l. c.

liniflora. 12. A. caulibus erectis inferne ramoſis ſuffruticoſis, foliis ſubulatis, floribus geminis. *Syſt. veget. p.* 425. *Iacqu. auſtr. t.* 445.

In alpibus auſtriacis. Fl. Iulio, Auguſto. Per.

Folia ſubulata, acuminata, lævia, craſſa, erecta, dorſo convexa, facie plana. Caules plerumque uniflori. Foliola calycina lanceolata, acuta, patentiſſima, glaberrima. Petala obovata, calyce paulo longiora. Capſula ovato - oblonga. Iacqu. l. c.

CHERLERIA. Calyx pentaphyllus. Nectaria quinque, emarginata, petaloidea. Antheræ alternæ ſteriles. Capſula unilocularis, trivalvis, triſperma.

Sedoides. 1. Cherleria. *Sp. pl. p.* 608. *Iacqu. auſtr. t.* 284. *Scop. carn. tom.* 1. *p.* 312 *Hall. hiſt. n.* 859.

In ſummis alpium jugis. Fl. Iulio, Auguſto. Per.

PENTAGYNIA.

SEDUM. Calyx quinquefidus. Corolla pentapetala. Squamæ nectariferæ quinque, ad baſin germinis extrorſum inſertæ.

Telephium. 1. S. foliis planiuſculis ſerratis, corymbo folioſo, caule erecto. *Sp. pl. p.* 616. *Fl. dan. t.* 686. *Curtis lond. Scop. carn. tom.* 1. *p.* 323.

Sedum foliis confertis, ſerratis, umbellis denſis terminalibus. *Hall. hiſt. n.* 954.

Telephium auſtriacum. *Cluſ. hiſt. p. LXVI.*

L'Orpin Repriſe. Regn. bot.

In collibus, montoſis petroſis. Fl. Iulio, Auguſto. Per.

daſyphyl-lum. 2. S. foliis oppoſitis ovatis obtuſis carnoſis, caule infirmo, floribus ſparſis. *Sp. pl. p.* 618. *Iacqu. hort. tom.* 2. *p.* 71. *t.* 153. *Curtis lond. Scop. carn. tom.* 1. *p.* 324.

Sedum foliis conicis, obtuſis, glaucis, reticulatis, caule ramoſo viscido. *Hall. hiſt. n.* 961.

Sedum minus ſeptimum. *Clus. hiſt. p. LX.*

L' Orpin glauque. Bulliard. herb. t. 11.

In maceriis, muris vetuſtis Styriæ, Carnioliæ. Fl. Iunio, Iulio. Per.

reflexum. 3. S. foliis ſubulatis ſparſis baſi ſolutis: inferioribus recurvatis. *Sp. pl. p.* 618. *Fl. dan. t.* 113.

Sedum foliis ſemiteretibus, acutis, ariſtatis, caule ſupremo multifido, umbellato. *Hall. hiſt. n.* 967.

Sedum minus quintum. *Clus. hiſt. p. LX.*

In Tyroli in rupeſtribus, maceriis circa *Sterzing, Mittewalde, Vindel, Brunegen.* Fl. æſtate. Per.

Stamina duodecim. Germina ſex.

hiſpanicum. 4. S. foliis teretiusculis acutis; radicalibus faſciculatis, cyma pubescente. *Sp. pl. p.* 618. *Iacqu. auſtr. app. t.* 47.

In montanis, & alpinis Styriæ, Carinthiæ, Carnioliæ. Fl. æſtate. Per.

album. 5. S. foliis oblongis obtuſis teretiusculis ſeſſilibus patentibus, cyma ramoſa. *Sp. pl. p.* 619. *Fl. dan. t.* 66. *Curtis lond. Scop. carn. tom.* 1. *p.* 324.

Sedum caule glabro, foliis teretibus, umbellis ramoſis, floribus petiolatis. *Hall. hiſt. n.* 959.

L' Orpin a feuilles cylindriques. Bulliard herb. t. 179.

In muris, ſaxoſis apricis, & alibi. Fl. Iunio, Iulio. Per.

Oederi. 6. foliis ſemiteretibus ſparſis baſi ſolutis, caule ramoſo adſcendente, floribus cymoſis. *Retzii ſcand. p.* 103. *Fl. dan. t.* 59.

Sedum caule ramoſo proſtrato, foliis teretibus remotis, floribus ſeſſilibus. *Hall. hiſt. n.* 964.

Sedum ſaxatile. *Allion. pedem. tom.* 2. *p.* 121. *t.* 65. *f.* 6.

Sedum ſaxatile teretifolium floſculis luteis, conceptaculis ſeminum ſtellatis, & viridantibus. *Scheuch. it.* 6. *p.* 518.

In Styria in rupeſtribus alpium judenburgenſium prope *Burgerſee.* Fl. Iulio, Auguſto. Per.

acre. 7. S. foliis ſubovatis adnato-ſeſſilibus gibbis erectiusculis alternis, cyma trifida. *Sp. pl. p.* 619.

Sedum foliis conicis, confertis, caulibus ramoſis, ſupremis trifidis. *Hall. hiſt. n.* 966.

Sedum minus octavum cauſticum. *Clus. hiſt. p. LXI.*

La Vermiculaire brulante. Bulliard. herb. t. 31.

Habitat & floret cum S. albo. Per.

ſexangulare. 8. S. foliis ſubovatis adnato-ſeſſilibus gibbis erectiusculis ſexfariam imbricatis. *Sp. pl. p.* 620. *Curtis lond. Retzii obs. faſc.* 1. *p.* 18. *Scop. carn. tom.* 1. *p.* 325.

Sedum foliis teretibus, ternatis, caulibus ſimplicibus trifidis. *Hall. hiſt. n.* 965.

In Carnioliæ muris, locis ſylvaticis, aſperis. Fl. Iulio, Auguſto. Per.

Huic folia inſipida, præcedenti acria ſunt.

villoſum. 9. S. caule erecto, foliis planiusculis, pedunculisque ſubpiloſis. *Sp. pl. p.* 620. *Wulfen in Iacqu. collect. tom.* 3. *p.* 39.

Sedum hirſutum & viscidum, foliis linearibus, obtuſis, planiusculis. *Hall. hiſt. n.* 962.

Sedum minus tertium ſive paluſtre. *Clus. hiſt. p. LIX.*

In Carinthia in uliginoſis alpium fladnizenſium ad S. Iohannem, & in turfoſis declivibus *in Stangalpe. Wulfen.* Fl. Iulio. Per.

atra-tum. 10. S. caule erecto, floribus corymboſis faſtigiatis. *Syſt. veget. p.* 431. *Iacqu. auſtr. t.* 8.

Sedum caule folioſo, erecto, umbellato, humillimo, foliis teretibus. *Hall. hiſt. n.* 963.

Sedum hæmatodes. *Scop. carn. tom.* 1. *p.* 323.

In alpinis, & ſubalpinis. Fl. æſtate.

OXALIS. Calyx quinquepartitus. Petala quinque. Filamenta decem, quorum quinque longiora. Capſula quinquelocularis, longitudinaliter dehiscens. Semina ex arillo bivalvi elaſtice exilientia.

Aceto-ſella. 1. O. ſcapis unifloris, foliis ternatis, foliolis obcordatis, ſtylis æqualibus. *Iacqu. oxal. p.* 114. *t.* 80. *f.* 1. *Sp. pl. p.* 620. *Fl. dan t.* 980. *Curtis lond.*

Oxys Acetoſella. *Scop. carn. tom.* 1. *p.* 326.

L' Alleluia. Regn. bot.

In ſylvis umbroſis, humidis alpinis. Fl. Aprili, Majo, Iunio. Per.

cornicu-lata. 2. O. caule proſtrato radicante; pedunculis bifloris; ſtylis ſubæqualibus. *Iacqu. oxal. p.* 9. & 30. *t.* 5. *Syſt. veget. p.* 434.

Oxys flavo flore. *Clus. hiſt. p. CCXLVIII.*

Trifolium acetoſum corniculatum. *Bauh. pin.* 330.

In muris vetuſtis, rupeſtribus, maceriis Litoralis, nullibi copioſius quam *Lauranæ.* Fl. ab Aprili ad ſerum Autumnum.

AGROSTEMA. Calyx monophyllus, coriaceus. Petala quinque, unguiculata: Limbo obtuſo, indiviſo. Capſula unilocularis.

Githago. 1. A. hirſuta, calycibus corollam æquantibus, petalis integris nudis. *Sp. pl. p.* 624. *Curtis lond. Fl. dan. t.* 576. *Scop. carn. tom.* 1. *p.* 310.

Lychnis calycibus longiſſime caudatis. *Hall. hiſt. n.* 926.

Inter ſegetes. Fl. Iunio, Iulio.

LYCHNIS. Calyx monophyllus, oblongus, lævis. Petala quinque, unguiculata: Limbo ſæpe fiſſo. Capſula quinquelocularis.

Flos cuculi. 1. L. petalis quadrifidis, fructu ſubrotundo. *Sp. pl. p.* 625. *Fl. dan. t.* 590. *Curtis lond. Scop. carn. tom.* 1. *p.* 311.

Lychnis petalis quadrifidis. *Hall. hiſt. n.* 921.

Odontitis Plinii ſimplici flore. *Clus hiſt. p.* 292.

In humidis. Fl. Majo, Iunio. Per.

quadridentata. 2. L. petalis quadridentatis, caule dichotomo, foliis glabris recurvis. *Syſt. veget. p.* 435.

Viscago foliis teneris recurvis, petalis quadrifidis. *Hall. hiſt. n.* 918.

Silene quadrifida. *Sp. pl. p.* 602. *Iacqu. auſtr. t.* 120.

Lychnis quadrifida. *Scop. carn. tom.* 1. *p.* 307.

In alpibus. Fl. æſtate. Per.

Caules modo ſolitarii, modo cespitoſe crescentes, modo uniflori, modo multiflori. Petala nunc leviſſime quadridentata; denticulis lateralibus brevioribus, nunc tantum emarginata. Styli in aliis floribus quatuor, in aliis quinque. Iacqu. l. c.

Viscaria. 3. L. petalis ſubintegris. *Syſt. veget. p.* 435. *Scop. carn. tom.* 1. *p.* 311.

Lychnis foliis lanceolatis, floribus verticillatis. *Hall. hiſt. n.* 927.

Silene ſylveſtris quarta. *Clus. hiſt. p.* 289.

In pratis, montanis ſylvaticis. Fl. Majo, Iunio. Per.

alpina. 4. L. petalis bifidis, floribus corymboſis. *Sp. pl. p.* 626. *Fl. dan. t.* 65. *Allion. pedem. tom.* 2. *p.* 85.

Lychnis foliis linearibus, lanceolatis, floribus umbellatis capitatis. *Hall. hiſt. n.* 922. *t.* 17.

In Carnia in ſummis jugis montis *Croſtis. I. Hoſt.* Fl. Iulio, Auguſto. Per.

Calyx inflatus. Laminæ petalorum profunde ſectæ, purpureæ. Squamæ coronatrices lineares. Styli quinque flore longiores. Allion. l. c.

dioica. 5. L. floribus dioicis, *Sp. pl. p.* 626. *Fl. dan. t.* 792. *Curtis lond.*

Lychnis floribus ſexu diſtinctis. *Hall. hiſt. n.* 923.

Lychnis ſylveſtris alba ſimplex. *Bauh. pin.* 204.

Lychnis ſylveſtris ſeu aquatica purpurea ſimplex. *Bauh. pin.* 204.

Melandrium Plinii genuinum. *Clus. hiſt. p.* 294.

In pratis, ſylvaticis, ad vias, in ſubalpinis. Fl. æſtate. Per.

Variat flore albo, & purpureo.

CERASTIUM. Calyx pentaphyllus. Petala bifida. Capſula unilocularis, apice dehiſcens.

* *Capſulis oblongis.*

vulgatum. 1. C. foliis oblongo-ovatis hirſutis, caulibus diffuſis hirſutie nuda. *Curtis lond. Sibthorp. oxon. p.* 146.

Ceraſtium viscoſum. *Sp. pl. p.* 627.

In arvis, ad vias, paſcuis ſterilibus. Fl. Majo. Per.

viſco-ſum. 2. C. erectum villoſo-viſcoſum. *Curtis lond. Sibthorp. oxon. p.* 147.

Ceraſtium vulgatum. *Sp. pl. p.* 627.

In arvis, ad vias. Fl. Aprili.

ſemide-candrum. 3. C. floribus pentandris, petalis emarginatis. *Sp. pl. p.* 627. *Curtis lond.*

In paſcuis, cultis, & alibi. Fl. Majo.

arvenſe. 4. C. foliis lineari-lanceolatis obtuſis glabris, corollis calyce majoribus. *Sp. pl. p.* 627. *Curtis lond. Fl. dan. t.* 626.

Myoſotis foliis linearibus lanceolatis, petalis calyce duplo longioribus. *Hall. hiſt. n.* 889.

In arvis, ad vias. Fl. Aprili, Majo. Per.

alpinum. 5. C. foliis ovato-lanceolatis, caule diviſo, capſulis oblongis. *Sp. pl. p.* 628. *Fl. dan. t.* 6. 976. *Retzii ſcand. p.* 110.

Myoſotis foliis ovato-lanceolatis, ſubhirſutis, petalis calyce duplo longioribus. *Hall. hiſt. n.* 888.

In alpibus. Fl. æſtate. Per.

Variat foliis glabris, ſubhirſutis & hirſutiſſimis, caulibus uni- & biſloris. Retzii obs. faſc. 3. p. 32.

** *Capſulis ſubrotundis.*

aquaticum. 6. C. foliis cordatis ſeſſilibus, floribus ſolitariis, fructibus pendulis. *Sp. pl. p.* 629. *Curtis lond.*

Alſine foliis cordatis ſeſſilibus, floribus ſolitariis, fructibus pendulis. *Hall. hiſt. n.* 885.

Stellaria aquatica. *Scop. carn. tom.* 1. *p.* 319.

Ad ſepes humidas, fluvios, lacus, & alibi. Fl. æſtate. Per.

SPERGULA. **Calyx pentaphyllus. Petala quinque, integra. Capſula ovata, unilocularis, quinquevalvis.**

arvenſis. 1. S. foliis verticillatis, floribus decandris. *Sp. pl. p.* 630. *Curtis lond.*

Alſine foliis verticillatis, ſeminibus rotundis. *Hall. hiſt. n.* 873.

Stellaria arvenſis. *Scop. carn. tom.* I. *p.* 318.

Inter ſegetes, & alibi. Fl. æſtate.

pentandra. 2. S. foliis verticillatis, floribus pentandris. *Sp. pl. p.* 630.

Stellaria pentandra. *Scop. carn. tom.* I. *p.* 318.

In herbidis Tergeſti. *Scop.*

nodoſa. 3. S. foliis oppoſitis ſubulatis lævibus, caulibus ſimplicibus. *Sp. pl. p.* 630. *Curtis lond. Fl. dan. t.* 96.

Alſine foliis ſuperioribus faſciculatis, imis conjugatis. *Hall. hiſt. n.* 871.

In Auſtriæ, Styriæ arenoſis udis. Fl. æſtate. Per.

ſubulata. 4. S. foliis oppoſitis ſubſecundis lineari-ſubulatis carinatis ciliatis, floribus decandris. *N. Act. holm.* 1789. *p.* 45. *t.* 3. *f.* 3. *Retzii ſcand. p.* 111.

Spergula laricina. *Fl. dan. t.* 858.

In Carnioliæ ſubalpinis. *Zois.* Fl. æſtate. Per.

DODECANDRIA.

MONOGYNIA.

ASARUM. Calyx coriaceus, tri- ſeu quadrifidus, germini inſidens. Corolla nulla. Capſula coriacea, intra ſubſtantiam calycis.

europaeum. 1. A. foliis reniformibus obtuſis, binis. *Sp. pl. p.* 633. *Fl. dan. t.* 633. *Scop. carn. tom.* 1. *p.* 228.

Aſarum foliis reniformibus ſubhirſutis. *Hall. hiſt. n.* 1547.

Le Cabaret. Regn. bot.

In nemoribus. Fl. primo vere. Per.

PORTULACA. Calyx bifidus, apice compreſſus. Corolla pentapetala. Capſula calyce tecta, unilocularis, circumſciſſa.

oleracea. 1. P. foliis cuneiformibus, floribus ſeſſilibus. *Sp. pl. p.* 638. *Scop. carn. tom.* 1. *p.* 329.

Portulaca foliis dilatatis, lævibus, ſeſſilibus, confertis. *Hall. hiſt. n.* 968.

Portulaca anguſtifolia, ſive ſylveſtris. *Bauh. pin.* 288. *Zann. iſt. p.* 218. *t.* 262.

Le Pourpier. Regn. bot.

In cultis, ruderatis, ad vias. Fl. æſtate.

LYTHRUM. Calyx denticulis duodecim: alternis minoribus. Petala ſex, calycis inciſuris inſerta. Capſula bilocularis, polyſperma.

Salicaria. 1. L. foliis oppoſitis cordato-lanceolatis, floribus ſpicatis dodecandris. *Sp. pl. p.* 640. *Fl. dan. t.* 671. *Curtis lond. Scop. carn. tom.* 1. *p.* 329.

Salicaria foliis lanceolatis, ſubhirſutis, floribus ſpicatis. *Hall. hiſt. n.* 854.

Lyſimachia purpurea communis, major. *Cluſ. hiſt. p. LI.*

La Salicaire. Regn. bot.

Ad fluvios, rivulos, lacus, in pratis humidis. Fl. æſtate. Per.

virgatum. 2. L. foliis oppoſitis lanceolatis, panicula virgata, floribus dodecandris, ternis. *Syſt. veget. p.* 446. *Iacqu. auſtr. t.* 7.

Lythrum auſtriacum. *Iacqu. en. p.* 80.

Lyſimachia purpurea ſecunda ſive major. *Cluſ. hiſt. p. LII.*

In Auſtria in pratis inſularum danubialium, & prope *Bruck an der Leitha*, *Hainburg*. Fl. Iunio, Iulio. Per.

Hyſſopifolia. 3. L. foliis alternis, linearibus, floribus hexandris. *Sp. pl. p.* 642. *Iacqu. auſtr. t.* 133. *Scop. carn. tom.* 1. *p.* 330.

Salicaria foliis linearibus, floribus in alis ſeſſilibus. *Hall. hiſt. n.* 855.

Hyſſopifolia, ſive Gratiola minor. *Bauh. pin.* 218. *prodr.* 108.

In inundatis, paluſtribus, uliginoſis Auſtriæ, Pannoniæ, Litoralis, Forijulii. Fl. æſtate.

DIGYNIA.

AGRIMONIA. Calyx quinquedentatus, altero calyce obvallatus. Petala quinque. Calyx collo coarctatus, induratus, ſemina duo includens.

Eupatoria. 1. A. fructibus hiſpidis, foliis caulinis pinnatis: foliolis oblongo-ovatis, ſpicis elevatis, petalis calyce duplo longioribus. *Ait. Kew. tom.* 2. *p.* 129. *Sp. pl. p.* 643. *Fl. dan. t.* 588. *Curtis lond. Scop. carn. tom.* 1. *p.* 331.

Agrimonia foliis pinnatis, pinnis alterne minimis. *Hall. hiſt. n.* 991.

L' Aigremoine. Regn. bot.

In pratis, montanis ſylvaticis, ad ſepes, & alibi. Fl. Iunio, Iulio. Per.

Agrimonoides. 2. A. foliis caulinis ternatis, fructibus glabris. *Sp. pl. p.* 643. *Scop. carn. tom.* 1. *p.* 331. *Wulfen in Iacqu. collect. tom.* 2. *p.* 208.

Agrimonoides. *Col. ecphr.* 1. *p.* 145. *t.* 144.

In umbroſis Carnioliæ. Fl. Majo, Iunio. Per.

TRIGYNIA.

RESEDA. Calyx monophyllus, partitus. Petala laciniata, unico ad baſin mellifero. Capſula ore dehiſcens, & hians, unilocularis.

Luteola. 1. R. foliis lanceolatis integris, baſi utrinque unidentatis, calycibus quadrifidis. *Syſt. veget. p.* 448.

Reſeda foliis ellipticis, obtuſe lanceolatis, undulatis, calycibus quadrifidis. *Hall. hiſt. n.* 1058.

Luteola herba ſalicis folio. *Bauh. pin.* 100.

La Gaude. Regn. bot.

Ad pagos, vias, in ruderatis. Fl. æſtate.

Folia omnia integra, pleraque baſi utrinque unidentata: alia plana, alia undulata. Radicalia aggregata, caulina alternatim ſparſa. Calyx quadripartitus. Petala quinque: ſupremo mellifero, quinque- vel ſexpartito; lateralibus duobus oppoſitis trifidis, infimis duobus minimis in pleriſque floribus abſentibus. Stamina plurima. Capſula ſexangularis.

lutea. 2. R. foliis omnibus trifidis: inferioribus pinnatis. *Sp. pl. p.* 645. *Iacqu. auſtr. t.* 353. *Scop. carn. tom.* 1. *p.* 332.

Reſeda hexapetala, foliis pinnatis, undulatis, calyce ſexfido. *Hall. hiſt. n.* 1056.

La Reſeda jaune. Bulliard herb. t. 281.

In montanis, cretaceis, glareoſis, ad vias. Fl. æſtate. Per.

Folia circa medium trifida: laciniis ſublinearibus, ad oras undulatis, quandoque trifidis, rarius verſus radicem ſimplicia. Calyx hexaphyllus: foliolis ad baſin usque diviſis. Petala quatuor, ſubrotunda, crenulato-ciliata emittunt ex dorſi ſuperiore parte appendices aut integras, aut inciſas: ſuperiora duo duas cum intermedia ſubulata; inferiora unicam. Stamina plura quam duodecim. Capſula oblonga, triangularis. Iacqu. l. c.

Phyteuma. 3. R. foliis integris trilobisque, calycibus ſexpartitis maximis. *Sp. pl. p.* 645. *Iacqu. auſtr. t.* 132. *Scop. carn. tom.* 1. *p.* 332.

Reſeda foliis radicalibus integris, caulinis integris & ſemitrilobis, calycibus ſexfidis, florem ſuperantibus. *Hall. hiſt. n.* 1057.

Reſedæ affinis Phyteuma. *Bauh. pin.* 100. *prodr.* 42.

In Auſtria ad margines agrorum prope Neoſtadium, *Neunkirchen*, *Ebergäſſing*, alibique. Flumine ad latera viæ, quæ *dal Lazzaretto ai Piopi* ducit. Fl. æſtate.

Folia obverſe lanceolata, obtuſiſſima, alterna, integra, aut lobata. Calyx in maximum excreſcens conſtat foliolis ſenis, obtuſis. Petala quatuor, ſubrotunda, exigua, ſubciliata, dentato-emarginata, concava; quorum duo ſuperiora abeunt in cornu bifidum erectum, diviſum in lacinias ſubulatas, acutas: lateralia duo minora emittunt cornu integrum. Stamina circiter duodecim. Capſula ſexangularis. Iacqu. l. c.

EUPHORBIA. Calyx monophyllus, ventricoſus. Petala quatuor ſeu quinque. Stamina ſucceſſive prorumpentia. Germen pedicellatum. Capſula tricocca.

caneſcens. 1. E. dichotoma, foliis integris ſubrotundis piloſis, floribus ſolitariis axillaribus, caulibus procumbentibus. *Sp. pl. p.* 652. *Iacqu. collect. tom.* 5. *t.* 2. *f.* 3.

Tithymalus exiguus villoſus, nummulariæ folio. *Tourn. inſt.* 87.

In Litorali ad vias, in vineis, hortis. Fl. Iulio, Auguſto.

Tota planta albis pilis canescit. Folia ſemicordata, alia unico, alia utroque latere plus vel minus crenulata. pauca emarginata. Flores pedunculati, axillares, ſolitarii. Petala quatuor, crenulata, alba, interne ad baſin linea transverſali rubra notata. Capſula triquetra. Differt ab Euphorbia Chamæſyce, quæ, exceptis capſulis, quæ ſubhirſutæ ſunt, tota glabra eſt.

Peplis. 2. E. dichotoma, foliis integerrimis ſemicordatis, floribus ſolitariis axillaribus, caulibus procumbentibus. *Sp. pl. p.* 652.

Tithymalus Peplis. *Scop. carn. tom.* 1. *p.* 340.

Peplis. *Cluſ. hiſt. p. CLXXXVII.*

Habitat & floret cum præc.

Peplus. 3. E. umbella trifida: dichotoma, involucellis ovatis, foliis integerrimis obovatis, petiolatis. *Sp. pl. p.* 653. *Curtis lond.*

Tithymalus foliis rotundis, ſtipulis floralibus cordatis obtuſis, petalis argute conriculatis. *Hall. hiſt. n.* 1049.

Peplus ſeu Eſula rotunda. *Bauh. pin.* 292.

In cultis, ruderatis, ad vias. Fl. ab Auguſto ad ſerum autumnum.

Caulis erectus, plerumque baſi ramoſus. Folia obovata, integerrima. Involucrum conſtat foliolis tribus, ovatis, acutis. Involucelli foliola ſunt ſemicordata. Calyx ventricoſus. Petala quatuor, lunata. Capſula tricocca.

falcata. 4. E. umbella trifida: dichotoma, involucellis ſubcordatis mucronatis, foliis lanceolatis obtuſiuſculis. *Syſt. veget. p.* 451. *Iacqu. auſtr. t.* 121.

In cultis. Fl. Iulio, Auguſto.

Caulis ramoſus, folia ſunt obverſe lanceolata, baſi attenuata, plerumque mucronata. Similia ſunt involucri foliola. Involucella ſunt ex ovato ſubrotunda, interdum etiam magis ovata, obliqua, ſcilicet latere altero, quo oppoſita magis ab invicem divergere ſolent, latiore, hinc veluti falcato, integerrima, aut obſoletiſſime denticulata. Umbella univerſalis trifida, quandoque quadrifida. Flores ſeſſiles. Calyx campanulatus, interne hirſutus. Petala quatuor, lunata. Capſula glabra. Iacqu. l. c.

exigua. 5. E. Umbella trifida: dichotoma: involucellis lanceolatis, foliis linearibus. *Sp. pl. p.* 654. *Fl. dan. t.* 592. *Curtis lond.*

Tithymalus foliis linearibus, ſtipulis lanceolatis ariſtatis. *Hall. hiſt. n.* 1048.

In arvis, & alibi. Fl. æſtate.

Caulis ramoſus. Folia linearia, alia acutiuscula, quædam apice retuſa. Involucrum conſtat foliolis lanceolato-linearibus. Involucella ſunt ovato-oblonga, acuminata. Petala lunata. Germen ſubrotundum. Capſula tricocca, lævis.

diffuſa. 6. E. umbella quadrifida vel quinquefida: dichotoma, caule diffuſiſſimo, foliis cuneiformibus integerrimis alternis. *Syſt. veget. p.* 451. *Iacqu. miscell. tom.* 2. *p.* 311. *ic. t.* 88.

In Auſtriæ arvis. Fl. a Iulio ad Septembrim.

Tota planta eſt glabra. Caulis erectus, ſupra originem valde ramoſus. Folia, ut involucri foliola, ſunt ſubſeſſilia, cuneiformia, truncata, mucronata, integerrima. Involucellorum foliola ſunt ſeſſilia, obtuſa cum mucrone, oblonga, ſic ut altera pagina ad baſin verſus latus producatur, altera ex adverſo ſit obliterata, in ſuperioribus diviſionibus latiora, breviora, & ſæpe retuſa, quandoque bi- vel tridentata. Flores omnes ſeſſiles, & fertiles. Calyx turbinatus. Petala quatuor, acute lunata, quorum duo duplo latiora. Capſula glabra. Semina rugoſa. Iacqu. l. c.

epithymoides. 7. E. umbella quinquefida: bifida, involucellis ovatis, foliis lanceolatis obtuſis ſubtus villoſis. *Sp. pl. p. 656. Iacqu. auſtr. t. 344.*

In aſperis ſaxoſis Auſtriæ, Litoralis. Fl. Majo, Iunio. Per.

Caulis erectus. Folia oblonga, obtuſa. Umbella quinqueſida, quadriſida, aut triſida; cum floſculo centrali pedunculato erecto, & ſterili. Involucri foliola interdum emarginata, cæterum foliis caulinis ſimilia. Umbellulæ in quolibet radio ſunt ternæ cum floſculo centrali pedunculato, ex integro fertiles. Harum duæ exteriores ſunt iterum biſidæ cum floſculo medio, tertia autem, quæ interior eſt, uniflora, involucello diphyllo donata. Caulis, folia, involucra rubedine magis minusve tinguntur; alias etiam tota planta livide rubet. Calyx campanulatus, ſemiquinqueſidus: laciniis oblongis, conniventi-canaliculatis, ſuperne dentatis. Petala ſunt orbiculata, concava: in floribus ſterilibus quina, in fertilibus tria, duo, vel quatuor. Capſula hirſute muricata. Iacqu. l. c.

dulcis. 8. E. umbella quinquefida: bifida, involucellis ſubovatis, foliis lanceolatis obtuſis integerrimis. *Sp. pl. p. 656. Iacqu. auſtr. t. 213.*

In montoſis. Fl. Aprili, Majo. Per.

Radix tuberoſa. Caulis erectus. Folia oblonga, obtuſa, ſubſeſſilia, integerrima, aut tenuiſſime ſerrulata. Involucri foliola ſunt acuta.

Involucella cordata, acutiuſcula, ſemper ſerrulata. Flores minuti, alii germine carent, & præcipue centrales, alii plerique fertiles. Petala quatuor, ſubrotunda, integerrima. Germen hirſutum. Capſula verrucoſa, & villoſa. Iacqu. l. c.

carniolica. 9. E. umbella quinquefida, radiis nutantibus, involucris, involucellis & foliis lanceolatis. *Syſt. veget. p. 452. Iacqu. auſtr. app. t. 14.*

Tithymalus piloſus, *Scop. carn. tom. 1. p. 336. t. 21.*

In Carniolia in ſylvis circa Idriam, & convallibus alpium wochinenſium, tolmienſium. Fl. Aprili, Majo. Per.

Radix nodoſa. Folia obverſe lanceolata, integerrima, breviter petiolata. Umbella terminalis quinquefida fit ex radiis debilibus fereque cernuis, cum pedunculo centrali unifloro, etiam cernuo. Radii bifidi. Ex foliorum ſummorum axillis ramuli etiam exeunt breves, bifidi, biflorique. Calyx *campanulatus pilis albis pubeſcit, quinquedentatus. Petala quinque, ſubrotunda, integra, lutea. Flores ob germinis defectum plerique ſteriles.* Capſula *verrucoſa.* Iacqu. l. c.

ſaxatilis. 10. E. umbella quinquefida: bifida, involucris & involucellis cordatis, foliis oblongis glabris. *Syſt. veget. p. 452. Iacqu. auſtr. t. 345.*

In Auſtria in ſaxoſis rupeſtribus prope *Baaden*, & in monte *Gieſing* arci *Stüchſenſtein* imminenti. Fl. Majo. Per.

Tota glabra. Caules procumbentes emittunt ramulos plures adſcendentes, roſula terminatos folioſa. Folia ſunt oblonga, obtuſa, integerrima, ſeſſilia, interdum parumper emarginata, glauca, ſumma quandoque ex oblongo cordata. Flores ſunt ſeſſiles. Centralis abortat, petalis donatur quinis. Reliquis fertilibus petala ſunt quatuor, acute lunata. Calyx extus glaber, intus valde hirſutus, ore multidentato. Capſula glabra, trigono-ovata. Iacqu. l. c.

Paralias. 11. E. umbella ſubquinquefida: bifida, involucellis cordato-reniformibus, foliis ſurſum imbricatis. *Sp. pl. p.* 657. *Iacqu. hort. tom.* 2. *p.* 88. *t.* 188.

Tithymalus maritimus. *Bauh. pin.* 292.

Tithymalus Paralias. *Bauh. hiſt.* 3. *p.* 674. *Zann. iſt. p.* 259. *t.* 40.

In arenoſis Litoralis ad mare. Fl. Iunio. Per.

Tota glabra. Radix lignoſa. Caulis & rami veſtiuntur foliis erectis, lineari-oblongis, ſubacutis, craſſiuſculis, cum exiguo mucrone, integerrimis, glaucis, imbricatis. Umbella quadrifida, vel quinquefida. Involucri foliola ſunt ſubovata; involucellorum cordato-ſubrotunda. Calyx ſubrotundus, quadriſulcatus, quadridentatus. Petala quatuor, ſubpeltata, bi- vel quadridentata. Capſula rugoſa. Flores ſteriles ſine ordine fertilibus intermixti. Iacqu. l. c.

ſegetalis. 12. E. umbella quinquefida: dichotoma, involucellis cordatis, acutis: ſuperioribus latioribus. *Syſt. veget. p.* 452. *Iacqu. auſtr. t.* 450.

In Auſtria inter ſegetes. Fl. æſtate.

Tota planta glaberrima. Folia linearia, acuta, integerrima. Flores omnes fertiles & ſeſſiles. Calyx glaber. Petala quatuor, acute lunata. Germen verrucoſulum. Iacqu. l. c.

helioſcopia. 13. E. umbella quinquefida: trifida: dichotoma, involucellis obovatis, foliis cuneiformibus ſerratis. *Sp. pl. p.* 658. *Curtis lond. Fl. dan. t.* 725.

Tithymalus foliis petiolatis, ſubrotundis, ſerratis, ſtipulis rotundis, ſerratis. *Hall. hiſt. n.* 1050.

Tithymalus helioſcopius. *Scop. carn. tom.* 1. *p.* 337.

In cultis, ad vias. Fl. a Iunio ad ſerum autumnum.

Caulis vel ſimplex, vel baſi ramoſus. Folia obovata, glabra, ſerrata. Hisce ſimilia ſunt involucri, involucellorumque foliola. Petala rotundata. Capſula ſubrotunda, glabra.

verrucoſa. 14. E. umbella quinquefida: ſubtrifida: bifida, involucellis ovatis, foliis lanceolatis ſerrulatis villoſis, capſulis verrucoſis. *Sp. pl. p.* 658.

Tithymalus foliis ellipticis ſerratis, ſtipulis umbellaribus quinis, floralibus obtuſe quadrangulis, fructibus undique exaſperatis. *Hall. hiſt. n.* 1052.

Tithymalus myrſinites, fructu verrucæ ſimili. *Bauh. pin.* 291.

In Auſtriæ montanis ſylvaticis, & in inſulis danubialibus. Fl. Majo, Iunio. Per.

Caules decumbentes vel erecti. Folia ſeſſilia, lanceolata, aculeato-ſerrulata, juniora villoſa, dein calveſcentia. Involucri foliola, & involucella ſunt breviora, ac latiora, cæterum foliis caulinis ſimilia. Petala rotunda. Capſulæ muricatæ.

platyphylla. 15. E. umbella quinquefida: trifida: dichotoma, involucellis carina piloſis, foliis ſerratis lanceolatis, capſulis verrucoſis. *Sp. pl. p.* 660. *Iacqu. auſtr. t.* 376.

In cultis, & incultis Auſtriæ. In ſterilibus, aſperis Carnioliæ, prope *Prewald*, *Seſſana*. Fl. æſtate.

Folia ſubſeſſilia, oblonga, infima obtuſa, ſuperiora magis acuta, facie glabra, dorſo & ad oras hirſutula, inferiora integerrima, ſuperiora tenuiſſime ſerrulata. Umbella habet radios a tribus ad quinque, trifidos, vel quadrifidos, dein tertia diviſione bifidos, vel trifidos. Foliola involucri univerſalis ſunt caulinis ſimilia; ſubſequarum vero umbellarum magis acuta, ſumma cordata ſine baſeos emarginatione. Flos centralis pedunculatus, pentapetalus, plerumque abortiens. Reliqui flores tetrapetali, & omnes fertiles ſunt. Calyx campanulatus, hirſutiſſimus. Petala rotunda. Germen ovatum & hirſutum. Iacqu. l. c.

Eſulæ. 16. E. umbella multifida: bifida, involucellis ſubcordatis, petalis ſubbicornibus, ramis ſterilibus, foliis uniformibus. *Sp. pl. p.* 660.

In Auſtria, Pannonia ad vias, agrorum vinearumque margines, in herbidis. Fl. æſtate. Per.

Folia lineari-lanceolata, glabra, ſeſſilia, obtuſiuſcula, mucronata. Involucri foliola ovata. Involucella cordiformia. Petala lunata. Fructus glaber.

gerardiana. 17. E. umbella multifida: dichotoma, involucellis ſubrotundis, petalis integerrimis, ramis nullis, foliis uniformibus. *Syſt. veget. p.* 454. *Iacqu. auſtr. t.* 436.

In collibus agri vindobonenſis, ad danubii ripas. Fl. Iunio, Iulio. Per.

Folia ſeſſilia, ſublinearia. Involucri foliola foliis caulinis ſimilia, ſed paulo breviora, & latiora. Involucella late ſubrotunda, ſeſſilia. Umbellæ radii ſunt bis vel ter bifidi. Flores omnes ſeſſiles, fertiles. Calyx glaber. Petala quatuor, ſubrotunda, integerrima. Capſula glabra. Iacqu. l. c.

Cypariſſias. 18. E. umbella multifida: dichotoma, involucellis ſubcordatis, ramis ſterilibus foliis ſetaceis, caulinis lanceolatis. *Sp. pl. p.* 661. *Iacqu. auſtr. t.* 435.

La petite Eſule. Regn. bot.

In pratis, locis arenoſis, collibus, montanis ſylvaticis. Fl. Aprili, Majo. Per.

Folia caulina linearia: ramea ſubſetacea. Involucri foliola foliis caulinis æmula. Involucella cordata, integra. Flos medius in diviſione radii principe infima eſt hermaphroditus, tamen ſterilis. Reliqui ſunt feminei & fertiles. Femineis eſt calyx campanulatus, quadridentatus: denticulis ad apicem bifidis. Petala quatuor. Stamina nulla. Germen ovatum. Capſula triquetra, muricato-rugoſa. Hermaphroditis eſt calyx quinquedentatus.

Corolla pentapetala. Stamina circiter octo. Occurrunt & in hac ſpecie, & in Eſula caules ſimplices, ſteriles, roſula terminati folioſa.

Myrſinites. 19. E. umbella ſuboctofida: bifida, involucellis ſubovatis, foliis ſpatulatis patentibus carnoſis mucronatis, margine ſcabris. *Sp. pl. p.* 661.

Tithymalus Myrſinites legitimus. *Clus. hiſt. p. CLXXXIX.*

In Litorali ad mare. Fl. Iunio, Iulio. Per.

Caules deflexi. Folia ſpathulata, margine aculeato-ſerrulata, mucronata. Umbella ſit ex radiis a ſeptem ad novem. Involucri foliola ovata. Involucella ſubcordata. Flores maſculi hermaphroditis intermixti. Calyx ſerratus. Petala bicornia. Capſulæ glabræ.

paluſtris. 20. E. umbella multifida: ſubtrifida: bifida, involucellis ovatis, foliis lanceolatis, ramis ſterilibus. *Sp. pl. p.* 662. *Fl. dan. t.* 866. *Iacqu. miscell. tom.* 2. *p.* 314.

In paluſtribus Auſtriæ, Pannoniæ, & in editiſſimo Auſtriæ monte *Oetſcherberg*. Fl. Majo, Iunio. Per.

Folia lanceolata, ſeſſilia, ad lentem ſuperne argute ſerrata, cæterum integerrima, piloſa. Umbellulæ ſunt quadrifidæ, vel bifidæ. Flores principes centrales in omnibus umbellis pedunculati, maſculi, pentapetali: reliqui hermaphroditi, tetrapetali. Calyx campanulatus, pilis albis inſperſus. Petala integra, transverſe ſubovata, obtuſiſſima. Capſula verrucoſa. Iacqu. l. c.

amygdaloides. 21. E. umbella multifida: dichotoma, involucellis perfoliatis orbiculatis, foliis obtuſis. *Sp. pl. p.* 662.

Euphorbia ſylvatica. *Iacqu. auſtr. t.* 375. *auct. Cl. Sibthorp.*

In montanis ſylvaticis, alpium convallibus. Fl. primo vere. Per.

Folia ſunt lanceolata, vel elliptica, ſuperne latiora, in brevem petiolum attenuata, obtuſa, integerrima. Involucri foliola ex ſubrotundo obverſe ovata, obtuſa. Involucella monophylla: inferiora plana; ſumma concava pelviformia, utrinque inciſo-emarginata, ut ex binis foliolis late triangularibus, obtuſisque videantur in unum conflata. Flores ſeſſiles poſiti inter radios plerique pentapetali ſteriles & decidui. Terminales fere ſoli fertiles, tetrapetali. Calyx ſubrotundus. Petala lunata. Germen glabrum. Iacqu. l. c.

nicenſis. 22. E. umbella multifida: dichotoma, involucellis ſubrotundis cordatis mucronatis, foliis ſubcuneiformibus integerrimis glabris. *Allion. pedem. tom. 1. p. 285. t. 69. f. 1. Iacqu. collect. tom. 2. p. 324. ic. t. 485.*

In Auſtria circa *Lanzendorf, Himberg, Ebergäſſing, Bruck an der Leitha*, vicinisque locis. Fl. Iunio. Per.

Tota glabra. Caulis *teres, purpuraſcens, adſcendens, vel erectus, ſimplex, aut e ſummorum foliorum alis ramulum unum alterumve aphyllum cum umbella bifida emmittens. Folia ſeſſilia, ſubcuneiformia, integerrima, acutula cum mucrone reflexo, firma, craſſula, glauca. Umbella conſtat radiis a quinque ad decem, bis dichotomis. Involucri foliola foliis ſimillima, ſed breviora & latiora. Involucella ſunt cordato-ſubrotunda, ſeſſilia, integerrima cum exiguo mucrone reflexo, ex flavo-virentia. Flores glabri, calyce turbinato. Primores umbellæ pentapetali, maſculi, ſteriles. Tales etiam in umbellis majoribus magisque diviſis flores primæ dichotomiæ. Reliqui tetrapetali, hermaphroditi, fertiles, Petala breviter bicornia, nonnunquam ſine ullis cornubus integerrima, lutea. Capſula glabra.* Iacqu. l. c.

ſalicifolia. 23. E. umbella multifida: dichotoma, involucellis reniformi-cordatis, foliis lanceolatis villoſis.

In Pannonia inter *Bruck an der Leitha* & *Neuſidelſee* ad margines ſylvarum, vias, ſepes. Fl. Majo. Per.

Caulis erectus, ſtriatus. Folia ſeſſilia, alternatim ſparſa, lanceolata, aut ovato-lanceolata, utrinque villoſa, obtuſa, glauca. Involuceri foliola foliis æmula ſunt, ſed breviora. Involucella ſunt reniformi-cordata, apice compreſſa, &, uti tota pars ſuperior plantæ, flaveſcentia. Calyces ventricoſi, villoſi. Petala in omnibus floribus quatuor, bicornia, glabra. Capſulæ glabræ, rugoſulæ.

DODECAGYNIA.

SEMPERVIVUM. Calyx ſex- ad duodecimpartitus. Petala ſex ad duodecim. Stamina & capſulæ tot, quot petala.

tectorum. 1. S. foliis ciliatis, propaginibus patentibus. *Sp. pl. p.* 664. *Scop. carn. tom.* 1. *p.* 325. *Curtis lond. Fl. dan. t.* 601.

La grande Ioubarbe. Regn. bot.

In tectis, muris vetuſtis. Fl. Iunio, Iulio. Per.

globiferum. 2. S. foliis ciliatis, propaginibus globoſis. *Sp. pl. p.* 665. *Iacqu. auſtr. app. t.* 40.

Sedum roſulis hirſutis, ciliatis, petalis ſubulato-lanceolatis duodenis. *Hall. hiſt. n.* 950.

In Carinthia in rupeſtribus alpium fladnizenſium, reichenavienſium; in Tyroli in alpibus leontinis. *Wulfen.* Fl. Iulio, Auguſto. Per.

arachnoideum. 3. S. foliis pilis intertextis, propaginibus globoſis. *Sp. pl. p.* 665. *Iacqu. auſtr. app. t.* 42.

Sedum roſulis araneoſis calveſcentibus, petalis connatis nervoſis. *Hall. hiſt. n.* 952.

In alpibus carinthiacis. In Tyroli in montanis rupeſtribus, ad latera viarum circa *Sterzing*, *Mittewalde*, *Millerwald*, alibique. Fl. æſtate. Per.

hirtum. 4. S. foliis caule petalorumque apicibus hirtis. *Sp. pl.* p. 665. *Iacqu. auſtr.* t. 12.

Cotyledon altera ſecunda. *Clus. hiſt.* p. LXIII.

In collibus Auſtriæ aſperis. Fl. Iulio, Auguſto, Septembri. Per.

montanum. 5. S. foliis integerrimis, propaginibus patulis. *Syſt. veget.* p. 456. *Iacqu. auſtr. app.* t. 41.

Sedum foliis ſemipatulis hirſutis, petalis novenis ſubhirſutis. *Hall. hiſt.* n. 951.

Cotyledon altera tertia. *Clus. hiſt.* p. LXIV.

In alpibus ſtyriacis, carinthiacis, tyrolenſibus. Fl. Iulio, Auguſto. Per.

ICOSANDRIA.

MONOGYNIA.

CACTUS. Calyx monophyllus, ſuperus, imbricatus. Corolla multiplex. Bacca unilocularis, polyſperma.

Opuntia. 1. C. articulato - prolifer laxus, articulis ovatis, ſpinis ſetaceis. *Sp. pl. p.* 669.

Opuntia foliis ovatis, compreſſis, ſpinis ſetaceis. *Hall. hiſt. n.* 1099.

Opuntia vulgo herbariorum. *Bauh. hiſt.* 1. *p.* 154. *Miller ic.* 191.

Flumine in *Belvedere* in maceriis vineas circumdantibus. Per.

MYRTUS. Calyx quadri- ſeu quinquefidus, ſuperus. Petala quatuor ſeu quinque. Bacca bi- ſeu trilocularis. Semina pauca, diſſepimentis affixa.

communis. 1. M. floribus ſolitariis: involucro diphyllo. *Sp. pl. p.* 673. *Scop. carn. tom.* 1. *p.* 341.

Le Myrthe. Regn. bot.

In rupeſtribus Litoralis inter *Moſchienizze* & *Fianona*, Duini. Copioſiſſime in inſula Auxerio. Fl. Iunio. Fr.

PUNICA. Calyx quinquefidus, ſuperus. Petala quinque, calyci inſerta. Pomum ſepto transverſo in duas concamerationes diviſum, quarum ſuperior ſubnovemlocularis, inferior ſubtrilocularis. Diſſepimentis membranaceis. Semina ſucculenta plurima.

Granatum. 1. P. foliis lanceolatis, caule arboreo. *Sp. pl. p.* 676. *Miller illuſtr. ic. Scop. carn. tom.* 1. *p.* 341.

Punica ſpinoſa, foliis nitentibus, ellipticis, integerrimis, floribus ſeſſilibus. *Hall. hiſt. n.* 1098.

In Litorali circa Duinum, Tergeſtum in hortis, & vineis neglectis, ad ſepes. Flumine in ſimilibus locis, & ad pedem montis *Vogliack* dicti ad latera viæ, que *ai molini* ducit. Fl. Iunio.

AMYGDALUS. Calyx quinquefidus, inferus. Petala quinque. Drupa nuce poris perforata.

Perſica. 1. A. foliorum ſerraturis omnibus acutis, floribus ſeſſilibus ſolitariis. *Sp. pl. p.* 676.

Le Pecher. Regn. bot.

Flumine ad vineas, aliisque aſperis locis. Fl. primo vere. Arb.

communis. 2. A. foliorum ſerraturis infimis glanduloſis, floribus ſeſſilibus, geminis. *Sp. pl. p.* 677.

Amygdalus foliis glabris, ovato-lanceolatis, ſerraturis imis dentibus & petiolis glanduloſis. *Hall. hiſt. n.* 1078.

L' Amandier. Regn. bot.

Occurrit per Forojulium, Iſtriam, Litorale. Inveni quoque in Pannonia in collibus inter *Bruck an der Leitha* & *Neuſidelſee* non procul a pago *Iojos*. Fl. Februario, Martio, Aprili. Arb.

nana. 3. A. foliis baſi attenuatis. *Sp. pl. p.* 677. *Curtis magaz.* 161. *Pallas roſſic. t.* 6.

In Pannonia in ſylvis, ad agrorum margines circa *Zorndorf*, pago ſesquihora circiter a *Bruck an der Leitha* diſſito. *Egregius hortulanus Lübeck.* Fl. primo vere. Fr.

PRUNUS. Calyx quinquefidus, inferus. Petala quinque. Drupæ nux ſuturis prominulis.

Padus. 1. P. floribus racemoſis, foliis deciduis baſi ſubtus biglanduloſis. *Sp. pl. p.* 677. *Fl. dan. t.* 205. *Scop. carn. tom.* 1. *p.* 343.

Padus foliis ovato-lanceolatis ſerratis. *Hall. hiſt. n.* 1086.

In ſylvis, ad ſepes. Fl. Aprili, Majo. Arb.

Mahaleb. 2. P. floribus corymboſis terminalibus, foliis ovatis. *Sp. pl. p.* 678. *Iacqu. auſtr. t.* 227. *Scop. carn. tom.* 1. *p.* 343.

Ceraſus foliis ſubrotundis ſerratis, pedunculis multifloris. *Hall. hiſt. n.* 1084.

Mahaleb. *Clus. hiſt. p.* 64.

In ſaxoſis, rupeſtribus Litoralis, Forijulii. In ſimilibus locis in Auſtria prope *Baaden*, *Medling*, *Briel.* Fl. Aprili, Majo. Arb.

Ceraſus. 3. P. umbellis ſubpedunculatis, foliis ovato-lanceolatis glabris conduplicatis. *Syſt. veget. p.* 463.

Ceraſus foliis glabris, ſerratis, ovato-lanceolatis, mucrone producto. *Hall. hiſt. n.* 1083.

Prunus Chamæceraſus. *Iacqu. collect. tom.* 1. *p.* 133. *ic. t.* 90.

Chamæceraſus. *Clus. hiſt. p.* 64.

In Auſtria, Pannonia in apricis aſperis, dumetis, ad margines ſylvarum, ad vineas. Fl. Majo. Arb.

Variat pro diverſa ſoli natura, ac expoſitione a pedali ad humanam magnitudinem. In hortis in arborem excreſcit.

avium. 4. P. umbellis ſeſſilibus, foliis ovato-lanceolatis ſubtus pubescentibus conduplicatis, petiolis biglandulofis. *Sibthorp. oxon. p.* 154. *Syſt. veget. p.* 463.

Ceraſus foliis ovato-lanceolatis, ſerratis, inferne ſubhirſutis, mucrone producto. *Hall. hiſt. n.* 1082.

Prunus Ceraſus. *Scop. carn. tom.* 1. *p.* 343.

Le Meriſier. Regn. bot.

In ſylvis cum aliis arboribus magnitudine certans. In Auſtria, Pannonia ad ſepes quoque, alibique in apricis ſiccis vel fruticis, vel demiſſæ arbusculæ modo crescens. Fl. Aprili, Majo.

domeſtica. 5. P. pedunculis ſubſolitariis, foliis lanceolato-ovatis convolutis, ramis muticis. *Sp. pl. p.* 680.

Prunus foliis ſerratis, hirſutis, ovato-lanceolatis, floribus longe petiolatis. *Hall. hiſt. n.* 1079.

Le Prunier. Regn. bot.

Ad pagos, vineas. Fl. primo vere. Arb.

inſititia. 6. P. pedunculis geminis, foliis ovatis ſubvilloſis convolutis, ramis ſpinescentibus. *Sp. pl. p.* 680. *Hall. hiſt. n.* 1081.

Habitat & floret cum præc. Fr.

ſpinoſa. 7. P. pedunculis ſolitariis, foliis lanceolatis glabris, ramis ſpinoſis. *Sp. pl. p.* 681. *Fl. dan. t.* 926. *Scop. carn. tom.* 1. *p.* 342.

Prunus ſpinoſa, foliis glabris, ſerratis, ovato-lanceolatis, floribus breviter petiolatis. *Hall. hiſt. n.* 1080.

Le Prunelier. Regn. bot.

In collibus apricis, ſylvis, ad ſepes, vias. Fl. primo vere. Fr.

DIGYNIA.

CRATÆGUS. Calyx quinquefidus. Petala quinque. Bacca infera, biſperma.

Aria. 1. C. foliis ovatis inciſis ſerratis ſubtus tomentoſis. *Syſt. veget. p.* 464. *Fl. dan. t.* 302.

Mespilus foliis ovatis, ſerratis, ſubtus tomentoſis. *Hall. hiſt. n.* 1089.

Mespilus Aria. *Scop. carn. tom.* 1. *p.* 345.

Aria cognominata. *Clus. hiſt. p.* 9.

In ſylvis montanis, & ſubalpinis, ascendit quoque ad alpes. Fl. Majo, Iunio. Arb.

In alpinis creſcens folia habet magis diviſa, dorſo minus tomentoſa. In demiſſioribus contra.

torminalis. 2. C. foliis cordatis ſeptangulis: lobis infimis divaricatis. *Sp. pl. p.* 681. *Iacqu. auſtr. t.* 443. *Fl. dan. t.* 793.

Mespilus foliis ſerratis ſeptilobis primis divergentibus. *Hall. hiſt. n.* 1088.

In Pannoniæ, Croatiæ, Litoralis, Iſtriæ, Auſtriæ ſylvis. Fl. Majo. Arb.

Folia ſunt ſeptemlobata: lobis ſuperioribus apicem verſus vergentibus; duobus infimis divaricatis. At in agro vindobonenſi in ſylvula, Laacherwald *dicta, vidi arbores, in quarum foliis lobi duo ultimi a reliqua folii ſuſtantia perfecte ſeparati ſunt, ita ut folia ſint quinquloba, alata.*

Oxyacantha. 3. C. foliis obtuſis ſubtrifidis ſerratis. *Syſt. veget. p.* 465. *Iacqu. auſtr. t.* 292. *f.* 2.

In Auſtriæ, Pannoniæ collibus, montoſis ſylvaticis. Fl. primo vere. Fr.

monogyna. 4. C. foliis ſubtrifidis acutis, floribus monogynis. *Syſt. veget. p.* 465. *Iacqu. auſtr. t.* 292. *f.* 1.

Mespilus Oxyacantha. *Scop. carn. tom.* 1. *p.* 344.

Ad ſepes, vias, in ſylvis. Fl. primo vere. Fr.

TRIGYNIA.

SORBUS. Calyx quinquefidus. Petala quinque. Bacca infera, triſperma.

aucupa-ria. 1. S. foliis pinnatis utrinque glabris. *Sp. pl. p.* 683. *Miller illuſtr. ic. Hall. hiſt. n.* 1091.

Mespilus aucuparia. *Scop. carn. tom.* 1. *p.* 346.

Aucuparia Sorbus. *Clus. hiſt. p.* 9.

In ſylvis montanis, & ſubalpinis. Fl. Majo, Iunio. Arb.

domeſti-ca. 2. S. foliis pinnatis ſubtus villoſis. *Sp. pl. p.* 684. *Iacqu. auſtr. t.* 447.

Mespilus foliis pinnatis, ſubtus villoſis. *Hall. hiſt. n.* 1092.

Sorbus legitima. *Clus. hiſt. p.* 9.

Le Sorbier commum. Regn. bot.

Ad pagos, vineas. Fl. Majo, Iunio. Arb.

PENTAGYNIA.

MESPILUS. Calyx quinquefidus. Petala quinque. Bacca infera, pentaſperma.

germa-nica. 1. M. ſpinoſa, foliis lanceolatis ſubtus tomentoſis, floribus ſeſſilibus ſolitariis. *Sp. pl. p.* 684. *Pallas roſſic. t.* 13. *f.* 1.

Mespilus foliis elliptico - lanceolatis, ſerratis, calycibus longiſſimis periſtentibus. *Hall. hiſt. n.* 1094.

Mespilus Pyracantha. *Socp. carn. tom.* 1. *p.* 347.

Le Neſſlier. Regn. bot.

In Iſtria, Litorali, Carniolia calidiore ſpontanea in ſylvaticis, aſperis, ad ſepes occurrit. In Auſtria reliquisque provinciis in cultis invenitur. Fl. Majo. Arb.

In ſpontanea rami principes ſpinis validis armantur; folia, flores, fructus multo minores ſunt. In culta deſiderantur ſpinæ; folia, flores, fructus multo majores ſunt quam in ſpontanea. Cl. Linnæus differentiam ſpecificam ad cultam arborem fecit. Hinc factum, quod Cl. Scopoli M. germanicam, in qua ſpinas detexit, ſub nomine M. Pyracanthæ nobis propoſuerit.

Amelanchier. 2. M. inermis, foliis ovalibus ſerratis ſubtus hirſutis. *Syſt. veget. p.* 466. *Jacqu. auſtr. t.* 300. *Scop. carn. tom.* 1. *p.* 347.

Pyrus foliis ovatis, ſerratis, ſubtus tomentoſis, calveſcentibus. *Hall. hiſt. n.* 1095.

Vitis idæa tertia. *Clus. hiſt. p.* 61.

In collibus, montanis ſylvaticis Auſtriæ, Carnioliæ calidioris, Forijulii, Litoralis. Fl. Aprili, Majo. Fr.

Folia juniora marginibus conniventia, dorſo hirſuta. Adultiora paulatim ſeſe explicant, ac maximam pubis partem abjiciunt. Aetate provecta ſiunt penitus plana, glaberrima.

Chamæ-Mespilus. 3. M. inermis, foliis ovalibus acute ſerratis glabris, floribus corymboſo-capitatis. *Syſt. veget. p.* 466. *Scop. carn. tom.* 1. *p.* 345.

Mespilus foliis ovatis ſerratis glabris. *Hall. hiſt. n.* 1090.

Cratægus Chamæ-Mespilus. *Jacqu. auſtr. t.* 231.

Cotoneaſter forte Gesneri. *Clus. hiſt. p.* 62.

In Auſtriæ, Styriæ, Carnioliæ alpinis, & ſubalpinis. Fl. Iunio. Fr.

Cotoneaſter. 4. M. inermis, foliis ovatis integerrimis: ſubtus tomentoſis. *Syſt. veget. p.* 466. *Fl. dan. t.* 112. *Pallas roſſic. t.* 15. *Scop. carn. tom.* 1. *p.* 346. *Hall. hiſt. n.* 1093.

Chamæ-Mespilus Gesneri. *Clus. hiſt. p.* 60.

In collibus, montoſis ſylvaticis, occurrit quoque in alpinis, & ſubalpinis. Fl. Majo, Iunio. Fr.

In locis apricis, ſiccioribus creſcens humilis eſt frutex, folia habens parva, ſuborbiculata. Inter alios frutices, in ſubumbroſis, ſolo feliciori habitans, in elatum fruticem excreſcit, veſtiturque foliis amplioribus, ovatis. Variat quoque, pro diverſa ſoli nutura, pedunculis ſimplicibus, & ramoſis; ſtylis à tribus ad quinque; & cum his ſeminum numero.

PYRUS. Calyx quinquefidus. Petala quinque. Pomum inferum, quinqueloculare, polyſpermum.

communis. 1. P. foliis ſerratis glabris, floribus corymboſis. *Syſt. veget. p.* 466. *Scop. carn. tom.* 1. *p.* 348.

Pyrus foliis ovato-lanceolatis, ſerratis, glabris. *Hall. hiſt n.* 1096.

In ſylvis. Fl. Aprili, Majo. Arb.

Pyrus nivalis Iacqu. auſtr. t. 107. *colitur in Auſtria, Styria, Iſtria, propagaturque omnibus iis methodis, quibus aliæ hortenſes varietates propagantur. E ſeminibus terræ concreditis oritur P. communis ramis ſpineſcentibus, nempe Pyraſter horticolarum.*

Malus. 2. P. foliis ſerratis, umbellis ſeſſilibus. *Sp. pl. p.* 686. *Miller illuſtr. ic. Scop. carn. tom.* 1. *p.* 348.

Pyrus foliis ovatis acuminatis, ſubtus hirſutis, petiolis florigeris breviſſimis. *Hall. hiſt. n.* 1097.

Le Pomier. Regn. bot.

In ſylvis, ad ſepes. Fl. Majo. Arb.

Cydonia. 3. P. foliis integerrimis, floribus ſolitariis. *Sp. pl. p.* 687. *Iacqu. auſtr. t.* 342.

Le Coignaſſier. Regn. bot.

In Auſtria, Iſtria ad vineas, vias, ſepes. Fl. Majo. Arb.

SPIRÆA. **Calyx quinquefidus. Petala quinque. Piſtilla quinque, vel plura: Capſulæ bivalves. Semina ſuturæ internæ affixa.**

ſalicifolia. 1. S. foliis lanceolatis obtuſis ſerratis nudis, floribus duplicato - racemoſis. *Sp. pl. p.* 700. *Pallas roſſic. t.* 21. 22.

Spiræa Theophraſti forte. *Clus. hiſt. p.* 84.

Frutex ſpicatus, foliis ſerratis ſalignis. *Bauh. pin.* 475.

In Carnia prope *Argons. I. Hoſt.* In Carniolia in agro labacenſi. *Zois.* In Carinthia inter *Gurk* & *Weitensfeld*, tum in valle *Gledniz. Wulfen.* Ego in Styria ſuperiore in agro knittelfeldenſi ad ripas fluvii *Ingering.* Fl. a Iunio ad Septembrim. Fr.

chamædrifolia. 2. S. foliis ovatis inciſo - ſerratis glabris, umbellis pedunculatis. *Sp. pl. p.* 701. *Iacqu. hort. tom.* 2. *p.* 66. *t.* 140.

Spiræa ulmifolia. *Scop. carn. tom.* 1. *p.* 349. *t.* 22.

In Carniolia *in der Wochein* ad originem fluminis *Feiſtriz*; in rupeſtribus circa Idriam. In ſimilibus locis in Forojulio, Carnia. Fl. primo vere. Fr.

Folia alterna, breviter petiolata, inciſo ſerrata, dorſo copioſiori, facie rariori villo inſperſa, dein calveſcentia: alia ovata, quædam lanceolata, multa ex lanceolato ovata.

Aruncus. 3. S. foliis ſupradecompoſitis, ſpicis paniculatis, floribus dioicis. *Sp. pl. p.* 702. *Pallas roſſic. t.* 36. *Scop. carn. tom.* 1. *p.* 350.

Filipendula dioica, foliis duplicato - pinnatis, pinnis quinquefoliis, trifoliis & ſimplicibus. *Hall. hiſt. n.* 1134.

Barba capræ floribus oblongis. *Bauh. pin.* 163.

In montanis, & ſubalpinis. Fl. Iulio, Auguſto. Per.

Filipendula. 4. S. foliis interrupte pinnatis: foliolis lineari-lanceolatis interrupte ſerratis glaberrimis, floribus cymoſis. *Syſt. veget. p.* 472. *Fl. dan. t.* 635. *Scop. carn. tom.* 1. *p.* 350.

Filipendula foliis pinnatis, minimis intermixtis, extrema trilobata. *Hall. hiſt. n.* 1136.

Filipendula vulgaris. *Clus. hiſt. p.* CCXI.

La Filipendule. Regn. bot.

In pratis. Fl. æſtate. Per.

Ulmaria. 5. S. foliis interrupte pinnatis: foliolis ovatis biſerratis ſubtus canis, floribus cymoſis. *Syſt. veget. p.* 472. *Curtis lond. Fl. dan. t.* 547. *Scop. carn. tom.* 1. *p.* 351.

Filipendula pinnis acute ſerratis, minimis intermixtis, extrema trilobata maxima. *Hall. hiſt. n.* 1135.

Ulmaria. *Clus. hiſt. p.* CXCVIII.

La Reine des Prés. Regn. bot.

In pratis udis, ad margines ſylvarum demiſſarum, ad fluvios. Fl. Iunio. Per.

POLYGYNIA.

ROSA. Calyx urceolatus, quinquefidus, carnoſus, collo coarctatus. Petala quinque calycis collo inſerta. Semina plurima, hiſpida, calycis interiori lateri affixa.

rubiginoſa. 1. R. germinibus ovatis, pedunculis hiſpidis, petiolis cauleque aculeatis: aculeis recurvis, foliolis ovatis ſubtus glanduloſo-piloſis. *Ait. Kew. tom.* 2. *p.* 206. *Syſt. veget. p.* 473. *Iacqu. auſtr. t.* 50.

Roſa ſpinis aduncis, foliis ſubtus rubiginoſis. *Hall. hiſt. n.* 1103.

Roſa ſuavifolia. *Fl. dan. t.* 870.

In Auſtriæ collibus, ad margines agrorum, copioſiſſime in inſulis danubialibus. Fl. Iunio, Iulio. Fr.

pumila. 2. R. germinibus ovatis petiolis pedunculisque hiſpidis, caule ſuperne aculeatiſſimo. *Syſt. veget. p.* 474. *Iacqu. auſtr. t.* 198.

Roſa pumila. *Cluſ. hiſt. p.* 117.

In Auſtriæ, Pannoniæ collibus, montoſis ſylvaticis. Fl. Majo, Iunio. Fr.

alpina. 3. R. germinibus ovatis glabris, pedunculis petiolisque hiſpidis, caule inermi. *Syſt. veget. p.* 474. *Iacqu. auſtr. t.* 279.

Roſa inermis, foliis ſeptenis glabris, calycis ſegmentis indiviſis. *Hall. hiſt. n.* 1107.

In Auſtriæ, Styriæ ſubalpinis, & alpinis. Fl. Iunio, Iulio. Fr.

canina. 4. R. germinibus ovatis pedunculisque glabris, caule petiolisque aculeatis. *Syſt. veget. p.* 474. *Curtis lond.*

L' Eglantier, ou le Roſier ſauvage. Regn. bot.

In collibus, montoſis ſylvaticis, ad ſepes, vias. Fl. Iunio. Fr.

collina. 5. R. germinibus ovatis ſubglabris, pedunculis petiolisque glanduloſo - hirſutis, caule aculeato. *Syſt. veget. p.* 474. *Iacqu. auſtr. t.* 197.

In collibus agri vindobonenſis. Fl. Majo. Fr.

ſpinoſiſſima. 6. R. germinibus globoſis glabris, pedunculis hiſpidis, caule petiolisque aculeatiſſimis. *Syſt. veget. p.* 473.

Roſa campeſtris odorato flore. *Cluſ. hiſt. p.* 116.

In Auſtriæ, Pannoniæ ſterilibus. Fl. Majo, dein autumno. Fr.

RUBUS. Calyx quinquefidus. Petala quinque. Bacca compoſita, acinis monoſpermis, in capitulum collectis.

idæus. 1. R. foliis quinato-pinnatis ternatisque, caule aculeato, petiolis canaliculatis. *Sp. pl. p.* 706. *Fl. dan. t.* 788. *Scop. carn. tom.* 1. *p.* 355.

Rubus caule ſpinoſo, ſubrecto, foliis quinatis & ternatis, ſubtus tomentoſis, fructibus hirſutis. *Hall. hiſt. n.* 1108.

Rubus idæus vulgaris. *Clus. hiſt. p.* 117.

In ſylvis. Fl. Majo, Iunio. Fr.

cæſius. 2. R. foliis ternatis ſubnudis: lateralibus bilobis, caule aculeato tereti. *Sp. pl. p.* 706. *Scop. carn. tom.* 1. *p.* 356.

Rubus caule aculeato proſtrato, foliis ternatis. *Hall. hiſt. n.* 1110.

Rubus minor. *Dod. pempt.* 742.

In dumetis, fruticoſis, agris, vineis, alibique. Fl. Majo, Iunio. Fr.

fruticoſus. 3. R. foliis quinato-digitatis ternatisque, caule petiolisque aculeatis. *Sp. pl. p.* 707. *Miller. illuſtr. ic. Scop. carn. tom.* 1. *p.* 356.

Rubus caule ſpinoſo, ſerpente, foliis pinnatis & ternatis, ſubtus tomentoſis, baccis lævibus. *Hall. hiſt. n.* 1109.

Le Ronce. Regn. bot.

In collibus apricis aſperis, ad ſepes, maceries. Fl. æſtate. Fr.

Cum larva Phalenæ mori, ante evolutionem foliorum Mori albæ, ex ovo exit, tunc teneris Rubi fruticoſis foliis hanc nutriunt fluminenſes feminæ. Foliorum pulverem in nimiis ventris profluviis ad ſcrupulum unum quater de die a Liburniæ ruricolis optimo cum effectu exhibitum fuiſſe, vidi.

ſaxatilis. 4. R. foliis ternatis nudis, flagellis reptantibus herbaceis. *Sp. pl. p.* 708. *Fl. dan. t.* 134. *Scop. carn. tom.* I. *p.* 357.

Rubus caule herbaceo, foliis glabris, ternatis, acinis pauciſſimis. *Hall. hiſt. n.* 1111.

Rubus ſaxatilis alpinus. *Clus. hiſt. p.* 118.

In alpinis, & ſubalpinis. Fl. Majo, Iunio. Per.

FRAGARIA. Calyx decemfidus. Petala quinque. Receptaculum ſeminum ovatum, pulpoſum, deciduum.

vesca. 1. F. flagellis reptantibus. *Syſt. veget. p.* 476.

Fragaria foliis ternatis, flagellis reptans. *Hall. hiſt. n.* 1112.

Potentilla vesca. *Scop. carn. tom.* I. *p.* 363.

Le Fraiſier. Regn. bot.

In ſylvis, pascuis. Fl. æſtate. Per.

POTENTILLA. Calyx decemfidus. Petala quinque. Semina ſubrotunda, nuda, receptaculo parvo exſucco affixa.

Anſerina. 1. P. foliis pinnatis ſerratis, caule repente, pedunculis unifloris. *Syſt. veget. p.* 477. *Curtis lond. Fl. dan. t.* 544. *Scop. carn. tom.* I. *p.* 358.

Fragaria caule repente, foliis pinnatis, ſerratis, tomentoſis, petiolis longis unifloris. *Hall. hiſt. n.* 1126.

L' Argentine. Regn. bot.

In pratis, ad fluvios, vias. Fl. æſtate. Per.

rupeſtris. 2. P. foliis pinnatis alternis: foliolis quinis ovatis, crenatis, caule erecto. *Sp. pl. p.* 711. *Iacqu. auſtr. t.* 114. *Scop. carn. tom.* I. *p.* 358.

Fragaria foliis pinnatis, pinis ſeptenis, ſubrotundis, ſerratis. *Hall. hiſt. n.* 1127.

Quinquefolium quintum fragiferum. *Clus. hiſt. p. CVII.*

In pratenſibus, ſylvaticis Auſtriæ, Pannoniæ, Styriæ, Carnioliæ. Fl. Majo, Iunio Per.

Folia radicalia & caulina ima ſunt pinnata, ſumma ternata.

ſupina. 3. P. foliis pinnatis, caule dichotomo decumbente. *Sp. pl. p.* 711. *Iacqu. auſtr. t.* 406.

Pentaphyllum ſupinum Potentillæ facie. *Clus hiſt. p. CVIII.*

In agri vindobonenſis cultis, ruderatis. Fl. primo ætatis anno ab Auguſto ad Octobrim, altero jam Majo.

recta. 4. P. foliis ſeptenatis, foliolis lanceolatis ſerratis utrinque ſubpiloſis, caule erecto. *Sp. pl. p.* 711. *Iacqu. auſtr. t.* 383. *Scop. carn. tom.* 1. *p.* 359.

Fragaria caule erecto, foliis quinis & ſeptenis, acute ſerratis. *Hall. hiſt. n.* 1121.

In muris vetuſtis, in ſylvaticis, aſperis. Fl. Majo, Iunio, Iulio. Per.

Folia omnia ſunt digitata: infima longe petiolata & ſeptenata; ſumma quinata & ſubſeſſilia.

argentea. 5. P. foliis quinatis cuneiformibus inciſis ſubtus tomentoſis, caule erecto. *Sp. pl. p.* 712. *Fl. dan. t.* 865. *Scop. carn. tom.* 1. *p.* 360.

Fragaria caule erecto, foliis quinatis, apice ſerratis, ſubtus tomentoſis. *Hall. hiſt. n.* 1124.

In muris vetuſtis, ruderatis, apricis aſperis. Fl. Iunio. Per.

opaca. 6. P. folis radicalibus quinatis cuneiformibus ſerratis, caulinis ſuboppoſitis, ramis filiformibus decumbentibus. *Sp. pl. p.* 713. *Iacqu. collect. tom.* 1. *p.* 33. *ic. t.* 91.

Quinquefolii quarti flavo flore ſecunda ſpecies. *Cluſ. hiſt. p. CVI.*

In collibus apricis. Fl. Aprili, Majo. Per.

verna. 7. P. foliis radicalibus quinatis acute ſerratis, retuſis, caulinis ternatis, caule declinato. *Sp. pl. p.* 712.

Quinquefolii quarti flavo flore prima ſpecies. *Cluſ. hiſt. p. CVI.*

In apricis aſperis, collibus herbidis, montoſis ſylvaticis. Fl. primo vere. Per.

In apricis aſperis, collibus editioribus creſcens habet caulem brevem, erectum; folia omnia ternata, magis tomentoſa.

aurea. 8. P. foliis radicalibus quinatis ſerratis acuminatis, caulinis ternatis, caule declinato. *Sp. pl. p.* 712. *Iacqu. en. p.* 245.

Potentilla verna. *Fl. dan. t.* 114.

Quinquefolium tertium aureo flore. *Cluſ. hiſt. p. CVI.*

In alpinis, & ſubalpinis. Fl. æſtate. Per.

alba. 9. P. foliis quinatis apice conniventi-ſerratis, caulibus procumbentibus, receptaculis hirſutis. *Sp. pl. p.* 713. *Iacqu. auſtr. t.* 115.

Potentilla nitida. *Scop. carn. tom.* 1. *p.* 362.

Quinquefolium primum majus flore albo. *Cluſ. hiſt. p. CV.*

In montoſis ſylvaticis. Fl. primo vere, dein & Autumno. Per.

caules-cens. 10. P. foliis quinatis apice conniventi-ſerratis, caulibus multifloris decumbentibus, receptaculis hirſutis, petalis oblongis. *Syſt. veget. p.* 478. *Iacqu. auſtr. t.* 220.

Fragaria foliis ovatis acute ſerratis, oris ſericeis, pappo disci longiſſimo. *Hall. hiſt. n.* 1123.

In fiſſuris rupium Auſtriæ circa *Schottwien*, in monte *Gieſing* arci *Stüchſenſtein* imminenti, alibique. In ſimilibus locis in reliquis provinciis, alpinis, & ſubalpinis. Fl. Iulio, Auguſto. Per.

cluſiana. 11. P. foliis quinatis apice conniventi-ſerratis, caulibus multifloris decumbentibus, receptaculis hirſutis, petalis ſubrotundis. *Syſt. veget. p.* 478. *Iacqu. auſtr. t.* 116.

Quinquefolium ſecundum minus albo flore. *Cluſ. hiſt. p.* CV.

In rupeſtribus, ſaxoſis alpium auſtriacarum, ſtyriacarum, carniolicarum. Fl. æſtate. Per.

Potentilla caulescens habet foliola apice dentibus a tribus ad ſeptem notata. P. cluſianæ ſunt foliola tri- raro pluries dentata. In P. cluſiana omnia foliola omni ætate marginibus connivent. In caulescente adulta patent, &, dum juniora ſunt, connivent. P. caulescenti eſt calyx hirſutus, petala oblonga, receptaculum ex albo piloſum. P. cluſiana donatur calyce hirſuto, purpureo, petalis ſubrotundis.

nitida. 12. P. foliis ſubternatis tomentoſis conniventi-tridentatis, caulibus unifloris, receptaculis lanatis. *Syſt. veget. p.* 478. *Iacqu. auſtr. app. t.* 25.

In rupeſtribus editiſſimarum alpium carniolicarum, carinthiacarum, tyrolenſium. Fl. Iulio, Auguſto. Per.

reptans. 13. P. foliis quinatis, caule repente, pedunculis unifloris. *Sp. pl. p.* 714. *Curtis lond. Scop. carn. tom.* 1. *p.* 361.

Fragaria foliis quinatis, ſerratis, petiolis unifloris, caule reptante. *Hall. hiſt. n.* 1118.

La Quinte-Feuille. Regn. bot.

In pratis, ruderatis, ad vias, alibique. Fl. æſtate. Per.

TORMENTILLA. Calyx octofidus. Petala quatuor. Semina rotunda, nuda, receptaculo parvo, exſucco affixa.

erecta. 1. T. caule erectiusculo, foliis ſeſſilibus. *Sp. pl. p.* 716. *Curtis lond. Fl. dan t.* 589.

Fragaria tetrapetala foliis caulinis ſeſſilibus, quinatis. *Hall. hiſt. n.* 1117.

Potentilla Tormentilla erecta. *Scop. carn. tom.* 1. *p.* 360.

La Tormentille. Regn. bot.

In pratis, paſcuis alpinis, & ſubalpinis. Fl. æſtate. Per.

GEUM. Calyx decemfidus. Petala quinque. Semina ſtylo longo ariſtata.

urbanum. 1. G. floribus erectis, ariſtis uncinatis nudis, foliis caulinis ternatis, radicalibus lyrato-pinnatifidis. *Ait. Kew. tom.* 2. *p.* 218. *Sp. pl. p.* 716. *Curtis lond. Fl. dan. t.* 672.

Geum foliis pinnatis, pinna ultima trilobata, floribus patulis, tubis aduncis. *Hall. hiſt. n.* 1130.

Caryophyllata urbana. *Scop. carn. tom.* 1. *p.* 364.

Caryophyllata vulgaris. Geum Plinii. *Clus. hiſt. p. CII.*

La Benoite. Regn. bot.

In ſylvaticis, ad ſepes. Fl. Iunio. Per.

rivale. 2. G. floribus nutantibus, ariſtis uncinatis villoſis, petalis retuſis ſubrotundo-cuneiformibus, foliis pinnatis. *Ait. Kew. tom.* 2. *p.* 218. *Sp. pl. p.* 717. *Fl. dan. t.* 722.

Geum foliis pinnatis, pinnis acutis, extrema maxima, trilobata, floribus conniventibus, nutantibus. *Hall. hiſt. n.* 1129.

Caryophyllata rivalis. *Scop. carn. tom.* 1. *p.* 365.

Caryophyllata montana prima. *Cluſ. hiſt. p. CIII.*

Ad rivulos, & in aliis humidis locis ſubalpinis, & alpinis. Fl. Iunio, Iulio. Per.

hybridum. 3. G. flore nutante, calyce folioſo corollam polypetalam excedente. *Syſt. veget. p.* 480. *Wulfen in Iacqu. miscell. tom.* 2. *p.* 33. *ic. t.* 93.

Caryophyllata montana tertia. *Cluſ. hiſt. p. CIV.*

In alpe inferioris Carinthiæ non procul a *Pleyberg Pezzen* dicta, prope ſtabula. *Wulfen.* In Tyroli in paſcuis Pompeagi montis. *Scopoli.* Fl. Iunio. Per.

montanum. 4. G. caule unifloro, ariſtis rectis villoſis, foliis pinnatis pilolis: foliolo extimo maximo ſubrotundo; inferioribus ſenſim minoribus. *Ait. Kew. tom.* 2. *p.* 218. *Sp. pl. p.* 717. *Iacqu. auſtr. t.* 373.

Geum foliis pinnatis, extrema maxima, rotunde cordata. *Hall. hiſt. n.* 1131.

Caryophyllata montana. *Scop. carn. tom.* 1. *p.* 365.

Caryophyllata montana ſecunda. *Cluſ. hiſt. p. CIII.*

In alpibus. Fl. Iulio, Auguſto. Per.

reptans. 5. G. caulibus unifloris, ariſtis rectis villoſis, foliis pinnatis inciſis piloſis, flagellis reptantibus. *Ait. Kew. tom.* 2. *p.* 219. *Sp. pl. p.* 717. *Iacqu. auſtr. app. t.* 22.

Geum foliis pinnatis, trilobatis, inciſis, flagellis reptans. *Hall. hiſt. n.* 1132.

In abruptis præcipitiis ſummarum alpium *Winterthal*, *Eiſenhut*, tum etiam ſolo arenoſo ſupremarum viarum cavarum inter Carinthiam & Styriam ſupra alpes fladnizenſes. *Wulfen.* Ego in Tyroli in *Sellrainerthal* in arenoſis non procul a *Lüſenſerferner.* Fl. Iulio, Auguſto. Per.

DRYAS. Calyx quinque - ſeu decemfidus. Petala tot, quot laciniæ calycis. Semina caudata, piloſa.

octopetala. 1. D. floribus octopetalis, foliis ſimplicibus. *Sp. pl. p.* 717. *Fl. dan. t.* 31. *Scop. carn. tom.* 1. *p.* 366.

Dryas floribus octopetalis, foliis ovatis, ſubtus albiſſimis. *Hall. hiſt. n.* 1133.

Chamædrys tertia ſive montana. *Clus. hiſt. p.* 351.

In alpibus. Fl. Iunio, Iulio. Per.

COMARUM. Calyx decemfidus. Petala quinque, calyce minora. Receptaculum ſeminum ovatum, ſpongioſum, perſiſtens.

paluſtre. 1. Comarum. *Sp. pl. p.* 718. *Fl. dan. t.* 636.

Fragaria foliis pinnatis, petalis lingulatis minimis. *Hall. hiſt. n.* 1128.

Potentilla paluſtris. *Scop. carn. tom.* 1. *p.* 359.

In Carniolia in paludibus labacenſibus. *Scop.* Ego in Tyroli in *Sellrainerthal* locis udis. Fl. Iulio, Auguſto. Per.

POLYANDRIA.

MONOGYNIA.

CAPPARIS. Calyx tetraphyllus, coriaceus. Petala quatuor. Stamina longa. Bacca corticoſa, unilocularis, pedunculata.

ſpinoſa. 1. C. pedunculis unifloris ſolitariis, ſtipulis ſpinoſis, foliis annuis, baccis ovalibus. *Sp. pl. p.* 720. *Scop. carn. tom.* 1. *p.* 367. *Hall. hiſt. n.* 1077.

Le Caprier. Regn. bot.

In Litorali in rupibus ſub arce duinenſi. In Iſtria in muris civitatis Albonæ. In Forojulio Gradiscæ ad Liſonzium. Fr.

ACTÆA. Calyx tetraphyllus. Corolla tetrapetala. Bacca unilocularis. Semina ſemiorbiculata, gemino ordine incumbentia.

ſpicata. 1. A. racemo ovato, fructibus baccatis. *Sp. pl. p.* 722. *Fl. dan. t.* 498. *Scop. carn. tom.* 1. *p.* 368.

Actæa foliis circumſerratis, duplicato-pinnatis, pinnula extrema trilobata. *Hall. hiſt. n.* 1076.

Chriſtophoriana. *Clus. hiſt. p. LXXXVI.*

In opacis, aſperis ſylvarum montoſarum, & ſubalpinarum. Fl. æſtate. Per.

CHELIDONIUM. Calyx diphyllus. Corolla tetrapetala. Pericarpium ſiliquoſum, uni- vel biloculare, lineare.

majus. 1. C. pedunculis umbellatis. *Sp. pl. p.* 723. *Fl. dan. t.* 542. 676. *Scop. carn. tom.* 1. *p.* 369.

Chelidonium lobis foliorum ſubrotundis. *Hall. hiſt. n.* 1059.

Chelidonium majus vulgare. *Clus. hiſt. p.* CCIII.

La Chelidoine. Regn. bot.

Ad ſepes, ſylvarum margines, & alibi. Fl. æſtate. Per.

Glaucium. 2. C. pedunculis unifloris, foliis amplexicaulibus ſinuatis, caule glabro. *Sp. pl. p.* 724. *Fl. dan. t.* 585.

Glaucium foliis radicalibus ſemipinnatis, caulinis amplexicaulibus. *Hall. hiſt. n.* 1060.

Glaucium luteum. *Scop. carn. tom.* 1. *p.* 369.

Papaver corniculatum flavo flore. *Clus. hiſt. p.* XCI.

Le Pavot cornu. Regn. bot.

In Litorali ad mare. Fl. æſtate.

corniculatum. 3. C. pedunculis unifloris, foliis ſeſſilibus pinnatifidis, caule hiſpido. *Sp. pl. p.* 724.

Papaver corniculatum phœniceo flore. *Clus. hiſt. p.* XCI.

In Auſtria, Pannonia inter ſegetes, ad agrorum margines. Fl. Iunio, Iulio.

PAPAVER. Calyx diphyllus. Corolla tetrapetala. Capſula unilocularis, ſub ſtigmate perſiſtente poris dehiſcens.

Argemone. 1. P. capſulis clavatis hiſpidis, caule folioſo multifloro. *Sp. pl. p.* 725. *Fl. dan. t,* 867. *Scop. carn. tom.* 1. *p.* 370.

Papaver foliis hiſpidis, pinnatis, pinnis lobatis, capſulis ellipticis, hiſpidis. *Hall. hiſt. n.* 1061.

In agro goritienſi. *Wulfen.*

alpinum. 2. P. capſula hiſpida, ſcapo unifloro nudo hiſpido, foliis bipinnatis. *Sp. pl. p.* 725. *Iacqu. auſtr. t.* 83. *Scop. carn. tom.* 1. *p.* 371.

Papaver caule nudo, unifloro, foliis pinnatis, pinnis integris & diviſis, capitulis ſubrotundis hiſpidis. *Hall. hiſt. n.* 1062.

In alpium præcipitiis ſaxoſis, Fl. Iulio, Auguſto, Per.

Folia utrinque pilis albis hirta, bipinnatifida: pinnis exiguis, modo indiviſis, modo bifidis, trifidis; laciniis lanceolatis. Flos albus, vel flavus.

Rhœas. 3. P. capſulis glabris, globoſis, caule piloſo, multifloro, foliis pinnatifidis inciſis. *Sp. pl. p.* 726. *Curtis lond. Scop. carn. tom.* 1. *p.* 372.

Papaver foliis ſemipinnatis, hiſpidis, fructu ovato, glabro. *Hall. hiſt. n.* 1064.

Le Pavot rouge, ou Coquelicot. Regn. bot.

Inter ſegetes. Fl. Majo, Iunio.

Setæ, quibus caulis obſitus eſt, patent, & molliores ſunt quam in ſequente.

dubium. 4. P. capſulis oblongis glabris, caule multifloro ſetis adpreſſis, foliis pinnatifidis inciſis. *Sp. pl. p.* 726. *Iacqu. auſtr. t.* 25. *Curtis lond.*

Papaver foliis hiſpidis pinnatis, pinnis lobatis, fructu ovato lævi. *Hall. hiſt. n.* 1063.

In agri vindobonenſis collibus, vineis, inter ſegetes. Fl. Iunio.

Setæ caulinæ ſuperiores adpreſſæ, inferiores magis patentes. Variat quoque petalis albis, & rubris.

NYMPHÆA. Corolla polypetala. Calyx tetra- ſeu pentaphyllus. Bacca carnoſa, apice coronata, multilocularis.

lutea. 1. N. foliis cordatis integerrimis; calyce pentaphyllo petalis majore. *Sp. pl. p.* 729. *Fl. dan. t.* 603. *Scop. carn. tom.* 1. *p.* 372. *Cluſ. hiſt. p. LXXVII.*

Nymphæa calyce pentaphyllo petalis majori, fructu conico. *Hall. hiſt. n.* 1066.

In aquis ſtagnantibus, vel lente fluentibus. Fl. æſtate. Per.

alba. 2. N. foliis cordatis, integerrimis, calyce quadripartito. *Sp. pl. p.* 729. *Fl. dan. t.* 602. *Scop. carn. tom.* I. *p.* 373. *Cluſ. hiſt. p. LXXVII.*

Nymphæa calyce quadrifolio, petalis minori, fructu ſphærico. *Hall. hiſt. n.* 1067.

Le Nenufar. Regn. bot.

Cum præcedente. Fl. Iunio, Iulio. Per.

TILIA. Calyx quinquepartitus. Corolla pentapetala. Bacca ſicca, globoſa, quinquelocularis, quinquevalvis, baſi dehiscens.

europaea. I. T. floribus nectario deſtitutis. *Sp. pl. p.* 733. *Fl. dan. t.* 553.

Tilia foliis cordato-lanceolatis, ſpongiolis ad nervorum angulos ſeſſilibus. *Hall. hiſt. n.* 1030.

Le Tilleul. Regn. bot.

In ſylvis, ad pagos, fluvios. Fl. Iunio. Arb.

CISTUS. Calyx pentaphyllus: foliolis duobus minoribus. Corolla pentapetala. Capſula calyce obtecta.

ſalvifolius. I. C. arboreseens exſtipulatus, foliis ovatis petiolatis utrinque hirſutis. *Sp. pl. p.* 738. *Scop. carn. tom.* I. *p.* 374. *Wulfen in Iacqu. collect. tom.* 2. *p.* 120. *t.* 8.

Ciſtus fruticoſus, foliis petiolatis, ovatis, rugoſis, ſerratis. *Hall. hiſt. n.* 1031.

Ciſtus femina. *Cluſ. hiſt. p.* 70.

Ciſtus femina, folio Salviæ. *Bauh. pin.* 464.

In montanis Litoralis, copioſiſſime in inſulis venetis. Fl. Majo, Iunio.

Fumana. 2. C. ſuffruticoſus exſtipulatus procumbens, foliis alternis linearibus margine ſcabris, pedunculis unifloris. *Sp. pl. p.* 740. *Iacqu. auſtr. t.* 252. *Scop. carn. tom.* 1. *p.* 375.

Ciſtus foliis duris, linearibus, confertis, petiolis unifloris, calycibus glabris. *Hall. hiſt. n.* 1032.

Chamæciſtus ſextus. *Clus. hiſt. p.* 75.

In collibus apricis, aſperis. Fl. æſtate.

canus. 3. C. ſuffruticoſus exſtipulatus procumbens, foliis oppoſitis obovatis villoſis ſubtus tomentoſis, floribus ſubumbellatis. *Sp. pl. p.* 740. *Iacqu. auſtr. t.* 277. *Scop. carn. tom.* 1. *p.* 377.

Ciſtus caulibus duris, procumbentibus, foliis ellipticis, hirſutis, inferne cum calycibus tomentoſis. *Hall. hiſt. n.* 1035.

Chamæciſtus tertius. *Clus. hiſt. p.* 74.

Cum præcedente. Fl. Majo.

œlandicus. 4. C. ſuffruticoſus exſtipulatus procumbens, foliis oppoſitis oblongis ſubhirſutis, petiolis ciliatis, petalis emarginatis. *Sp. pl. p.* 741. *Iacqu. auſtr. t.* 399.

Ciſtus caulibus duris, procumbentibus, foliis conjugatis, ellipticis, ſubhirſutis, petiolis confertis unifloris, calycibus tomentoſis. *Hall. hiſt. n.* 1034.

Ciſtus alpeſtris. *Iacqu. en. p.* 248. *Scop. carn. tom.* 1. *p.* 375.

Chamæciſtus ſecundus. *Clus. hiſt. p.* 73.

In alpinis. Fl. Iulio, Auguſto.

Folia apud nos utrinque & ad oras pilis albis hirſuta, quædam tamen dorſo minus, fereque hic glabra. Iacqu. l. c.

Helianthemum. 5. C. ſuffruticoſus procumbens, ſtipulis lanceolatis, foliis oblongis revolutis ſubpiloſis. *Sp. pl. p.* 744. *Fl. dan. t.* 101. *Curtis lond. Scop. carn. tom.* 1. *p.* 378.

Ciſtus foliis conjugatis, ellipticis, hirſutis, integerrimis, petiolis unifloris ſubhirſutis. *Hall. hiſt. n.* 1033.

Chamæciſtus primus. *Clus. hiſt. p.* 73.

L' Heliantheme, ou la Fleur du ſoleil. Regn. bot.

In pratis, collibus, montoſis ſylvaticis, & alibi. Fl. æſtate.

DIGYNIA.

PÆONIA. Calyx pentaphyllus. Petala quinque. Germina duo. Stylus nullus. Stigmata compreſſa. Capſulæ polyſpermæ.

officinalis. 1. P. foliis decompoſitis nudis, foliolis lobatis: lobis lato-lanceolatis, capſulis rectiuſculis tomentoſis. *Retzii obs. fasc.* 3. *p.* 35. *Sp. pl. p.* 747. *Scop. carn. tom.* 1. *p.* 378.

Pæonia radice glanduloſa, foliis duplicato-pinnatis, pinnis ellipticis & trilobis, *Hall. hiſt. n.* 1187.

La Pivoine. Regn. bot.

In Carnioliæ, Litoralis montanis, aſperis, copioſiſſime circa *Saguria, Materia, Lipa, Lipizza.* Fl. Majo. Per.

Variat foliolis lobatis, & indiviſis.

TRIGYNIA.

DELPHINIUM. Calyx nullus. Petala quinque. Nectarium bifidum, poſtice cornutum. Capſulæ tres ſeu unica.

Conſolida. 1. D. unicapſulare, nectariis monophyllis, caule ſubdiviſo. *Sp. pl. p.* 748. *Fl. dan. t.* 683. *Scop. carn. tom.* I. *p.* 379.

Delphinium uniſiliquoſum, caule brachiato diffuſo, floribus ſparſis. *Hall. hiſt. n.* 1203.

Delphinium ſeu Conſolida regalis. *Clus. hiſt. p.* CCV.

Le Pied d'Alouette. Regn. bot.

Inter ſegetes, alibique. Fl. æſtate.

intermedium. 2. D. tricapſulare, nectariis diphyllis: labellis ovatis, bifidis: laciniis ovatis, foliis tripartitis, laciniis trifidis inciſis. *Ait. Kew. tom.* 2. *p.* 243.

Aconitum Lycoctonum flore Delphinii primum ſileſiacum. *Clus. hiſt. p.* XCIV.

In Styria in alpibus judenburgenſibus in præcipitiis celſiſſimi jugi, *die ſcharfe Eck* dicti. Fl. Iulio, Auguſto. Per.

Radix tuberoſa, difformis. Caulis cubitalis, ſæpe altior, inanis, ſuperne ut flores eleganti cæruleo colore tinctus, ſimplex vel paucis ramis, ex axillis foliorum ſuperiorum ortis, auctus. Folia alternatim ſparſa, villoſa, tandem calveſcentia, palmata: laciniis acutis, integris vel diviſis. Flores racemoſi. Bracteæ ad apicem pedunculi duæ, ad baſin unica. Petalum ſupremum in cornu tubulatum, rugoſum extenſum; quatuor reliqua includunt nectaria quatuor cæruleſcentiatra; quorum duo ſuperiora duriora, integra, glabra, calcarata; calcaribus poſtice protenſis, & tubo petali ſupremi involutis: inferiora duo teneriora, limbo (labello) pilis ex viridi flaveſcentibus barbato, bifido: laciniis ovatis. Cl. Linnæus hanc ſpeciem cum aliis duabus confudit, & omnes tres ſub nomine Delphinii elati propoſuit. Separarunt in horto kewenſi Angli.

Staphiſagria. 3. D. tricapſulare, nectariis diphyllis petalo brevioribus, foliis palmatis: lobis obtuſis. *Sp. pl. p.* 750.

La Staphiſaigre. Regn. bot.

In Iſtria creſcere dicitur. Nec mihi ſinum flanaticum diligentiſſime per plures menſes perluſtranti, nec aliis in loco, a Cl. *Wulfen* indicato, invenire licuit.

ACONITUM. Calyx nullus. Petala quinque: ſupremo fornicato. Nectaria duo, pedunculata, recurva. Capſulæ tres ſeu quinque.

Lycoctonum. 1. A. floribus trigynis, foliis palmatis multifidis villoſis. *Syſt. veget. p.* 504. *Iacqu. auſtr. t.* 380.

Aconitum Lycoctonum vulgare luteo flore. *Clus. hiſt. p.* XCIX.

In montanis, & ſualpinis. Fl. a Iulio ad Septembrim. Per.

Napellus. 2. A. floribus trigynis, foliis multipartitis, laciniis linearibus. *Wulfen in Kæl. ſpic. p.* 14.

Aconitum cernuum. *ibid. p.* 17.

Aconitum tauricum. *ibidem. p.* 15. *in Iacqu. collect. tom.* 2. *p.* 112. *ic. t.* 492.

Aconitum Lycoctonum quartum tauricum. *Clus. hiſt. p.* XCV.

Aconitum Lycoctonum octavum coma nutante. *Clus. hiſt. p.* XCVII.

In Carniolia in alpibus wochinenſibus. In Carinthia in Taurero malnizenſi, raſtadienſi, in alpibus prægratenſibus. Fl. Iulio, Auguſto.

Radix napiformis. Folia petiolata, alternatim ſparſa, utrinque glabra, facie ſaturate, dorſo pallide viridia, multipartita: laciniis linearibus. Caulem terminat racemus, qui quandoque uno alterove racemulo, ex foliorum axillis orto, augetur. Piſtilla tria, hisce raro quartum abortiens accedit. In apricis ſiccis caulis cubitalis

circiter; folia conferta; flores congeſti; florum color ſaturate cærulæus. In humidis, umbroſis, & in horto caulis multo altior; ante florum explicationem racemus debilis nutat, dein ſimior factus erigitur: folia circumſcriptione ampliora, inter ſe magis diſtantia; flores ſegregati; florum color dilute cæruleus.

variegatum. 3. A. floribus trigynis, foliorum laciniis ſemipartitis ſuperne latioribus. *Syſt. veget. p.* 504.

Lycoctonum cœruleum parvum. *Dod. pempt.* 438.

Lycoctonum cœruleum parvum facie Napelli. *Lob. hiſt.* 386.

Aconitum Lycoctonum decimum. Thora italica. *Clus. hiſt. p. XCVIII.* quoad figuram tantum.

In Carnioliæ ſylvis ſubalpinis, in fruticoſis inter *Prewald* & *Adlersberg*, alibique. Fl. a Iunio ad Auguſtum. Per.

Folia ſunt pallidiora, minus diviſa. Petalum ſupremum apice revolutum: duo lateralia inferne margine revoluta; infima duo obliqua. Omnia ex viridi & cæruleo variegata. Caulis, prout diverſa eſt ſoli natura, expoſitio, eſt cubitalis, vel altior.

Anthora. 4. A. floribus pentagynis, foliorum laciniis linearibus. *Sp. pl. p.* 751. *Iacqu. auſtr. t.* 382. *Scop. carn. tom.* 1. *p.* 381.

Anthora vulgaris. *Clus. hiſt. p. XCVIII.*

L' Aconit, ou l' Anthora. Regn. bot.

In Carniolia calidiore in Carſo. *Scop.* In Auſtria prope *Stüchſenſtein* in monte *Gieſing*. *Iacqu.* Fl. Auguſto, Septembri. Per.

Cammarum. 5. A. floribus ſubpentagynis, foliorum laciniis cuneiformibus inciſis acutis. *Syſt. veget. p.* 504. *Iacqu. auſtr. t.* 424.

Aconitum Napellus. *Iacqu. auſtr. t.* 381.

Aconitum Lycoctonum quintum neubergenſe. *Clus. hiſt. p.* XCVI.

Aconitum Lycoctonum ſextum, Napellus vulgaris. *Cluſ. hiſt. p.* XCVI.

Aconitum Lycoctonum nonum, judenburgenſe. *Cluſ. hiſt. p.* XCVII.

In ſubalpinis, & alpinis Auſtriæ, Styriæ. Fl. Iulio, Auguſto, Septembri. Per.

In demiſſioribus, umbroſis, ſucculentis caulis ſæpe bicubitalis, valde ramoſus; foliorum lobi minus profunde diviſi laciniæque breviores, latiores; flores in racemis pauciores, ſegregati, dilute cærulei. In editioribus, apricis caulis brevior, minus ramoſus; foliorum lobi magis diviſi, laciniæ anguſtiores; racemus conſtat floribus congeſtis, ſaturatæ cæruleis.

PENTAGYNIA.

AQUILEGIA. Calyx nullus. Petala quinque. Nectaria quinque, corniculata, inter petala. Capſulæ quinque, diſtinctæ.

vulgaris. 1. A. nectariis incurvis petalo ſubæqualibus, foliolis omnibus petiolatis. *Hænke in Iacqu. collect. tom. 2. p. 42. Sp. pl. p. 752. Scop. carn. tom. 1. p. 381.*

Aquilegia foliis trilobatis; lobis petiolatis tripartitis, ſubrotundis, obtuſe dentatis. *Hall. hiſt. n.* 1195.

Aquilina. *Cluſ. hiſt. p.* CCIV.

L' Ancolie. Regn. bot.

In montanis, ſylvaticis. Fl. Majo, Iunio. Per.

viſcoſa. 2. A. nectariis incurvis, caule ſubnudo ſubunifloro viſcoſo-piloſo, foliolis ſubtrilobis. *Syſt. veget. p. 505. Gouan. fl. p. 267. Magn. bot. 24. hort. 21.*

Aquilegia alpina. *Hænke in Iacqu. collect. tom. 2. p. 42.*

In Carniolia in alpibus wochinenſibus. In Tyroli in alpibus leontinis. Fl. Iulio, Auguſto. Per.

Vel ſola pube viſcida, qua caulis, pedunculi, piſtilla veſtita ſunt, diſtinguitur ab A. vulgari; vel ſola radice perenni, ab A. alpina, cui radix biennis eſt.

NIGELLA. Calyx nullus. Petala quinque. Nectaria quinque, trifida, intra corollam. Capſulæ quinque, connexæ.

arvenſis. 1. N. piſtillis quinis, petalis integris, capſulis turbinatis. *Sp. pl. p.* 753. *Scop. carn. tom.* 1. *p.* 382.

Nigella flore nudo, ſiliquarum cornubus longiſſimis. *Hall. hiſt. n.* 1194.

La Nielle des champs. Bulliard. herb. t. 126.

In arvis, ad agrorum margines, vineas. Fl. æſtate.

POLIGYNIA.

ANEMONE. Calyx nullus. Petala duorum triumve ordinum, in ſingula ſerie tria. Semina plura, ſtylum retinentia.

* *Flore ſubcalyculato.*

Hepatica. 1. A. foliis trilobis integerrimis. *Sp. pl. p.* 758. *Fl. dan. t.* 610. *Scop. carn. tom.* 1. *p.* 382. *Hall. hiſt. n.* 1156.

In nemoribus. Fl. Martio, Aprili. Per.

** *Pedunculo involucrato, ſeminibus caudatis.*

vernalis. 2. A. pedunculo involucrato, foliis pinnatis, flore erecto. *Sp. pl. p.* 759. *Fl. dan. t.* 29. *Wulfen in Iacqu. collect. tom.* 3. *p.* 10.

Anemone tubis caudatis, involucris multifidis, foliis pinnatis, pinnis latis, lobatis. *Hall. hiſt. n.* 1147. *t.* 21.

Anemone tubis caudatis, involucris multifidis, foliis hirſutis, pinnatis, pinnis acute lobatis. *Hall. hiſt. n.* 1148.

Anemone ſulphurea. *Allion. pedem. tom.* 2. *p.* 170.

Anemone Halleri. *Allion. pedem. tom.* 2. *p.* 170. *t.* 80. *f.* 2.

In Carinthiæ alpibus reichenavienſibus. In Tyroli in alpibus leontinis, & in montibus *in Neuſtift im Gerichte Stubay.* Fl. poſt ſolutas nives. Per.

Pulſatilla. 3. A. pedunculo involucrato, petalis rectis, foliis bipinnatis. *Sp. pl. p.* 759. *Fl. dan. t.* 153. *Scop. carn. tom.* 1. *p.* 386.

Anemone tubis caudatis, involucris unifoliis multifidis, foliis linearibus hirſutis, duplicato-pinnatis, pinnis tripartitis & quinquepartitis. *Hall. hiſt. n.* 1146.

Pulſatilla vulgaris dilutiore flore. *Clus. hiſt. p.* 246.

L' Anemone Pulſaltille. Bulliard *herb. t.* 49.

In collibus, montoſis ſylvaticis. Fl. primo vere. Per.

pratenſis. 4. A. pedunculo involucrato, petalis apice reflexis, foliis bipinnatis. *Sp. pl. p.* 760. *Fl. dan. t.* 611.

Pulſatilla vulgaris ſaturatiore flore. *Clus. hiſt. p.* 246.

Pulſatilla flore minore nigricante. *Bauh. pin.* 177.

In pratis ſiccioribus, collibus herbidis, montoſis apricis. Fl. primo vere. Per.

alpina. 5. A. foliis caulinis ternis connatis ſupradecompoſitis multifidis, ſeminibus hirſutis caudatis. *Sp. pl. p.* 760. *Iacqu. auſtr. t.* 85.

Anemome foliis triplicato-pinnatis, acute circumſerratis, caulinis ternatis, tubis caudatis. *Hall. hiſt. n.* 1149.

Anemone burſeriana. *Scop. carn. tom.* 1. *p.* 285.

Anemone ſulphurea. *Syſt. veget. p.* 509.

Anemone apiifolia. *Syſt. veget. p.* 510. *Wulfen in Iacqu. miscell. tom.* 2. *p.* 47. *t.* 4. *Scop. carn. tom.* 1. *p.* 285.

Anemone ſylveſtris secunda. *Clus hiſt. p.* 244.

Anemone ſylveſtris tertia. *Clus. hiſt. p.* 245.

Pulſatilla lutea. *Cam. epit.* 393. *Bauh. hiſt.* 3. *p.* 411.

In alpinis, & ſubalpinis. Fl. a Iunio ad Septembrim. Per.

Petala alba, per ſuccedentes gradus, in pallidum, citrinum & croceum colorem degenerant. Haller. l. c.

*** *Seminibus ecaudatis.*

hortenſis. 6. A. ſeminibus lanatis, foliis radicalibus trilobatis & multifidis, caulinis ovato-lanceolatis. *Hall. hiſt. n.* 1152. *Sp. pl. p.* 761. *Curtis magaz.* 123.

Anemone hortenſis latifolia ſimplici flore. *Clus. hiſt. p.* 249.

In agro fluminenſi in *Belvedere, Raſtozzine, Braida*, & alibi ad maceries, ſepes, in juniperetis. Fl. Februario, Martio. Per.

Radix tuberoſa, plantæ novellæ exiguæ olivæ magnitudine, adultæ forma & magnitudine varia, fibris paucis aucta. Folia dorſo purpuraſcentia, triloba, vel multifida. Caulis albido to-

mento veſtitus. Involucrum triphyllum: foliolis lanceolatis, vel integris, vel apice in duas tresve lacinias diviſis. Caulem terminat flos unicus, conſtans petalis duodecim, vel pluribus, purpureis, raro albis, lineari-lanceolatis, dorſo villöſis; externis quinque longioribus. Semina lanata.

ſylveſtris. 7. A. foliis digitatis, hirſutis, floribus pentapetalis, ſeminibus lanatis. *Hall. hiſt. n.* 1150. *Sp. pl. p.* 761. *Curtis magaz.* 54.

Anemone ſylveſtris prima. *Clus. hiſt. p.* 244.

Anemone blanche. Bulliard. herb. t. 59.

In collibus, montoſis ſylvaticis. Fl. Majo. Per.

baldenſis. 8. A. ſeminibus lanatis, foliis caulinis ternatis, triplicato-trifidis, lobulis lanceolatis, acute trifidis bifidisve. *Hall. hiſt. n.* 1151. *Syſt. veget. p.* 509.

Anemone fragifera. *Wulfen in Iacqu. miscell. tom.* 2. *p.* 55. *ic. t.* 103. *Syſt. veget. p.* 510.

Anemone alpina. *Scop. carn. tom.* 1. *p.* 384. *t.* 26.

Anemone alpina alba minor. *Bauh. pin.* 176. *prodr.* 94.

In alpibus Styriæ, Carnioliæ, Croatiæ, Carinthiæ, Tyrolis, Carniæ. Fl. a Iunio ad Auguſtum. Per.

trifolia. 9. A. foliis ternatis ovatis integris ſerratis, caule unifloro. *Sp. pl. p.* 762. *Scop. carn. tom.* 1. *p.* 383.

Anemone trifolia, flore albo. *Bauh. hiſt.* 3. *p.* 412.

In Carnioliæ alpium convallibus, alibique in nemoribus. Fl. Majo. Per.

nemoroſa. 10. A. ſeminibus acutis, foliolis inciſis, caule unifloro. *Sp. pl. p.* 762. *Fl. dan. t.* 549. *Scop. carn. tom.* 1. *p.* 383.

Anemone ſeminibus nudis, caule unifloro, foliis caulinis quinquepartitis, lobis tripartitis inciſis. *Hall. hiſt.* 1154.

Ranunculus ſylvarum. *Clus. hiſt. p.* 247.

L' Anemone des Bois. Bulliard herb. t. 3.

In nemorenſibus, ſylvaticis. Fl. primo vere. Per.

ranunculoides. 11. A. ſeminibus acutis, foliolis inciſis, petalis ſubrotundis, caule ſubbifloro. *Sp. pl. p.* 762. *Fl. dan. t.* 140. *Scop. carn. tom.* 1. *p.* 384.

Anemone biflora, ſeminibus nudis, foliis caulinis quinquepartitis, lobis tripartitis inciſis. *Hall. hiſt. n.* 1153.

Habitat & floret cum præc. Per.

narciſſiflora. 12. A. floribus umbellatis, ſeminibus depreſſo-ovalibus nudis. *Sp. pl. p.* 763. *Iacqu. auſtr. t.* 159.

Anemone ſeminibus nudis, rotundis, planis, foliis ſeptifidis imbricatis, floribus umbellatis. *Hall. hiſt. n.* 1155.

Ranunculus montanus ſecundus, & Ranunculi montani ſecundi ſpecies altera. *Clus. hiſt. p.* 235.

In Auſtriæ, Styriæ ſubalpinis, & alpinis. Fl. a Iunio ad Auguſtum. Per.

ATRAGENE. Calyx tetraphyllus. Petala duodecim. Semina caudata.

alpina. 1. A. foliis duplicato-ternatis ſerratis, petalis exterioribus quaternis. *Sp. pl. p.* 764. *Iacqu. auſtr. t.* 241.

Atragene foliis ternatis, foliolis trifoliis acutis. *Hall. hiſt. n.* 1145.

Atragene auſtriaca. *Scop. carn. tom.* 1. *p.* 387.

Clematis cruciata alpina. *Pon. in Clus. hiſt. p.* CCCXXI.

Clematis alpina geranifolia. *Bauh. pin.* 300. *prodr.* 135.

In rupeſtribus, ſaxoſis alpinis, & ſubalpinis. Fl. Iulio, Auguſto. Fr.

CLEMATIS. Calyx nullus. Petala quatuor, rarius quinque vel ſex. Semina caudata.

Viticella. 1. C. foliis compoſitis decompoſitisque: foliolis ovatis ſublobatis integerrimis. *Syſt. veget. p.* 512. *Scop. carn. tom.* 1. *p.* 388.

Clematis altera. *Clus. hiſt. p.* 122.

Clematis cœrulea vel purpurea repens. *Bauh. pin.* 300.

Clematis ſive flammula flore cœruleo ſcandens. *Bauh. hiſt.* 2. *p.* 128. *Zann. iſt. p.* 69. *t.* 80.

In Forojulio ad ſepes agri medenſis. *Scop.* Ego Duini; Flumine *al Ponſal*, *San Nicolò*, & alibi. Fl. Majo, Iunio. Fr.

Vitalba. 2. C. foliis pinnatis, foliolis cordatis ſcandentibus. *Sp. pl. p.* 766. *Iacqu. auſtr. t.* 308. *Curtis lond. Scop. carn. tom.* 1. *p.* 388.

Clematis caule ſcandente, foliis pinnatis, ovato-lanceolatis. *Hall. hiſt. n.* 1142.

Atragene Theophraſti. *Clus. hiſt p.* 122.

La Clematite des Haies. Bulliard herb. t. 89.

Ad ſepes, alibique. Fl. Auguſto, Septembri. Per.

Flammula. 3. C. foliis inferioribus pinnatis ſcandentibus laciniatis: ſummis ſimplicibus integerrimis lanceolatis. *Sp. pl. p.* 766.

Clematis cespitoſa. *Scop. carn. tom.* 1. *p.* 389.

Clematis maritima. *Allion. nic.* 122.

Clematis maritima repens. *Bauh. pin.* 300. *prodr.* 135. *Zann. iſt. p.* 68. *t.* 129.

In Litorali ad ſepes, in fruticoſis agri flu-minenſis, duinenſis, & alibi. Fl. Iunio, Iulio. Fr.

Radix emittit caules plures, ſarmentoſos, ramoſos, ſi frutices in vicinia, ſcandentes, ſi nulli, inter ſe et cum ramis intertextos, humi diffuſos, paucos breviores erectos. Folia ſumma ſunt ſimplicia, reliqua bipinnata, pinnata, & hisce mixta bigemina, ternataque: foliolis aliis lanceolatis, quibusdam ovatis, aliquibus linearibus; omnibus integris, aut bi- ſeu trifidis. Corolla conſtat petalis quatuor vel quinque, lanceolatis, lineatis, apice compreſſis. Stamina numeroſa. Piſtilla a quinque ad octo. Hanc C. maritimam repentem Bauhini eſſe, autopta teſtor. Quod C. Flammula ſit quoque C. maritima Allionii, candide in Auctario ſuo ad Floram pedemontanam ipſemet Cl. vir teſtatur dicens. „Deleatur C. maritima, que eſt vere Flammula. Ea, quam pro Flammula habui, varietas eſt C. erectæ.“

erecta. 4. C. foliis pinnatis: foliolis ovato-lanceolatis, integerrimis, caule erecto, floribus pentapetalis tetrapetalisque. *Sp. pl. p.* 767. *Iacqu. auſtr. t.* 291. *Scop. carn. tom.* I. *p.* 387.

Clematis caule erecto, foliis pinnatis, ovato-lanceolatis. *Hall. hiſt. n.* 1144.

Flammula Matthioli. *Clus. hiſt. p.* 124.

Ad ſepes, in fruticoſis, montoſis ſylvaticis. Fl. æſtate. Per.

integrifolia. 5. C. foliis ſimplicibus ovato-lanceolatis, floribus cernuis. *Sp. pl. p.* 767. *Iacqu. auſtr. t.* 363. *Scop. carn. tom.* I. *p.* 387.

Clematis cœrulea pannonica. *Clus. hiſt. p.* 123.

In pratis paluſtribus Duini, Aquilejæ. In ſimilibus locis in Auſtria circa *Bruck an der Leitha*; & in Pannonia ad lacum Peiſonis, alibique. Fl. Iunio, Iulio. Per.

THALICTRUM. Calyx nullus. Petala quatuor ſeu quinque. Semina ecaudata.

alpinum. 1. T. caule ſimplicisſimo ſubnudo, racemo ſimplici terminali. *Sp. pl. p.* 767. *Fl. dan. t.* 11.

In Tyroli in alpibus leontinis. Per.

minus. 2. T. foliis ſexpartitis, floribus cernuis. *Sp. pl. p.* 769. *Iacqu. auſtr. t.* 419. *Scop. carn. tom.* I. *p.* 390.

Thalictrum foliis parvulis, imis obtuſe trilobis, ſupremis acutis, panicula diffuſa, floribus nutantibus. *Hall. hiſt. n.* 1139.

In collibus herbidis, montoſis apricis, & in ipſis alpibus. Fl. Iunio, Iulio. Per.

Folia multiplicato-pinnata: foliolis breviſſime petiolatis vel ſeſſilibus, venoſis, ad apicem bis terve inciſis aut integerrimis, obtuſis vel acutis. Piſtilla tria vel quatuor. Iacqu. l. c.

majus. 3. T. foliis multipartitis, floribus cernuis, ramis floriferis ex iisdem axillis binis vel ternis. *Syſt. veget. p.* 513. *Iacqu. auſtr. t.* 420.

In Auſtriæ, Pannoniæ ſylvaticis. Fl. Iunio, Iulio. Per.

In majori piſtilla quatuor ad ſeptem; ex eodem caulis & ramorum principum nodo bini vel terni egrediuntur rami, qui in minori fere ſemper ſolitarii. Foliorum prona facies in majori magis ſaturate veluti ex atro viret, poſtica autem intenſe & ex albo glauca eſt; dum illa in minori viret lætius, hæc vero pallet. Iacqu. l. c.

anguſtifolium. 4. T. foliolis lanceolato-linearibus integerrimis. *Sp. pl. p.* 769. *Iacqu. hort. tom.* 3. *p.* 25. *t.* 43. *Scop. carn. tom.* I. *p.* 391.

Thalictrum foliis aſcendentibus, linearibus, panicula erecta. *Hall. hiſt. n.* 1137.

Thalictrum pratenſe anguſtiſſimo folio. *Bauh. pin.* 337. *prodr.* 146.

In Auſtriæ, Styriæ, Carnioliæ pratis udis. Fl. Iunio, Iulio. Per.

flavum. 5. T. caule folioſo ſulcato; panicula multiplici erecta. *Sp. pl. p.* 770.

Thalictrum foliis erectis, lobulis acute trilobis, ſpica adſcendente. *Hall. hiſt. n.* 1138.

Thalictrum nigricans. *Iacqu. auſtr. t.* 421. *auct. Cl. Sibthorp.*

In Auſtria, Pannonia in inſulis danubialibus, pratis paluſtribus. Fl. Iunio. Per.

Radix repens. Foliola lanceolata, acuta, facie ſaturate virentia, ſubtus pallidiora venisque rugoſula, utrinque glabra; alia integra, alia ter vel bis inciſa, idque inprimis in foliis inferioribus, ubi etiam latiora ſunt quam in ſuperioribus. Flores erecti. Piſtilla a duobus ad quinque. Iacqu. l. c.

aquilegifolium. 6. T. fructibus pendulis triangularibus rectis, caule tereti. *Sp. pl. p.* 770. *Iacqu. auſtr. t.* 318. *Scop. carn. tom.* 1. *p.* 389.

Thalictrum foliis triplicato-pinnatis, patulis, obtuſe trilobatis, ſtipulis ovatis, paniculis umbellatis. *Hall. hiſt. n.* 1141.

In alpinis, & ſubalpinis. Fl. æſtate. Per.

Caulis fiſtuloſus. Folia triternata, glabra, ad exortum & diviſiones munita ſtipulis rotundatis, perfoliatis. Foliola ſunt fere ſubrotunda, ſuperne obtuſe dentato-inciſa, ſubtus glauca, venuloſa. Petala quatuor, obverſe ovata, concava, ſtaminibus breviora. Flores paniculati. Stamina plurima. Piſtilla ſeptem vel octo. Capſulæ pendulæ, triangulares, trialatæ. Iacqu. l. c.

ADONIS. Calyx pentaphyllus. Petala absque nectario, quinque vel plura. Semina nuda.

autumnalis. 1. A. floribus octopetalis, fructibus ſubcylindricis. *Sp. pl. p.* 771.

Adonis autumnalis, corollis flavo-rubris. *Scop. carn. tom.* 1. *p.* 392.

Staphiſagria. 3. D. tricapſulare, nectariis diphyllis petalo brevioribus, foliis palmatis: lobis obtuſis. *Sp. pl. p.* 750.

La Staphiſaigre. Regn. bot.

In Iſtria creſcere dicitur. Nec mihi ſinum flanaticum diligentiſſime per plures menſes perluſtranti, nec aliis in loco, a Cl. *Wulfen* indicato, invenire licuit.

ACONITUM. Calyx nullus. Petala quinque: ſupremo fornicato. Nectaria duo, pedunculata, recurva. Capſulæ tres ſeu quinque.

Lycoctonum. 1. A. floribus trigynis, foliis palmatis multifidis villoſis. *Syſt. veget. p.* 504. *Iacqu. auſtr. t.* 380.

Aconitum Lycoctonum vulgare luteo flore. *Clus. hiſt. p.* XCIX.

In montanis, & ſualpinis. Fl. a Iulio ad Septembrim. Per.

Napellus. 2. A. floribus trigynis, foliis multipartitis, laciniis linearibus. *Wulfen in Kœl. ſpic. p.* 14.

Aconitum cernuum. *ibid. p.* 17.

Aconitum tauricum. *ibidem. p.* 15. *in Iacqu. collect. tom.* 2. *p.* 112. *ic. t.* 492.

Aconitum Lycoctonum quartum tauricum. *Clus. hiſt. p.* XCV.

Aconitum Lycoctonum octavum coma nutante. *Clus. hiſt. p.* XCVII.

In Carniolia in alpibus wochinenſibus. In Carinthia in Taurero malnizenſi, raſtadienſi, in alpibus prægratenſibus. Fl. Iulio, Auguſto.

Radix napiformis. Folia petiolata, alternatim ſparſa, utrinque glabra, facie ſaturate, dorſo pallide viridia, multipartita: laciniis linearibus. Caulem terminat racemus, qui quandoque uno alterove racemulo, ex foliorum axillis orto, augetur. Piſtilla tria, hisce raro quartum abortiens accedit. In apricis ſiccis caulis cubitalis

circiter; folia conferta; flores congeſti; florum color ſaturate cœrulæus. In humidis, umbroſis, & in horto caulis multo altior; ante florum explicationem racemus debilis nutat, dein ſimior factus erigitur: folia circumſcriptione ampliora, inter ſe magis diſtantia; flores ſegregati; florum color dilute cæruleus.

variegatum. 3. A. floribus trigynis, foliorum laciniis ſemipartitis ſuperne latioribus. *Syſt. veget. p.* 504.

Lycoctonum cœruleum parvum. *Dod. pempt.* 438.

Lycoctonum cœruleum parvum facie Napelli. *Lob. hiſt.* 386.

Aconitum Lycoctonum decimum. Thora italica. *Clus. hiſt. p. XCVIII.* quoad figuram tantum.

In Carnioliæ ſylvis ſubalpinis, in fruticoſis inter *Prewald* & *Adlersberg*, alibique. Fl. a Iunio ad Auguſtum. Per.

Folia ſunt pallidiora, minus diviſa. Petalum ſupremum apice revolutum: duo lateralia inferne margine revoluta; infima duo obliqua. Omnia ex viridi & cæruleo variegata. Caulis, prout diverſa eſt ſoli natura, expoſitio, eſt cubitalis, vel altior.

Anthora. 4. A. floribus pentagynis, foliorum laciniis linearibus. *Sp. pl. p.* 751. *Iacqu. auſtr. t.* 382. *Scop. carn. tom.* 1. *p.* 381.

Anthora vulgaris. *Clus. hiſt. p. XCVIII.*

L' Aconit, ou l' Anthora. Regn. bot.

In Carniolia calidiore in Carſo. *Scop.* In Auſtria prope *Stüchſenſtein* in monte *Gieſing. Iacqu.* Fl. Auguſto, Septembri. Per.

Cammarum. 5. A. floribus ſubpentagynis, foliorum laciniis cuneiformibus inciſis acutis. *Syſt. veget. p.* 504. *Iacqu. auſtr. t.* 424.

Aconitum Napellus. *Iacqu. auſtr. t.* 381.

Aconitum Lycoctonum quintum neubergenſe. *Clus. hiſt. p.* XCVI.

Aconitum Lycoctonum ſextum, Napellus vulgaris. *Cluſ. hiſt. p.* XCVI.

Aconitum Lycoctonum nonum, judenburgenſe. *Cluſ. hiſt. p.* XCVII.

In ſubalpinis, & alpinis Auſtriæ, Styriæ. Fl. Iulio, Auguſto, Septembri. Per.

In demiſſioribus, umbroſis, ſucculentis caulis ſæpe bicubitalis, valde ramoſus; foliorum lobi minus profunde diviſi laciniæque breviores, latiores; flores in racemis pauciores, ſegregati, dilute cærulei. In editioribus, apricis caulis brevior, minus ramoſus; foliorum lobi magis diviſi, laciniæ anguſtiores; racemus conſtat floribus congeſtis, ſaturatæ cæruleis.

PENTAGYNIA.

AQUILEGIA. Calyx nullus. Petala quinque. Nectaria quinque, corniculata, inter petala. Capſulæ quinque, diſtinctæ.

vulgaris. 1. A. nectariis incurvis petalo ſubæqualibus, foliolis omnibus petiolatis. *Hænke in Iacqu. collect. tom.* 2. *p.* 42. *Sp. pl. p.* 752. *Scop. carn. tom.* 1. *p.* 381.

Aquilegia foliis trilobatis; lobis petiolatis tripartitis, ſubrotundis, obtuſe dentatis. *Hall. hiſt. n.* 1195.

Aquilina. *Cluſ. hiſt. p.* CCIV.

L' Ancolie. Regn. bot.

In montanis, ſylvaticis. Fl. Majo, Iunio. Per.

viſcoſa. 2. A. nectariis incurvis, caule ſubnudo ſubunifloro viſcoſo-piloſo, foliolis ſubtrilobis. *Syſt. veget. p.* 505. *Gouan. fl. p.* 267. *Magn. bot.* 24. *hort.* 21.

Aquilegia alpina. *Hænke in Iacqu. collect. tom.* 2. *p.* 42.

In Carniolia in alpibus wochinenſibus. In Tyroli in alpibus leontinis. Fl. Iulio, Auguſto. Per.

Vel ſola pube viſcida, qua caulis, pedunculi, piſtilla veſtita ſunt, diſtinguitur ab A. vulgari; vel ſola radice perenni, ab A. alpina, cui radix biennis eſt.

NIGELLA. Calyx nullus. Petala quinque. Nectaria quinque, trifida, intra corollam. Capſulæ quinque, connexæ.

arvenſis. 1. N. piſtillis quinis, petalis integris, capſulis turbinatis. *Sp. pl. p.* 753. *Scop. carn. tom.* 1. *p.* 382.

Nigella flore nudo, ſiliquarum cornubus longiſſimis. *Hall. hiſt. n.* 1194.

La Nielle des champs. Bulliard. herb. t. 126.

In arvis, ad agrorum margines, vineas. Fl. æſtate.

POLIGYNIA.

ANEMONE. Calyx nullus. Petala duorum triumve ordinum, in ſingula ſerie tria. Semina plura, ſtylum retinentia.

* *Flore ſubcalyculato.*

Hepatica. 1. A. foliis trilobis integerrimis. *Sp. pl. p.* 758. *Fl. dan. t.* 610. *Scop. carn. tom.* 1. *p.* 382. *Hall. hiſt. n.* 1156.

In nemoribus. Fl. Martio, Aprili. Per.

** *Pedunculo involucrato, ſeminibus caudatis.*

vernalis. 2. A. pedunculo involucrato, foliis pinnatis, flore erecto. *Sp. pl. p.* 759. *Fl. dan. t.* 29. *Wulfen in Iacqu. collect. tom.* 3. *p.* 10.

Anemone tubis caudatis, involucris multifidis, foliis pinnatis, pinnis latis, lobatis. *Hall. hiſt. n.* 1147. *t.* 21.

Anemone tubis caudatis, involucris multifidis, foliis hirſutis, pinnatis, pinnis acute lobatis. *Hall. hiſt. n.* 1148.

Anemone ſulphurea. *Allion. pedem. tom.* 2. *p.* 170.

Anemone Halleri. *Allion. pedem. tom.* 2. *p.* 170. *t.* 80. *f.* 2.

In Carinthiæ alpibus reichenavienſibus. In Tyroli in alpibus leontinis, & in montibus *in Neuſtift im Gerichte Stubay.* Fl. poſt ſolutas nives. Per.

Pulſatilla. 3. A. pedunculo involucrato, petalis rectis, foliis bipinnatis. *Sp. pl. p.* 759. *Fl. dan. t.* 153. *Scop. carn. tom.* 1. *p.* 386.

Anemone tubis caudatis, involucris unifoliis multifidis, foliis linearibus hirſutis, duplicato-pinnatis, pinnis tripartitis & quinquepartitis. *Hall. hiſt. n.* 1146.

Pulſatilla vulgaris dilutiore flore. *Clus. hiſt. p.* 246.

L' Anemone Pulſaltille. Bulliard *herb. t.* 49.

In collibus, montoſis ſylvaticis. Fl. primo vere. Per.

pratenſis. 4. A. pedunculo involucrato, petalis apice reflexis, foliis bipinnatis. *Sp. pl. p.* 760. *Fl. dan. t.* 611.

Pulſatilla vulgaris ſaturatiore flore. *Clus. hiſt. p.* 246.

Pulſatilla flore minore nigricante. *Bauh. pin.* 177.

In pratis ſiccioribus, collibus herbidis, montoſis apricis. Fl. primo vere. Per.

alpina. 5. A. foliis caulinis ternis connatis ſupradecompoſitis multifidis, ſeminibus hirſutis caudatis. *Sp. pl. p.* 760. *Iacqu. auſtr. t.* 85.

Anemome foliis triplicato-pinnatis, acute circumſerratis, caulinis ternatis, tubis caudatis. *Hall. hiſt. n.* 1149.

Anemone burſeriana. *Scop. carn. tom.* 1. *p.* 285.

Anemone ſulphurea. *Syſt. veget. p.* 509.

Anemone apiifolia. *Syſt. veget. p.* 510. *Wulfen in Iacqu. miſcell. tom.* 2. *p.* 47. *t.* 4. *Scop. carn. tom.* 1. *p.* 285.

Anemone ſylveſtris ſecunda. *Clus hiſt. p.* 244.

Anemone ſylveſtris tertia. *Clus. hiſt. p.* 245.

Pulſatilla lutea. *Cam. epit.* 393. *Bauh. hiſt.* 3. *p.* 411.

In alpinis, & ſubalpinis. Fl. a Iunio ad Septembrim. Per.

Petala alba, per ſuccedentes gradus, in pallidum, citrinum & croceum colorem degenerant. Haller. l. c.

*** *Seminibus ecaudatis.*

hortenſis. 6. A. ſeminibus lanatis, foliis radicalibus trilobatis & multifidis, caulinis ovato-lanceolatis. *Hall. hiſt. n.* 1152. *Sp. pl. p.* 761. *Curtis magaz.* 123.

Anemone hortenſis latifolia ſimplici flore. *Clus. hiſt. p.* 249.

In agro fluminenſi in *Belvedere, Raſtozzine, Braida*, & alibi ad maceries, ſepes, in juniperetis. Fl. Februario, Martio. Per.

Radix tuberoſa, plantæ novellæ exiguæ olivæ magnitudine, adultæ forma & magnitudine varia, fibris paucis aucta. Folia dorſo purpuraſcentia, triloba, vel multifida. Caulis albido to-

mento veſtitus. Involucrum triphyllum: foliolis lanceolatis, vel integris, vel apice in duas tresvè lacinias diviſis. Caulem terminat flos unicus, conſtans petalis duodecim, vel pluribus, purpureis, raro albis, lineari-lanceolatis, dorſo villoſis; externis quinque longioribus. Semina lanata.

ſylveſtris. 7. A. foliis digitatis, hirſutis, floribus pentapetalis, ſeminibus lanatis. *Hall. hiſt. n.* 1150. *Sp. pl. p.* 761. *Curtis magaz.* 54.

Anemone ſylveſtris prima. *Cluſ. hiſt. p.* 244.

Anemone blanche. **Bulliard.** *herb. t.* 59.

In collibus, montoſis ſylvaticis. Fl. Majo. Per.

baldenſis. 8. A. ſeminibus lanatis, foliis caulinis ternatis, triplicato-trifidis, lobulis lanceolatis, acute trifidis bifidisve. *Hall. hiſt. n.* 1151. *Syſt. veget. p.* 509.

Anemone fragifera. *Wulfen in Iacqu. miscell. tom.* 2. *p.* 55. *ic. t.* 103. *Syſt. veget. p.* 510.

Anemone alpina. *Scop. carn. tom.* 1. *p.* 384. *t.* 26.

Anemone alpina alba minor. **Bauh.** *pin.* 176. *prodr.* 94.

In alpibus Styriæ, Carnioliæ, Croatiæ, Carinthiæ, Tyrolis, Carniæ. Fl. a Iunio ad Auguſtum. Per.

trifolia. 9. A. foliis ternatis ovatis integris ſerratis, caule unifloro. *Sp. pl. p.* 762. *Scop. carn. tom.* 1. *p.* 383.

Anemone trifolia, flore albo. **Bauh.** *hiſt.* 3. *p.* 412.

In Carnioliæ alpium convallibus, alibique in nemoribus. Fl. Majo. Per.

nemoroſa. 10. A. ſeminibus acutis, foliolis inciſis, caule unifloro. *Sp. pl. p.* 762. *Fl. dan. t.* 549. *Scop. carn. tom.* 1. *p.* 383.

Anemone ſeminibus nudis, caule unifloro, foliis caulinis quinquepartitis, lobis tripartitis inciſis. *Hall. hiſt.* 1154.

Ranunculus ſylvarum. *Clus. hiſt. p.* 247.

L' Anemone des Bois. Bulliard herb. t. 3.

In nemorenſibus, ſylvaticis. Fl. primo vere. Per.

ranunculoides. 11. A. ſeminibus acutis, foliolis inciſis, petalis ſubrotundis, caule ſubbifloro. *Sp. pl. p.* 762. *Fl. dan. t.* 140. *Scop. carn. tom.* 1. *p.* 384.

Anemone biflora, ſeminibus nudis, foliis caulinis quinquepartitis, lobis tripartitis inciſis. *Hall. hiſt. n.* 1153.

Habitat & floret cum præc. Per.

narciſſiflora. 12. A. floribus umbellatis, ſeminibus depreſſo-ovalibus nudis. *Sp. pl. p.* 763. *Iacqu. auſtr. t.* 159.

Anemone ſeminibus nudis, rotundis, planis, foliis ſeptifidis imbricatis, floribus umbellatis. *Hall. hiſt. n.* 1155.

Ranunculus montanus ſecundus, & Ranunculi montani ſecundi ſpecies altera. *Clus. hiſt. p.* 235.

In Auſtriæ, Styriæ ſubalpinis, & alpinis. Fl. a Iunio ad Auguſtum. Per.

ATRAGENE. Calyx tetraphyllus. Petala duodecim. Semina caudata.

alpina. 1. A. foliis duplicato-ternatis ſerratis, petalis exterioribus quaternis. *Sp. pl. p.* 764. *Iacqu. auſtr. t.* 241.

Atragene foliis ternatis, foliolis trifoliis acutis. *Hall. hiſt. n.* 1145.

Atragene auſtriaca. *Scop. carn. tom.* 1. *p.* 387.

Clematis cruciata alpina. *Pon. in Clus. hiſt. p.* CCCXXI.

Clematis alpina geranifolia. *Bauh. pin.* 300. *prodr.* 135.

In rupeſtribus, ſaxoſis alpinis, & ſubalpinis. Fl. Iulio, Auguſto. Fr.

CLEMATIS. Calyx nullus. Petala quatuor, rarius quinque vel ſex. Semina caudata.

Viticella. 1. C. foliis compoſitis decompoſitisque: foliolis ovatis ſublobatis integerrimis. *Syſt. veget. p.* 512. *Scop. carn. tom.* 1. *p.* 388.

Clematis altera. *Clus. hiſt. p.* 122.

Clematis cœrulea vel purpurea repens. *Bauh. pin.* 300.

Clematis ſive flammula flore cœruleo ſcandens. *Bauh. hiſt.* 2. *p.* 128. *Zann. iſt. p.* 69. *t.* 80.

In Forojulio ad ſepes agri medenſis. *Scop.* Ego Duini; Flumine *al Ponſal*, *San Nicoló*, & alibi. Fl. Majo, Iunio. Fr.

Vitalba. 2. C. foliis pinnatis, foliolis cordatis ſcandentibus. *Sp. pl. p.* 766. *Iacqu. auſtr. t.* 308. *Curtis lond. Scop. carn. tom.* 1. *p.* 388.

Clematis caule ſcandente, foliis pinnatis, ovato-lanceolatis. *Hall. hiſt. n.* 1142.

Atragene Theophraſti. *Clus. hiſt p.* 122.

La Clematite des Haies. Bulliard herb. t. 89.

Ad ſepes, alibique. Fl. Auguſto, Septembri. Per.

Flammula. 3. C. foliis inferioribus pinnatis ſcandentibus laciniatis: ſummis ſimplicibus integerrimis lanceolatis. *Sp. pl. p.* 766.

Clematis cespitoſa. *Scop. carn. tom.* 1. *p.* 389.

Clematis maritima. *Allion. nic.* 122.

Clematis maritima repens. *Bauh. pin.* 300. *prodr.* 135. *Zann. iſt. p.* 68. *t.* 129.

In Litorali ad ſepes, in fruticoſis agri fluminenſis, duinenſis, & alibi. Fl. Iunio, Iulio. Fr.

Radix emittit caules plures, ſarmentoſos, ramoſos, ſi frutices in vicinia, ſcandentes, ſi nulli, inter ſe et cum ramis intertextos, humi diffuſos, paucos breviores erectos. Folia ſumma ſunt ſimplicia, reliqua bipinnata, pinnata, & hisce mixta bigemina, ternataque: foliolis aliis lanceolatis, quibusdam ovatis, aliquibus linearibus; omnibus integris, aut bi- ſeu trifidis. Corolla conſtat petalis quatuor vel quinque, lanceolatis, lineatis, apice compreſſis. Stamina numeroſa. Piſtilla a quinque ad octo. Hanc C. maritimam repentem Bauhini eſſe, autopta teſtor. Quod C. Flammula ſit quoque C. maritima Allionii, candide in Auctario ſuo ad Floram pedemontanam ipſemet Cl. vir teſtatur dicens. „Deleatur C. maritima, que eſt vere Flammula. Ea, quam pro Flammula habui, varietas eſt C. erectæ.“

erecta. 4. C. foliis pinnatis: foliolis ovato-lanceolatis, integerrimis, caule erecto, floribus pentapetalis tetrapetalisque. *Sp. pl. p.* 767. *Iacqu. auſtr. t.* 291. *Scop. carn. tom.* 1. *p.* 387.

Clematis caule erecto, foliis pinnatis, ovato-lanceolatis. *Hall. hiſt. n.* 1144.

Flammula Matthioli. *Clus. hiſt. p.* 124.

Ad ſepes, in fruticoſis, montoſis ſylvaticis. Fl. æſtate. Per.

integrifolia. 5. C. foliis ſimplicibus ovato-lanceolatis, floribus cernuis. *Sp. pl. p.* 767. *Iacqu. auſtr. t.* 363. *Scop. carn. tom.* 1. *p.* 387.

Clematis cœrulea pannonica. *Clus. hiſt. p.* 123.

In pratis paluſtribus Duini, Aquilejæ. In ſimilibus locis in Auſtria circa *Bruck an der Leitha*; & in Pannonia ad lacum Peiſonis, alibique. Fl. Iunio, Iulio. Per.

THALICTRUM. Calyx nullus. Petala quatuor ſeu quinque. Semina ecaudata.

alpinum. 1. T. caule ſimpliciſſimo ſubnudo, racemo ſimplici terminali. *Sp. pl. p.* 767. *Fl. dan. t.* 11.

In Tyroli in alpibus leontinis. Per.

minus. 2. T. foliis ſexpartitis, floribus cernuis. *Sp. pl. p.* 769. *Iacqu. auſtr. t.* 419. *Scop. carn. tom.* 1. *p.* 390.

Thalictrum foliis patulis, imis obtuſe trilobis, ſupremis acutis, panicula diffuſa, floribus nutantibus. *Hall. hiſt. n.* 1139.

In collibus herbidis, montoſis apricis, & in ipſis alpibus. Fl. Iunio, Iulio. Per.

Folia multiplicato - pinnata: foliolis breviſſime petiolatis vel ſeſſilibus, venoſis, ad apicem bis terve inciſis aut integerrimis, obtuſis vel acutis. Piſtilla tria vel quatuor. Iacqu. l. c.

majus. 3. T. foliis multipartitis, floribus cernuis, ramis floriferis ex iisdem axillis binis vel ternis. *Syſt. veget. p.* 513. *Iacqu. auſtr. t.* 420.

In Auſtriæ, Pannoniæ ſylvaticis. Fl. Iunio, Iulio. Per.

In majori piſtilla quatuor ad ſeptem; ex eodem caulis & ramorum principum nodo bini vel terni egrediuntur rami, qui in minori fere ſemper ſolitarii. Foliorum prona facies in majori magis ſaturate veluti ex atro viret, poſtica autem intenſe & ex albo glauca eſt; dum illa in minori viret lætius, hæc vero pallet. Iacqu. l. c.

anguſtifolium. 4. T. foliolis lanceolato - linearibus integerrimis. *Sp. pl. p.* 769. *Iacqu. hort. tom.* 3. *p.* 25. *t.* 43. *Scop. carn. tom.* 1. *p.* 391.

Thalictrum foliis aſcendentibus, linearibus, panicula erecta. *Hall. hiſt. n.* 1137.

Thalictrum pratenſe anguſtiſſimo folio. *Bauh. pin.* 337. *prodr.* 146.

In Auſtriæ, Styriæ, Carnioliæ pratis udis. Fl. Iunio, Iulio. Per.

flavum. 5. T. caule folioſo ſulcato ; panicula multiplici erecta. *Sp. pl. p.* 770.

Thalictrum foliis erectis, lobulis acute trilobis, ſpica adſcendente. *Hall. hiſt. n.* 1138.

Thalictrum nigricans. *Iacqu. auſtr. t.* 421. *auct. Cl. Sibthorp.*

In Auſtria, Pannonia in inſulis danubialibus, pratis paluſtribus. Fl. Iunio. Per.

Radix repens. Foliola lanceolata, acuta, facie ſaturate virentia, ſubtus pallidiora veniſque rugoſula, utrinque glabra ; alia integra, alia ter vel bis inciſa, idque inprimis in foliis inferioribus, ubi etiam latiora ſunt quam in ſuperioribus. Flores erecti. Piſtilla a duobus ad quinque. Iacqu. l. c.

aquilegifolium. 6. T. fructibus pendulis triangularibus rectis, caule tereti. *Sp. pl. p.* 770. *Iacqu. auſtr. t.* 318. *Scop. carn. tom.* 1. *p.* 389.

Thalictrum foliis triplicato-pinnatis, patulis, obtuſe trilobatis, ſtipulis ovatis, paniculis umbellatis. *Hall. hiſt. n.* 1141.

In alpinis, & ſubalpinis. Fl. æſtate. Per.

Caulis fiſtuloſus. Folia triternata, glabra, ad exortum & diviſiones munita ſtipulis rotundatis, perfoliatis. Foliola ſunt fere ſubrotunda, ſuperne obtuſe dentato-inciſa, ſubtus glauca, venuloſa. Petala quatuor, obverſe ovata, concava, ſtaminibus breviora. Flores paniculati. Stamina plurima. Piſtilla ſeptem vel octo. Capſulæ pendulæ, triangulares, trialatæ. Iacqu. l. c.

ADONIS. Calyx pentaphyllus. Petala absque nectario, quinque vel plura. Semina nuda.

autumnalis. 1. A. floribus octopetalis, fructibus ſubcylindricis. *Sp. pl. p.* 771.

Adonis autumnalis, corollis flavo-rubris. *Scop. carn. tom.* 1. *p.* 392.

Adonis miniata. *Iacqu. auſtr. t.* 354.

Inter ſegetes, in arvis reſtibilibus. Fl. Majo, Iunio.

Petala ſunt minii colore tincta. Semel tamen in Auſtriæ arvis prope Bertholdsdorf *inveni varietatem corollis citrinis.*

flammea. 2. A. floribus octopetalis, calycibus hirſutis, fructibus cylindricis. *Syſt. veget. p.* 514. *Iacqu. auſtr. t.* 355.

Adonis autumnalis, floribus phœniceis. *Scop. carn. tom.* 1. *p.* 392.

Cum præcedente. Fl. æſtate.

Habitus idem cum autumnali. Radix quaſi inſipida, cum amara ſit autumnali. Folia multifida, ut in illa, at laciniæ ſunt anguſtiores & longiores. In præcedente calyces ſunt glaberrimi, qui hirſutuli ſunt in flammea. Petala ſunt anguſtiora, quam in autumnali. Iacqu. l. c.

vernalis. 3. A. flore dodecapetalo, fructu ovato. *Sp. pl. p.* 771. *Scop. carn. tom.* 1. *p.* 392. *Hall. hiſt. n.* 1157.

Adonis apenina. *Iacqu. auſtr. t.* 44.

Pſeudoelleborus niger. *Cluſ. hiſt. p.* 333.

In Auſtriæ, Pannoniæ collibus, montoſis ſylvaticis. Fl. primo vere. Per.

Variat ramis floriferis, & ſterilibus.

RANUNCULUS. Calyx pentaphyllus. Petala intra ungues poro mellifero. Semina nuda.

Flammula. 1. R. foliis ovato-lanceolatis petiolatis, caule declinato. *Sp. pl. p.* 772. *Fl. dan. t.* 572. *Scop. carn. tom.* 1. *p.* 394.

Ranunculus caule declinato, foliis elliptico-lanceolatis, ſubſerratis. *Hall. hiſt. n.* 1182.

La petite Douve. Bulliard. herb. t. 15.

In udis. Fl. æſtate. Per.

Etiamſi ſæpe e geniculis radices agat, rependoque ſe multiplicet; male tamen a nonnullis pro varietate R. reptantis habetur. Retzii ſcand. p. 130.

Lingua. 2. R. foliis lanceolatis, caule erecto. *Sp. pl. p.* 773. *Fl. dan. t.* 754. *Scop. carn. tom.* 1. *p.* 395.

Ranunculus erectus, ramoſus, foliis enſiformibus ſubſerratis. *Hall. hiſt. n.* 1181.

In Auſtriæ, Pannoniæ, Carnioliæ aquis ſtagnantibus. Fl. Iulio, Auguſto Per.

pyrenæus. 3. R. foliis linearibus indiviſis, caule erecto ſtriato ſubbifloro. *Syſt. veget. p.* 515. *Wulfen in Iacqu. miscell. tom.* 1. *p.* 154. *t.* 18. *f.* 1.

Ranunculus radicibus fasciculatis; imo caule reticulato; foliis gramineis. *Hall. hiſt. n.* 1180.

In Tyroli in pratis alpium leontinarum. *Wulfen.* Per.

parnaſſifolius. 4. R. foliis ſubovatis nervoſis lineatis integerrimis petiolatis. *Sp. pl. p.* 774. *Wulfen in Iacqu. collect. tom.* 1. *p.* 191. *t.* 9. *f.* 3.

Ranunculus radicibus fasciculatis, imo caule reticulato, foliis cordiformibus. *Hall. hiſt. n.* 1179.

In Tyroli in alpibus leontinis. Per.

Ficaria. 5. R. foliis cordatis angulatis petiolatis, caule unifloro polypetalo. *Retzii ſcand. p.* 130. *Sp. pl. p.* 774. *Fl. dan. t.* 499. *Curtis lond. Scop. carn. tom.* 1. *p.* 395.

Caltha paluſtris vulgaris. *Clus. hiſt. p. CXIV.*

L' Eclairette. Regn. bot.

In cultis, pratis, ſylvis, & alibi. Fl. Aprili. Per.

Thora. 6. R. foliis reniforimbus ſubtrilobis crenatis: caulino ſeſſili, floralibus lanceolatis, caule ſubbifloro. *Sp. pl. p.* 775. *Iacqu. auſtr. t.* 442. *Scop. carn. tom.* 1. *p.* 395.

Ranunculus foliis duris, venoſis, reniformibus, rotunde crenatis, convexitate retuſa, acute ſerrata. *Hall. hiſt. n.* 1178.

Thora. Valdenſium, & Ranunculus grumoſa radice tertius. *Clus. hiſt. p.* 239.

In alpibus, Fl. æſtate. Per.

auricomus. 7. R. foliis radicalibus reniformibus crenatis inciſis; caulinis digitato-linearibus, caule multifloro. *Sp. pl. p.* 775. *Fl. dan. t.* 665. *Curtis lond. Scop. carn. tom.* 1. *p.* 397.

Ranunculus foliis radicalibus integris & ſemitrilobatis, rotunde crenatis, caulinis multipartitis, linearibus integerrimis. *Hall. hiſt. n.* 1177.

In pratis, montanis ſylvaticis. Fl. Majo. Per.

ſceleratus. 8. R. foliis inferioribus palmatis: ſummis digitatis, fructibus oblongis. *Sp. pl. p.* 776. *Curtis lond. Scop. carn. tom.* 1. *p.* 398.

Ranunculus foliis lævibus, ſemitrilobatis, rotunde ſerratis, fructu ovato. *Hall. hiſt. n.* 1175.

La Renoncule des Marais. Regn. bot.

Ad foſſas, lacus, in uliginoſis, paluſtribus. Fl. Majo, Iunio.

aconitifolius. 9. R. caule ramoſo multifloro, foliis venoſis, quinquelobis rhomboideis, acutis, acute ſerratis. *Hall. hiſt. n.* 1164. *Sp. pl. p.* 776. *Socp. carn. tom.* 1. *p.* 393.

Ranunculus montanus quartus. *Clus. hiſt. p.* 236.

In alpinis, & ſubalpinis. Fl. a Iunio ad Auguſtum. Per.

illyricus. 10. R. foliis ternatis integerrimis lanceolatis. *Sp. pl. p.* 776. *Iacqu. auſtr. t.* 222.

Ranunculus grumoſa radice quartus. *Clus. hiſt. p.* 239.

In Auſtria, Pannonia in herbidis, ad agrorum margines. Fl. Majo, Iunio. Per.

rutæfolius. 11. R. foliis ſupradecompoſitis, caule ſimpliciſſimo unifolio, unifloro, radice tuberoſa. *Sp. pl. p.* 777. *Wulfen in Iacqu. collect. tom.* 1. *p.* 186. *t.* 6. 7.

Ranunculus caule unifloro, foliis imbricatis pinnatis, pinnis lobatis, lobis obtuſe palmatis. *Hall. hiſt. n.* 1165.

Ranunculus præcox rutæfolius. *Clus hiſt. p.* 232.

In alpibus. Per.

glacialis. 12. R. calycibus hirſutis, caule bifloro, foliis multifidis. *Sp. pl. p.* 777. *Fl. dan. t.* 19. *Wulfen in Iacqu. collect. tom.* 1. *p.* 189. *t.* 8. 9. *f.* 1. 2.

Ranunculus caule bifloro, calyce hirſuto. *Fl. lapp. n.* 233. *t.* 3. *f.* 1.

Ranunculus foliis pulpoſis, repetito trilobatis, calycibus villoſis. *Hall. hiſt. n.* 1166.

In Tyroli in vicinia glaciei. Fl. Iulio, Auguſto. Per.

nivalis. 13. R. calyce hirſuto, caule unifloro, foliis radicalibus palmatis: caulinis multipartitis ſeſſilibus. *Sp. pl. p.* 778. *Iacqu. auſtr. t.* 325. 326.

Ranunculus caule unifloro, foliis glabris, reniformibus, ſemiquinquelobis. *Hall. hiſt. n.* 1168.

In alpibus. Fl. Iulio, Auguſto. Per.

alpeſtris. 14. R. foliis radicalibus ſubcordatis obtuſis tripartitis: lobis trilobatis: caulino lanceolato integerrimo, caule ſubunifloro. *Sp. pl. p.* 778. *Iacqu. auſtr. t.* 110. *Scop. carn. tom.* 1. *p.* 393.

Ranunculus uniflorus, foliis lævibus, ſemitrilobis, rotunde ſerratis, caulinis lingulatis. *Hall. hiſt. n.* 1165.

Ranunculi montani primi ſpecies prima, & altera. *Clus. hiſt. p.* 234.

In alpibus. Fl. Iulio, Auguſto. Per.

bulboſus. 15. R. calycibus retroflexis, pedunculis ſulcatis, caule erecto multifloro, foliis compoſitis. *Sp. pl. p.* 778. *Curtis lond. Fl. dan. t.* 551. *Scop. carn. tom.* 1. *p.* 400.

Ranunculus radice ſubrotunda, foliis hirſutis, ſemiſeptilobis, lobis petiolatis. *Hall. hiſt. n.* 1174.

La Renoncule bulbeuſe. Bulliard herb. t. 27.

In pratis, collibus, montanis ſylvaticis, & alibi. Fl. Majo, Iunio. Per.

repens. 16. R. calycibus patulis, pedunculis ſulcatis, ſtolonibus repentibus, foliis compoſitis. *Sp. pl. p.* 779. *Curtis lond. Fl. dan. t.* 795. *Scop. carn. tom.* 1. *p.* 398.

Ranunculus caule reptante radicato, foliis ſemitrilobatis, lobis petiolatis. *Hall. hiſt. n.* 1173.

La Renoncule rampante. Bulliard herb. t. 77.

In cultis, ruderatis, pratis, ad fluvios. Fl. æſtate. Per.

acris. 17. R. calycibus patulis pedunculis teretibus, foliis tripartito-multifidis: ſummis linearibus. *Sp. pl. p.* 779. *Curtis lond. Scop. carn. tom.* 1. *p.* 398.

Ranunculus foliis hirſutis ſemitrilobatis, lobis lateralibus bipartitis, foliis caulinis ſemitrilobis. *Hall. hiſt. n.* 1169.

La Renoncule acre. Bulliard. herb. t. 109.

In pratis, & aliis herbidis locis. Fl. æſtate. Per.

lanuginoſus. 18. R. calycibus patulis; pedunculis teretibus, caule petioliſque hirtis, foliis trifidis lobatis crenatis holoſericeis. *Sp. pl. p.* 779. *Fl. dan t.* 397. *Scop. carn. tom.* 1. *p.* 399.

Ranunculus folioſus, ramoſus, erectus, foliis ſemitrilobatis, hirſutis, vaginis ſericeis. *Hall. hiſt. n.* 1172.

In ſylvis. Fl. æſtate. Per.

arvenſis. 19. R. ſeminibus aculeatis, foliis ſuperioribus decompoſitis linearibus. *Sp. pl. p.* 780. *Fl. dan. t.* 219. *Scop. carn. tom.* 1. *p.* 400.

Ranunculus ſeminibus aculeatis, foliis tripartitis, lobis longe petiolatis, bipartitis & tripartitis acute inciſis. *Hall. hiſt. n.* 1176.

La Renoncule des Champs. Bulliard. herb. t. 117.

In arvis. Fl. Majo, Iunio.

murica-tus. 20. R. ſeminibus aculeatis, foliis ſimplicibus lobatis obtuſis glabris, caule diffuſo. *Sp. pl. p.* 780.

Ranunculus Apulei quibusdam. *Clus. hiſt. p.* 233.

Flumine in udis, & vineis. Fl. æſtate.

falca-tus. 21. R. foliis filiformi-ramoſis, ſcapo nudo unifloro, ſeminibus falcatis. *Sp. pl. p.* 781. *Iacqu. auſtr. t.* 48.

In agri vindobonenſis aggeribus, arvis, ruderatis, ad margines agrorum. Fl. Martio, Aprili.

aquati-lis. 22. R. foliis ſubmerſis capillaceis, emerſis peltatis. *Sp. pl. p.* 781. *Fl. dan. t.* 376. *Scop. carn. tom.* 1. *p.* 394.

In rivulis, fluviis, lacubus, foſſis. Fl. æſtate. Per.

Variat foliis omnibus capillaceis.

TROLLIUS. Calyx nullus. Petala circiter quatuordecim. Nectaria novem, plana, baſi introrſum perforata. Capſulæ plurimæ, ovatæ, polyſpermæ.

europæ-us. 1. T. corollis conniventibus, nectariis longitudine ſtaminum. *Sp. pl. p.* 782. *Fl. dan. t.* 133. *Scop. carn. tom.* 1. *p.* 401. *Hall. hiſt. n.* 1189.

Ranunculus glomerato flore. *Clus. hiſt. p.* 237.

Ranunculus montanus aconiti folio, flore globoſo. *Bauh. pin.* 182.

In pratis demiſſis, ſubalpinis, & alpinis. Fl. a Majo ad Auguſtum. Per.

ISOPYRUM. Calyx nullus. Petala quinque. Nectaria trifida, tubulata. Capſulæ recurvæ, polyſpermæ.

thalictroides. 1. I. ſtipulis ovatis, petalis obtuſis. *Sp. pl. p.* 783. *Iacqu. auſtr. t.* 105. *Scop. carn. tom.* 1. *p.* 402.

Ranunculus præcox alter, Thalictri folio. *Cluſ. hiſt. p.* 233.

In umbroſis Auſtriæ, Pannoniæ, Carnioliæ, Carniæ. Fl. Aprili, Majo. Per.

Radix conſtat faſciculo fibrillarum teretium, ex quo poſtea elongatur fibrilla alia, horizontaliter pergens, quæ in decurſu ſimiles radicularum faſciculos deorſum terræ immittit, ex hinc iterum caulifera, atque ſic perennando ſe propagans. Folium radicale bis trifidum: foliolis obtuſis, apice bilobis vel trilobis. Caulina plerumque ſimilia. Summa ternata aut lobata. Germina ab uno ad tria, nunquam plura. Nectarii foliola receptaculo ante petala inſeruntur. Iacqu. l. c.

HELLEBORUS. Calyx nullus. Petala quinque ſeu plura. Nectaria bilabiata. Capſulæ polyſpermæ, erectiuſculæ.

hyemalis. 1. H. flore folio inſidente. *Sp. pl. p.* 783. *Iacqu. auſtr. t.* 202. *Hall. hiſt. n.* 1191.

Aconitum luteum minus. *Dod. purg. p.* 312.

L' Hellebore d' hiver. Bulliard herb. t. 35.

In agro vindobonenſi in montibus, qui inter *Hädersdorf* & *Kahlenberg* interjacent. *Iacqu.* In Carniolia in monte *Nanas. I. Hoſt.* Ego in Litorali circa *Moſchienizze.* Fl. Februario, Martio, Aprili. Per.

E radice tuberoſa exeunt folia gemina, alterum florigerum; ſterile alterum. Petala decidua. Iacqu. l. c.

niger. 2. H. ſcapo ſubbifloro ſubnudo, foliis pedatis. *Syſt. veget. p.* 519. *Iacqu. auſtr. t.* 201. *Scop. carn. tom.* 1. *p.* 403.

Helleborus legitimus. *Clus. hiſt. p.* 274.

L' Ellebore noir. Regn. bot.

In ſylvis ſubalpinis, montanis. Fl. Februario, Martio, Aprili. Per.

Nectariorum labium ſuperius emarginatum, inferius crenulatum. Germina quatuor vel quinque. Iacqu. l. c.

viridis. 3. caule bifido, ramis folioſis bifloris, foliis digitatis. *Sp. pl. p.* 784. *Iacqu. auſtr. t.* 106. *Scop. carn. tom.* 1. *p.* 403.

Veratrum nigrum alterum Dodonæi. *Clus. hiſt. p.* 275.

L' Ellebore a fleur verte. Regn. bot.

In Auſtriæ ſylvaticis prope *Hütteldorf, Mauerbach, Lilienfeld.* In petroſis aſperis Carnioliæ, Forijulii, Litoralis, Tyrolis calidioris. Fl. primo vere. Per.

CALTHA. Calyx nullus. Petala quinque. Nectaria nulla. Capſulæ plures, polyſpermæ.

paluſtris. 1. Caltha. *Sp. pl. p.* 784. *Curtis lond. Fl. dan. t.* 668. *Scop. carn. tom.* 1. *p.* 404. *Hall. hiſt. n.* 1188.

In udis. Fl. primo vere. Per.

DIDYNAMIA.

GYMNOSPERMIA.

AIUGA. Corollæ labium ſuperius minimum. Stamina labio ſuperiore longiora.

pyramidalis. 1. A. tetragono-pyramidalis villoſa, foliis radicalibus maximis. *Syſt. veget. p.* 525. *Fl. dan. t.* 185. *Retzii ſcand. p.* 139.

Bugula genvevenſis. *Scop. carn. tom.* 1. *p.* 416.

In montanis Carnioliæ. Fl. primo vere.

alpina. 2. A. caule ſimplici, foliis caulinis radicalia æquantibus. *Syſt. veget. p.* 525.

Bugula carneo flore. *Clus. hiſt. p. XLIII.*

In montanis, pratisque ſiccioribus Auſtriæ, Pannoniæ. Fl. Majo, Iunio. Per.

Variat floribus carneis & cœruleis.

reptans. 3. A. glabra ſtolonibus reptantibus. *Syſt. veget. p.* 525. *Curtis lond.*

Bugula flagellifera, foliis ovatis dentatis. *Hall. hiſt. n.* 282.

Bugula reptans. *Scop. carn. tom.* 1. *p.* 416.

Conſolida media pratenſis cœrulea. *Bauh. pin.* 260.

La Bugle, ou petite Conſoude. Regn. bot.

In ſylvis, pratis. Fl. Majo, Iunio. Per.

Variat floribus albis & cœruleis.

TEUCRIUM. Corollæ labium ſuperius ultra baſin diviſum: laciniis ad latera diſtantibus, referens corollam labio ſuperiore deſtitutam. Filamenta cum ſtylo inter diviſuram labii ſuperioris adſcendentia, prominentia.

Botrys. 1. T. foliis multifidis, floribus lateralibus ternis pedunculatis. *Syſt. veget. p.* 525. *Scop. carn. tom.* 1. *p.* 420.

Chamædrys foliis pinnatis, pinnis dentatis trilobatis, verticillis paucifloris. *Hall. hiſt. n.* 289.

Iva moſchata folio multifido. *Riv. mon. t.* 14.

In ſaxoſis, aſperis Auſtriæ, Carnioliæ calidioris, Litoralis. Fl. Iunio, Iulio.

Chamæpitys. 2. T. foliis trifidis linearibus integerrimis, floribus ſeſſilibus lateralibus ſolitariis, caule diffuſo. *Sp. pl. p.* 787. *Fl. dan. t.* 737.

Bugula foliis imis linearibus, caulinis tripartitis. *Hall. hiſt. n.* 284.

Bugula Chamæpitys. *Scop. carn. tom.* 1. *p.* 417.

L' Ivette. Regn. bot.

In arvis. Fl. æſtate.

Scorodonia. 3. T. foliis cordatis ſerratis petiolatis, racemis lateralibus ſecundis, caule erecto. *Sp. pl. p.* 789. *Curtis lond. Fl. dan. t.* 485. *Scop. carn. tom.* 1. *p.* 419.

Chamædrys foliis cordatis productis, ſpicis longiſſimis nudis, heteromallis. *Hall. hiſt. n.* 287.

Scordium alterum ſeu Salvia ſylveſtris. *Bauh. pin.* 247.

La Sauge des Bois. Regn. bot.

In ſylvis Carnioliæ, Croatiæ. Fl. Iunio, Iulio. Per.

Calycis lacinia ſuperior eſt maxima: hinc fit quod calyx bilabiatus appareat.

Scordium. 4. T. foliis oblongis ſeſſilibus dentato-ſerratis, floribus geminis axillaribus pedunculatis, caule diffuſo. *Sp. pl. p.* 790. *Fl. dan. t.* 593. *Scop. carn. tom.* 1. *p.* 418.

Chamædrys foliis mollibus, hirſutis, ellipticis, crenatis, verticillis paucifloris. *Hall. hiſt. n.* 288.

La Germandrée aquatique. Bulliard. herb. t. 205.

In paluſtribus, ad foſſas. Fl. æſtate. Per.

Chamædrys. 5. T. foliis cuneiformi-ovatis inciſis crenatis petiolatis, floribus ternis, caulibus procumbentibus ſubpiloſis. *Syſt. veget. p.* 527. *Scop. carn. tom.* 1. *p.* 418.

Chamædrys foliis firmis, ovatis, crenatis, verticillis paucifloris. *Hall. hiſt. n.* 286.

Chamædrys vulgaris, ſive ſecunda. *Clus. hiſt. p.* 351.

La Germandrée, ou petit Chêne. Regn. bot.

In collibus, montoſis apricis aſperis, ad vias, alibique. Fl. Iulio, Auguſto. Per.

flavum. 6. T. foliis cordatis obtuſe ſerratis: floralibus integerrimis concavis, caule fruticoſo, floribus ternis. *Sp. pl. p.* 791. *Scop. carn. tom.* 1. *p.* 419.

Teucrium vulgare fruticans primum. *Clus. hiſt. p.* 348.

In ſaxoſis, aſperis Litoralis, præcipue ad vinearum ſepes. Fl. Iulio.

Folia ſunt facie glabra, dorſo villoſula. Ex axillis bracteärum oppoſitarum prodeunt flores modo terni, modo quaterni, quandoque quini, raro plures.

montanum. 7. T. corymbo terminali, foliis lineari-lanceolatis integerrimis ſubtus tomentoſis. *Sp. pl. p.* 791. *Scop. carn. tom.* 1. *p.* 419.

Chamædrys foliis ellipticis integerrimis, contractis, ſubtus incanis, floribus capitatis. *Hall. hiſt. n.* 285.

Teucrium ſupinum. *Sp. pl. p.* 791. *Iacqu. auſtr. t.* 417.

Polium ſeptimum cum flore. *Clus. hiſt. p.* 363.

Polium pannonicum ſeptimum cum ſemine. *Clus. hiſt. p.* 363.

In apricis, ſaxoſis. Fl. æſtate. Per.

Folia ſunt integerrima, facie viridia & glabra, dorſo villoſo-incana, alia, præſertim juniora, margine revoluta, & linearia; alia margine plana, hæc adultiora tantum, & lanceolato-linearia.

Polium. 8. T. capitulis ſubrotundis, foliis oblongis obtuſis crenatis tomentoſis ſeſſilibus, caule proſtrato. *Sp. pl. p.* 792. *Scop. carn. tom.* 1. *p.* 420.

Polium montanum primum. *Clus. hiſt. p.* 361.

Polium maritimum ſupinum venetum. *Bauh. pin.* 221. *Zann. iſt. p.* 213. *t.* 35.

Le Thim blanc des montagnes, ou le Polion. Regn. bot.

In petroſis Litoralis circa *Moſchienizze*, *Berſez*, *Tergeſtum*, copioſiſſime in inſulis venetis. Fl. æſtate. Per.

Caules proſtrati, rami alii erecti, alii adſcendentes. Folia oppoſita, lineari-lanceolata, tomentoſa, crenulata, margine revoluta, pauca plana. Flores capitati: capitulis quibusdam ſeſſilibus, pedunculatis aliis.

SATUREIA. Perianthium quinquedentatum. Corollæ laciniæ ſubæquales. Stamina quatuor diſtantia.

montana. 1. S. pedunculis dichotomis lateralibus ſolitariis, foliis lineari-lanceolatis mucronatis. *Sp. pl. p.* 794. *Scop. carn. tom.* 1. *p.* 428. *t.* 30.

Satureja perennis. *Riv. mon.* 41.

In Carniolia, Croatiaque calidiore, & in toto Litorali, Tyrolique vinifero. Fl. a Iulio, ad Octobrim. Per.

rupeſtris. 2. S. pedunculis ramoſis lateralibus ſolitariis, foliis ovato - lanceolatis ſerrulatis. *Wulfen in Iacqu. collect. tom. 2. p. 130. ic. t. 494.*

Satureja thymifolia. *Scop. carn. tom. 1. p. 428. t. 29.*

In rupeſtribus Carnioliæ circa *Idriam*, prope *Haiduſchina.* In ſimilibus locis in Forojulio, & Carnia. Fl. cum præc. Per.

Folia ſunt oppoſita, ovato - lanceolata, utrinque punctata, apicem verſus denticulata. Calyx parum incurvus, ſtryatus, quinquedentatus: dentibus ſuperioribus tribus brevioribus approximatis; duobus inferioribus longioribus diſtantibus, omnibus margine aculeato - ciliatis. Corolla ex purpureo & albo variegata. Tubus incurvus. Labium utrumque margine reflexum. Erronee a quibusdam ſub duplici nomine: Saturejæ nempe rupeſtris, & S. thymifoliæ proponitur. Erronee ab aliis ut varietas Saturejæ Thymbræ, Tergeſti minime creſcentis, conſideratur.

HYSSOPUS. Corollæ labium ſuperius emarginatum: inferius trifidum; lacinia intermedia crenata, obcordata. Stamina recta, diſtantia.

officinalis. 1. S. ſpicis ſecundis, foliis lanceolatis. *Syſt. veget. p. 529. Iacqu. auſtr. t. 254.*

Hyſſopus foliis linearibus, punctatis, verticillis in ſpicam continuatis. *Hall. hiſt. n. 249.*

L' Hyſſope. Regn. bot.

In Carniolia ad pedem montis *Nanas*, in *Lipizza.* In agro fluminenſi ad radicem montium *mala Planina* dictorum, in monte majore, alibique locis herbidis, apricis, ſaxoſis. Fl. a Iunio ad Auguſtum. Per.

Caules oriuntur ex eadem radice plures, vix ſemipedales, quidam erecti, alii aſcendentes, pauci proſtrati. Folia ſunt lineari-lanceolata, acuta, punctata, aculeis ciliata. Corollæ vel rubræ, vel cœruleæ. Aegre ex habitu diſtinguitur a Satureja montana, cum qua ſemper creſcit. Et varietas Saturejæ montanæ, floribus rubris, magisque aggregatis, de qua Cl. Scopoli l. c. loquitur, mihi nil aliud quam Hyſſopus officinalis eſſe videtur.

NEPETA. Corollæ labium inferius lacinula intermedia crenata: faux margine reflexo. Stamina approximata.

Cataria. 1. N. floribus ſpicatis, verticillis ſubpedicellatis, foliis petiolatis cordatis dentato-ſerratis. *Sp. pl. p.* 796. *Fl. dan. t.* 580. *Scop. carn. tom.* 1. *p.* 430.

Cataria foliis cordatis, verticillis ſpicatis. *Hall. hiſt. n.* 246.

La Chataire commune. Bulliard. herb. t. 287.

Ad pagos, vias, in ruderatis, & alibi. Fl. æſtate. Per.

Occurrit in agro vindobonenſi varietas, odorem, odori meliſſæ officinalis valde ſimilem, ſpargens.

pannonica. 2. N. floribus panniculatis, foliis cordatis petiolatis obſolete crenatis. *Sp. pl. p.* 797.

Nepeta nuda. *Iacqu. auſtr. t.* 24.

In aſperis Pannoniæ, Auſtriæ. Fl. Iunio, Iulio. Per.

nuda. 3. N. florum racemis verticillatis nudis, foliis cordato-oblongis ſeſſilibus ſerratis. *Syſt. veget. p.* 529.

Nepeta pannonica. *Iacqu. auſtr. t.* 129.

In ceſpitoſis, ad ſepes, vias, ſylvarum margines Auſtriæ, Pannoniæ. Fl. cum præc. Per.

VERBENA. Calyx quinquedentatus: denticulo unico minimo. Corolla inæqualis. Tubus cylindraceus, incurvatus. Filamenta quatuor breviſſima. Antheræ duæ reliquis breviores. Semina quatuor.

officinalis. 1. V. ſpicis filiformibus paniculatis, foliis multifido-laciniatis, caule ſolitario. *Sp. pl. p.* 29. *Curtis lond. Fl. dan. t.* 628. *Scop. carn. tom.* 1. *p.* 433.

Verbena foliis rugoſis, tripartitis, ſpicis nudis, ſtrigoſis. *Hall. hiſt. n.* 219.

Verbena vulgaris. *Clus. hiſt. p. XLV.*

La Verveine. Regn. bot.

In pratis, ruderatis, ad pagos, vias. Fl. a Iulio ad Octobrim. Per.

SIDERITIS. Stamina intra tubum corollæ. Stigma brevius involvens alterum.

montana. 1. S. herbacea ebracteata, calycibus corolla majoribus ſpinoſis: labio ſuperiore trifido. *Syſt. veget. p.* 531. *Iacqu. auſtr. t.* 434.

Sideritis ſexta pannonica. *Clus. hiſt. p. XLI.*

Sideritis montana, parvo varioque flore. *Bauh. pin.* 233.

In arvis, locis aſperis Auſtriæ, Pannoniæ. Fl. æſtate.

romana. 2. S. herbacea ebracteata, calycibus ſpinoſis: lacinia ſuperiore majore ovata. *Sp. pl. p.* 802.

Sideritis, verticillis ſpinoſis, minor procumbens. *Moris. hiſt.* 3. *p.* 388. *ſect.* 11. *t.* 12. *f.* 5.

Tergeſti. *I. Hoſt.*

Calycis lacinia ſuperior eſt ovata, reliquiſque duplo latior. Corollæ labium ſuperius emarginatum. Linn. l. c.

MENTHA. Corolla ſubæqualis, quadrifida: lacinia latiore emarginata. Stamina erecta, diſtantia.

ſylveſtris. 1. M. ſpicis oblongis, foliis oblongis tomentoſis ſerratis ſeſſilibus, ſtaminibus corolla longioribus. *Sp. pl. p.* 804. *Fl. dan. t.* 484.

Mentha foliis elliptico-lanceolatis, ſerratis, ſubtus tomentoſis, ſpicis cylindricis, ſtaminibus flore duplo longioribus. *Hall. hiſt. n.* 227.

Ad foſſas, fluvios, in pratis paluſtribus, alibique. Fl. Iulio, Auguſto. Per.

viridis. 2. M. ſpicis oblongis, foliis lanceolatis nudis ſerratis ſeſſilibus, ſtaminibus corolla longioribus. *Sp. pl. p.* 804.

Mentha foliis elliptico-lanceolatis glabris, ſpicis cylindricis. *Hall. hiſt. n.* 229.

La Menthe a Epi. Regn. bot.

Occurrit ad pagos, rivulos. Fl. cum præc. Per.

aquatica. 3. M. floribus capitatis, foliis ovatis ſerratis petiolatis, ſtaminibus corolla longioribus. *Sp. pl. p.* 805. *Fl. dan. t.* 673. *Scop. carn. tom.* 1. *p.* 432.

Mentha foliis ovato-lanceolatis, ſerratis, verticillis pauciſſimis, teminante maximo. *Hall. hiſt. n.* 225.

In paluſtribus, aquoſis. Fl. Iulio. Per.

ſativa. 4. M. floribus verticillatis, foliis ovatis acutiuſculis ſerratis, ſtaminibus corolla longioribus. *Sp. pl. p.* 805. *Fl. dan. t.* 794.

In humidis, ad pagos in Tyroli circa Oenopontum, & Halam ad Oenum. Infrequens in Auſtria ad rivulos. Fl. Auguſto, Septembri. Per.

arvenſis. 5. M. floribus verticillatis, foliis ovatis acutis ſerratis, ſtaminibus corollam æquantibus. *Sp. pl. p.* 806. *Fl. dan. t.* 512. *Scop. carn. tom.* 1. *p.* 432.

Mentha floribus verticillatis, foliis hirſutis, ovato-lanceolatis, verticillis tomentoſis. *Hall. hiſt. n.* 223.

In arvis, paluſtribus. Fl. Iulio, Auguſto. Per.

auſtriaca. 6. M. floribus verticillatis, corollæ laciniis obtuſis, foliis ſubovatis villoſis, ſtaminibus corolla brevioribus. *Syſt. veget. p.* 533. *Iacqu. auſtr. t.* 430.

In Auſtria in arenoſis ad Danubii ripas. Fl. cum præc. Per.

Pulegium. 7. M. floribus verticillatis, foliis ovatis obtuſis ſubcrenatis, caulibus ſubteretibus, repentibus, ſtaminibus corolla longioribus. *Sp. pl. p.* 807. *Scop. carn. tom.* 1. *p.* 432.

Mentha caule proſtrato, foliis ſubrotundis, obiter dentatis, ſtaminibus exertis. *Hall. hiſt. n.* 221.

Pulegium. *Riv. mon. t.* 23.

Le Pouliot. Regn. bot.

In uliginoſis, paluſtribus. Fl. Iulio, Auguſto. Per.

GLECOMA. Calyx quinquefidus. Antherarum ſingulum par in formam crucis connivens.

hederacea. 1. G. foliis reniformibus crenatis. *Syſt. veget. p.* 534. *Fl. dan. t.* 789. *Curtis lond.*

Chamæclema caule procumbente, radicato, foliis reniformibus, rotunde crenatis. *Hall. hiſt. n.* 245.

Calamintha hederacea. *Scop. carn. tom.* 1. *p.* 423.

La Lierre terreſtre, ou la Terrette. Regn. bot.

In ſylvaticis, pomariis, ad ſepes, & alibi. Fl. primo vere. Per.

LAMIUM. Corollæ labium ſuperius integrum, fornicatum; labium inferius bilobum; faux utrinque margine dentata.

Orvala. 1. L. foliis cordatis inæqualiter arguteque ſerratis, corollis fauce inflata, calyce colorato. *Sp. pl. p.* 808. *Curtis magaz.* 172.

Lamium pannonicum. *Scop. carn. tom.* 1. *p.* 406. *t.* 27.

Orvala garganica. *Sp. pl. p.* 807.

Papia garganica, foliis urticæ alternis & eleganter inciſis, flore purpureo. *Mich. gen. p.* 20. *t.* 17.

Galeopſis maxima pannonica. *Clus. hiſt. p. XXXVI.*

In collibus, ad ſepes, in ceſpitoſis, ad aquas Carnioliæ, Croatiæ, Litoralis. Fl. primo vere. Per.

maculatum. 2. L. foliis cordatis acuminatis, verticillis decemfloris. *Sp. pl. p.* 809.

Lamium alba linea notatum. *Bauh. pin.* 231.

In ſylvis, ad ſepes Auſtriæ, Pannoniæ. Fl. æſtate. Per.

album. 3. L. foliis cordatis acuminatis ſerratis petiolatis, verticillis vigintifloris. *Sp. pl. p.* 809. *Fl. dan. t.* 594. *Curtis lond. Scop. carn. tom.* 1. *p.* 407.

Lamium foliis cordato - lanceolatis, ſerratis, verticillis multifloris. *Hall. hiſt. n.* 271.

Lamium album non fœtens, folio oblongo. *Bauh. pin.* 231.

L' Ortie blanche. Regn. bot.

Ad ſepes, pagos, in ſylvis. Fl. æſtate. Per.

purpureum. 4. L. foliis cordatis obtuſis petiolatis. *Sp. pl. p.* 809. *Fl. dan. t.* 523. *Curtis lond. Scop. carn. tom.* 1. *p.* 407.

Lamium foliis cordatis obtuſis, in ſummo ramo congeſtis. *Hall. hiſt. n.* 272.

Lamium purpureum fœtidum, folio ſubrotundo. *Bauh. pin.* 230.

In ruderatis, cultis, ad vias, ſepes. Fl. primo vere.

amplexicaule. 5. L. foliis floralibus ſeſſilibus amplexicaulibus obtuſis. *Sp. pl. p.* 809. *Fl. dan. t.* 752. *Curtis lond. Scop. carn. tom.* 1. *p.* 408.

Lamium foliis radicalibus petiolatis lobatis, ſuperioribus rotundis, amplexicaulibus, inciſis. *Hall. hiſt. n.* 273.

Lamium, folio caulem ambiente, minus. *Bauh. pin.* 231.

Habitat & Fl. cum præcedente.

GALEOPSIS. Corollæ labium ſuperius ſubcrenatum, fornicatum; labium inferius ſupra bidentatum.

Ladanum. 1. G. internodiis caulinis æqualibus, verticillis omnibus remotis, calycibus inermibus. *Syſt. veget. p.* 534 *Scop. carn. tom.* 1. *p.* 421.

Galeopſis foliis linearibus, ſerratis, verticillis diſſitis. *Hall. hiſt. n.* 266.

Ladanum ſegetum. *Riv. mon. t.* 24.

In ſterilibus aridis, in arvis, alibique. Fl. æſtate.

Variat foliis integerrimis, & plus vel minus dentatis.

Tetrahit. 2. G. internodiis caulinis ſuperne incraſſatis, verticillis ſummis ſubcontiguis, calycibus ſubpungentibus. *Syſt. veget. p.* 535. *Fl. dan. t.* 929. *Scop. carn. tom.* 1. *p.* 422.

Galeopſis caule hirto, foliis ovato-lanceolatis ſerratis. *Hall. hiſt. n.* 268.

Galeopſis caule hirto, foliis ovato-lanceolatis, ſerratis, flore calyce quadruplo majore. *Hall. hiſt. n.* 269.

In cultis, ſylvis, ad ſepes, in paluſtribus. Fl. æſtate.

GALEOBDOLON. Corollæ labium ſuperius erectum fornicatum; labium inferius trifidum: laciniis inæqualibus, media productiore. *Huds. angl. ed.* 2. 257.

Galeopſis. 1. G. verticillis ſexfloris, involucro tetraphyllo. *Huds. l. c. Curtis lond.*

Cardiaca foliis petiolatis, cordatis, verticillis folioſis. *Hall. hiſt. n.* 274.

Galeopſis Galeobdolon. *Sp. pl. p.* 810.

Leonurus Galeobdolon. *Scop. carn. tom.* 1. *p.* 409.

In umbroſis, ad ſepes. Fl. Aprili, Majo. Per.

BETONICA. Calyx ariſtatus. Corollæ labium ſuperius adſcendens, planiuſculum; tubus cylindricus.

officinalis. 1. B. ſpica interrupta, corollarum lacinia labii intermedia emarginata. *Sp. pl. p.* 810. *Fl. dan. t.* 726. *Curtis lond. Scop. carn. tom.* 1. *p.* 422.

Betonica foliis petiolatis, imis cordatis, ſuperioribus crenatis, ſpica brevi, inſidente foliis. *Hall. hiſt. n.* 264.

Betonica. *Clus. hiſt. p. XXXIX.*

La Betoine. Regn. bot.

In collibus herbidis, pratis, ſylvis. Fl. Iunio, Iulio. Per.

alopecuros. 2. B. ſpica baſi folioſa, corollis galea bifida. *Sp. pl. p.* 811. *Iacqu. auſtr. t.* 78. *Crantz faſc. p.* 256.

Sideritis alopecuros. *Scop. carn. tom.* I. *p.* 413. *t.* 28.

Betonica albo flore. *Clus. hiſt. p. XXXVIII.*

Horminum minus album, betonicæ facie. *Bauh. pin.* 239. *prodr.* 114.

In alpinis. Fl. Iulio. Per.

hirſuta. 3. B. ſpica baſi folioſa, corollis galea integra. *Syſt. veget. p.* 535. *Murr. in comm. gœtt.* 1779. *tom.* 2. *p.* 13. *t.* 3.

In herbidis montium Carniæ. *I. Hoſt.* Fl. Iulio. Per.

STACHYS. Corollæ labium ſuperius fornicatum; labium inferius lateribus reflexum: lacinia intermedia majore emarginata. Stamina deflorata verſus latera reflexa.

ſylvatica. 1. S. verticillis ſexfloris, foliis cordatis petiolatis. *Sp. pl. p.* 811. *Curtis lond. Scop. carn. tom.* I. *p.* 410.

Cardiaca foliis cordatis, ſerratis, verticillis nudis, ſpicatis. *Hall. hiſt. n.* 275.

Galeopſis legitima. *Clus. hiſt. p. XXXVI.*

L' Ortie morte. Regn. bot.

In umbroſis. Fl. æſtate. Per.

paluſtris. 2. S. verticillis ſubſexfloris, foliis lineari-lanceolatis ſemiamplexicaulibus ſeſſilibus. *Syſt. veget. p.* 535. *Curtis lond. Scop. carn. tom.* I. *p.* 411.

Stachys foliis ſubhirſutis, elliptico-lanceolatis, breviter petiolatis, verticillis ſpicatis. *Hall. hiſt. n.* 257.

Stachys paluſtris fœtida. *Bauh. pin.* 231.

In paluſtribus, ad foſſas, rivulos, ſepes humidas. Fl. æſtate. Per.

alpina. 3. S. verticillis multifloris, foliorum ſerraturis apice cartilagineis, corollis labio plano. *Sp. pl. p.* 812.

Stachys foliis hirſutis, cordatis, verticillis ſubtomentoſis. *Hall. hiſt. n.* 256.

Pſeudo-ſtachys alpina. *Bauh. pin.* 236. *prodr.* 113.

In Auſtriæ, Carnioliæ montanis ſylvaticis. Fl. æſtate. Per.

germanica. 4. S. verticillis multifloris, foliorum ſerraturis imbricatis, caule lanato. *Sp. pl. p.* 812. *Iacqu. auſtr. t.* 319. *Fl. dan. t.* 684. *Scop. carn. tom.* 1. *p.* 412.

Stachys tomentoſa, foliis imis cordatis, ſuperioribus elliptico-lanceolatis, verticillis denſiſſimis. *Hall. hiſt. n.* 253.

Stachys lanata. *Crantz fasc. p.* 267.

Stachys major germanica. *Bauh. pin.* 236.

In apricis ſ.cis. Fl. Iunio, Iulio. Per.

recta. 5. S. verticillis ſubſpicatis, foliis cordato-ellipticis crenatis ſcabris, caulibus adſcendentibus. *Syſt. veget. p.* 536. *Iacqu. auſtr. t.* 359.

Betonica foliis hirſutis, ovatis, rotunde crenatis, bracteis ovato-lanceolatis, ariſtatis, integerrimis. *Hall. hiſt. n.* 262.

Stachys Betonica. *Scop. carn. tom.* 1. *p.* 411. *Crantz fasc. p.* 264.

Sideritis prima vulgaris. *Clus. hiſt. p. XXXIX.*

In pratis ſiccioribus, collibus, montanisque apricis. Fl. æſtate. Per.

annua. 6. S. verticillis ſexfloris, foliis ovato-lanceolatis trinerviis lævibus petiolatis, caule erecto. *Syſt. veget. p.* 536. *Iacqu. auſtr. t.* 360.

Betonica foliis ovatis, crenatis, bracteis ovato-lanceolatis. *Hall. hiſt. n.* 263.

Sideritis ſecunda. *Cluſ. hiſt. p. XXXIX.*

In Auſtriæ, Pannoniæ arvis. Fl. a Iulio ad Septembrim.

BALLOTA. Calyx hypocrateriformis, quinquedentatus, decemſtriatus. Corollæ labium ſuperius crenatum, concavum.

nigra. 1. B. foliis cordatis indiviſis ſerratis, calycibus acuminatis. *Sp. pl. p.* 814. *Scop. carn. tom.* 1. *p.* 415.

Ballote foliis cordatis, ſerratis, verticillis nudis. *Hall. hiſt n.* 259.

Marrubium nigrum ſeu Ballote. *Bauh. hiſt.* 3. *p.* 318. *Blakw. herb. t.* 136.

La Ballote. Regn. bot.

Ad pagos, vias, ſepes, & alibi. Fl. æſtate. Per.

MARRUBIUM. Calyx hypocrateriformis, rigidus, decemſtriatus. Corollæ labium ſuperius bifidum, lineare, rectum.

peregrinum. 1. M. foliis ovato-lanceolatis ſerratis, calycum denticulis ſetaceis. *Sp. pl. p.* 815. *Iacqu. auſtr. t.* 160. *Crantz faſc. p.* 274.

Marrubium ſupinum. *Scop. carn. tom.* 1. *p.* 414.

Marrubium alterum pannonicum. *Cluſ. hiſt. p. XXXIV.*

In agro vindobonenſi ad vias, aggeres. In Litoroli prope Duinum. Fl. æſtate. Per.

vulgare. 2. M. dentibus calycinis ſetaceis uncinatis. *Sp. pl. p.* 816. *Scop. carn. tom.* 1. *p.* 414. *Cluſ. hiſt. p. XXXIV.*

Marrubium dentibus calycinis denis, recurvis. *Hall. hiſt. n.* 258.

Marrubium album vulgare. *Bauh. pin.* 230.

Le Marrube blanc. Regn. bot.

In ruderatis, ad vias. Fl. æſtate. Per.

LEONURUS. Antheræ punctis nitidis adſperſæ.

Cardiaca. 1. L. foliis caulinis lanceolatis, trilobis. *Sp. pl. p.* 817. *Fl. dan. t.* 727.

Cardiaca foliis tripartitis lanceolatis. *Hall. hiſt. n.* 276.

L' Agripaume, ou la Cardiaque. Regn. bot.

In ruderatis, ad ſepes, pagos Auſtriæ, Pannoniæ, Styriæ. Fl. æſtate. Per.

CHAITURUS. Calycis laciniæ erectæ. Corollæ labium ſuperius integrum fornicatum: inferius trifidum patens: tubus longitudine calycis. Stamina vix tubi longitudine.

leonuroides. 1. Chaiturus. *Ehrhart. han. magaz. ann.* 1781. 27. *p.* 421. *Wild. prodr. p.* 200.

Leonurus *Marrubiaſtrum*, foliis ovatis lanceolatisque ſerratis, calycibus ſeſſilibus ſpinoſis. *Sp. pl. p.* 117. *Iacqu. auſtr. t.* 405.

In Auſtria, Pannonia ad pagos, in ſylvis planitiei, præcipue ad fluvios Laitham, Danubium. Fl. a Iulio ad Septembrim.

PHLOMIS. Calyx angulatus. Corollæ labium ſuperius incumbens, compreſſum, villoſum.

tuberoſa. 1. P. involucris hiſpidis, ſubulatis, foliis cordatis ſcabris, caule herbaceo. *Sp. pl. p.* 819.

Galeopſis maxima, foliis hormini. *Bux. cent.* 1. *p.* 4. *t.* 6.

In ſylvaticis, & ad margines agrorum in Pannonia inter *Barendorf* & *Bruck an der Leitha.* Fl. Iunio, Iulio. Per.

CLINOPODIUM. Calyx bilabiatus. Involucrum multiſetum, verticillis ſubjectum.

vulgare. 1. C. capitulis ſubrotundis hispidis, bracteis ſetaceis. *Sp. pl. p.* 821. *Fl. dan. t.* 930. *Scop. carn. tom.* 1. *p.* 424.

Clinopodium foliis ovatis, lanceolatis, obiter dentatis, ſtipulis ſetaceis. *Hall. hiſt. n.* 239.

Clinopodium Matthioli. *Clus. hiſt. p.* 354.

In montoſis, cespitoſis, ad ſepes. Fl. æſtate. Per.

ORIGANUM. Calyx bilabiatus. Strobilus tetragonus ſpicatus, calyces colligens.

vulgare. 1. O. ſpicis ſubrotundis paniculatis conglomeratis, bracteis calyce longioribus ovatis. *Sp. pl. p.* 824. *Fl. dan. t.* 638. *Scop. carn. tom.* 1. *p.* 429.

Origanum foliis ovatis, umbellis coloratis, ſtaminibus exertis. *Hall. hiſt. n.* 233.

L' Origan commun. Bulliard. herb. t. 193.

In collibus, montoſis ſylvaticis. Fl. Iunio, Iulio. Per.

THYMUS. Calyx bilabiatus fauce villis clauſa.

Serpyllum. 1. T. floribus capitatis, caulibis repentibus, foliis planis obtuſis baſi ciliatis. *Sp. pl. p.* 825. *Curtis lond. Scop. carn. tom.* 1. *p.* 426.

Thymus foliis ovatis, baſi ciliatis. *Hall. hiſt. n.* 235.

Thymus foliis ellipticis hirſutis. *Hall. hiſt. n.* 236.

Serpyllum vulgare repens. *Clus. hiſt. p.* 359.

Serpyllum pannonicum primum, ſeu ſerpyllum auſtriacum citri odore. *Clus. hiſt. p.* 359.

Serpyllum anguſto glabroque folio. *Clus. hiſt. p.* 359.

Serpyllum anguſto lanuginoſoque folio, ſeu Serpyllum pannonicum tertium. *Clus. hiſt. p.* 360.

Le Serpolet. Regn. bot.

In campis, pratis, ſylvis, alibique. Fl. æſtate. Per.

vulgaris. 2. T. foliis revolutis ovatis, floribus verticillato-ſpicatis. *Sp. pl. p.* 825.

Thymum legitimum. *Clus. hiſt. p.* 357.

Le Thym. Regn. bot.

In aſperis Litoralis. Fl. Iunio. Per.

Acinos. 3. T. floribus verticillatis, pedunculis unifloris, caulibus erectis ſubramoſis, foliis acutis ſerratis. *Syſt. veget. p.* 542. *Curtis lond. Fl. dan. t.* 814. *Scop. carn. tom.* I. *p.* 426.

Clinopodium foliis ovatis dentatis, flore folio breviore. *Hall. hiſt. n.* 237.

Clinopodium vulgare. *Clus. hiſt. p.* 354.

Le Thym baſilic. Bulliard herb. t. 318.

In collibus, ad agrorum margines, ad vias, vineas. Fl. æſtate.

alpinus. 4. T. verticillis ſexfloris, foliis obtuſiuſculis concavis ſubſerratis. *Sp. pl. p.* 826. *Iacqu. auſtr. t.* 96. *Scop. carn. tom.* I. *p.* 425.

Clinopodium foliis ovatis acutis, ſerratis, flore folio majori, *Hall. hiſt. n.* 238.

Thymus montanus. *Crantz faſc. p.* 278.

Clinopodium auſtriacum. *Clus. hiſt. p.* 353.

In alpinis, & montanis. Fl. a Majo ad Septembrim. Per.

MELISSA. Calyx aridus, ſupra planiuſculus: labio ſuperiore ſubfaſtigiato. Corollæ labium ſuperius ſubfornicatum, bifidum; labio inferiore lobo medio cordato.

officinalis. 1. M. racemis axillaribus verticillatis: pedicellis ſimplicibus. *Sp. pl. p.* 827. *Scop. carn. tom.* 1. *p.* 429.

Meliſſa foliis cordatis, ſerratis, petiolis folio brevioribus. *Hall. hiſt. n.* 242.

La Meliſſe, ou Citronelle. Regn. bot.

In Auſtriæ ſylvaticis prope *Bertholdsdorf, Baaden.* In Litorali frequentiſſima in cepitoſis, ad ſepes, & alibi. Fl. Iulio, Auguſto. Per.

grandiflora. 2. M. pedunculis axillaribus dichotomis longitudine florum. *Sp. pl. p.* 827. *Wulfen in Iacqu. collect. tom.* 2. *p.* 134. *Curtis magaz.* 208.

Thymus grandiflorus. *Scop. carn. tom.* 1. *p.* 424.

Calamintha magno flore. *Bauh. pin.* 229. *Riv. mon.* 43.

In montanis, & ſubalpinis Croatiæ, Carnioliæ, Litoralis. Fl a Iunio ad Auguſtum.

Radix in meliori ſolo ſtolonibus reptat. Folia oppoſita, ſerrata. Pedunculi ramoſi. Bracteæ lineari-lanceolatæ, indiviſæ, ciliatæ. Calyx fauce barbatus.

Nepeta. 3. M. pedunculis axillaribus dichotomis folio longioribus, caule adſcendente hirſuto. *Syſt. veget. p.* 542.

Meliſſa foliis hirſutis, ovato-lanceolatis, ſerratis, petiolis florigeris ramoſis, folia ſuperantibus. *Hall. hiſt. n.* 240.

Calamintha pulegii odore ſeu Nepeta. *Bauh. pin.* 228. *Zann. iſt. p.* 46. *t.* 187.

Thymus Calamintha. *Scop. carn. tom.* 1. *p.* 225.

Calamintha officinalis. *Blakw. herb. t.* 167.

In Auſtria ſuperiore ad margines ſylvarum infrequens circa Lentiam. In Tyroli, Carinthia, Carniolia, Forojulio, Litorali frequentiſſima in montium declivibus, & ſaxoſis aſperis. Fl. æſtate. Per.

DRACOCEPHALUM. Calyx bilabiatus. Corollæ faux inflata: labium ſuperius concavum.

auſtriacum. 1. D. floribus ſpicatis, foliis mucronatis: caulinis pinnatifidis, rameis indiviſis. *Iacqu. collect. tom.* 1. *p.* 119. *ic. t.* 112. *Sp. pl. p.* 829.

Chamæpitys auſtriaca. *Cluſ. hiſt. p. CLXXXV.*

In Auſtria in montibus prope *Bertholdsdorf, Rodaun, Stahrenberg* retro *Fiſchau* inter *Wopfing* & *Oberpieſting. Cluſius.* Fl. Majo, Iunio. Per.

Folia ſeſſilia, juniora ciliata, mucronata, adultiora margine nuda, & apice inermia. Bracteæ trifidæ, mucronatæ. Calycis labium ſuperius integrum, marginibus reflexum; inferius trifidum.

HORMINUM. Calyx campanulatus: laciniis quatuor ſubæqualibus; quinta majore emarginata. Corollæ labium ſuperius concavum.

pyrenaicum. 1. H. foliis cordatis obtuſis, caule nudo. *Sp. pl. p.* 831. *Hall. hiſt. n.* 243.

Meliſſa pyrenaica. *Scop. ann.* 2. *p.* 55. *Iacqu. hort. tom.* 2. *p.* 86. *t.* 183.

Meliſſa pyrenaica, caule brevi, plantaginis folio. *Tourn. inſt.* 193.

In Carnioliæ alpibus wochinenſibus in monte *Sucha.* Fl. Iulio, Auguſto. Per.

MELITTIS. Calyx bilabiatus, tubo corollæ amplior. Corollæ labium ſuperius planum; labium inferius crenatum. Antheræ cruciatæ.

Meliſſophyllum. 1. Melittis. *Sp. pl. p.* 832. *Iacqu. auſtr. t.* 26. *Scop. carn. tom.* 1. *p.* 421.

Meliſſophyllum verum. *Fuchs. p.* 498. *Hall. hiſt. n.* 244.

Lamium pannonicum primum. *Clus. hiſt. p. XXXVI.*

La Meliſſe batarde. Regn. bot.

Ad margines ſylvarum, & in fruticoſis. Fl. Majo, Iunio. Per.

Folia petiolata, ovata, dentato-ſerrata, villoſiuscula & rugoſa. Corollæ ex purpureo alboque variæ, nunc ſaturatius coloratæ, nunc dilutius, ſæpe etiam totæ niveæ. Iacqu. l. c.

SCUTELLARIA. Calyx ore integro: poſt floreſcentiam clauſo, operculato.

galericulata. 1. S. foliis cordato-lanceolatis crenatis, floribus axillaribus. *Sp. pl. p.* 835. *Fl. dan. t.* 637. *Curtis lond.*

Caſſida foliis oblonge cordatis, crenatis, verticillis nudis, bifloris. *Hall. hiſt. n.* 280.

Caſſida galericulata. *Scop. carn. tom.* 1. *p.* 430.

La Toque tertianáire. Bulliard herb. t. 275.

Secus aquas, in ſylvis umbroſis, humidis. Fl. æſtate. Per.

haſtifolia. 2. S. foliis integerrimis: inferioribus haſtatis, ſuperioribus ſagittatis. *Sp. pl. p.* 835.

Caſſida haſtifolia. *Scop. carn. tom.* 1. *p.* 430.

Habitat & Fl. cum præcedente. Per.

PRUNELLA. Calyx bilabiatus. Filamenta bifurca, altero apice antherifera. Stigma bifidum.

vulgaris. 1. P. foliis omnibus ovato - oblongis ſerratis petiolatis. *Syſt. veget. p.* 547. *Curtis lond.*

Prunella foliis ovatis, oblongis, calycibus ſuperne truncatis. *Hall. hiſt. n.* 277.

Prunella grandiflora. *Iacqu. auſtr. t.* 317.

Prunella prima non vulgaris. *Clus. hiſt. p. XLII.*

La Brunelle. Regn. bot.

In pratis, collibus apricis, atque in montanis. Fl. æſtate. Per.

Variat flore minore, & ampliare; foliis integris, & laciniatis.

laciniata. 2. P. foliis ovato - oblongis petiolatis: ſupremis quatuor lanceolatis dentatis. *Sp. pl. p.* 837. *Iacqu. auſtr. t.* 376.

Prunella foliis imis ovatis oblongis, caulinis dentatis & ſemipinnatis. *Hall. hiſt. n.* 279.

Prunella ſecunda non vulgaris, albo flore. *Clus. hiſt. p. XLIII.*

In montanis, & in pratis ſiccioribus. Fl. Iunio, Iulio. Per.

ANGIOSPERMIA.

BARTSIA. Calyx bilobus: lobis emarginatis, apicibus coloratis. Corollæ labium ſuperius longius. Capſula bilocularis.

alpina. 1. B. foliis oppoſitis cordatis obtuſe ſerratis. *Sp. pl. p.* 839. *Fl. dan. t.* 43. *Scop. carn. tom.* 1. *p.* 434.

Stæhelina foliis cordatis, amplexicaulibus, ſerratis, floralibus coloratis. *Hall. hiſt. n.* 312.

In alpibus. Fl. Iulio, Auguſto. Per.

Folia obſcuri coloris, ſuprema, ut flores, violacea. Hall. l. c.

RHINANTHUS. Calyx quadrifidus, ventricoſus. Capſula bilocularis, obtuſa, compreſſa.

Criſta galli. 1. R. corollarum labio ſuperiore compreſſo breviore, calycibus glabris. *Pollich. palat. tom.* 2. *p.* 175. *Fl. dan. t.* 981.

Alectorolophus calycibus glabris. *Hall. hiſt. n.* 313.

Mimulus Criſta galli. *Scop. carn. tom.* 1. *p.* 434.

La Cocriſte glabre. Bulliard herb t. 125.

In pratis. Fl. Majo, Iunio.

Galea auriculis bifidis. Nectarium acuminatum. Calyces glabri. Scop. l. c.

Alectorolophus. 2. R. corollarum labio ſuperiore compreſſo breviore, calycibus villoſis. *Pollich. palat. tom.* 2. *p.* 177.

Alectorolophus calycibus hirſutis. *Hall. hiſt. n.* 314.

Mimulus Alectorolophus. *Scop. carn. tom.* 1. *p.* 435.

Habitat & Fl. cum præc.

Galea auriculis truncatis. Nectarium rotundatum. Calyces villoſi. Scop. l. c.

EUPHRASIA. Calyx quadrifidus, cylindricus. Capſula bilocularis, ovato-oblonga. Antheræ inferiores altero lobo baſi ſpinoſæ.

officinalis. 1. E. foliis ovatis lineatis argute dentatis. *Sp. pl. p.* 481. *Curtis lond. Scop. carn. tom.* 1. *p.* 435. *Hall. hiſt. n.* 303.

L' Euphraiſe officinale. Bulliard herb. t. 233.

In pratis, aliisque herbidis locis. Fl. Iulio, Auguſto, Septembri.

Variat foliis profunde & tenuiter diviſis, nec non exiguis nigricantibusque; floribus parvis, luteis, vel ex luteo, cœruleo & albo variegatis.

Odontites. 2. E. foliis linearibus: omnibus ſerratis. *Sp. pl. p.* 841. *Fl. dan. t.* 655. *Curtis lond. Scop. carn. tom.* 1. *p.* 435.

Odontites bracteis ſerratis hirſutis. *Hall. hiſt. n.* 304.

Ad foſſas, rivulos, in pratis. Fl. Iulio, Auguſto.

lutea. 3. E. foliis linearibus ſerratis: ſuperioribus integerrimis. *Sp. pl. p.* 842. *Iacqu. auſtr. t.* 398. *Scop. carn. tom.* 1. *p.* 436.

Odontites bracteis glabris integerrimis. *Hall. hiſt. n.* 305.

In ſiccis, glareoſis, Auſtriæ, Forijulii. Fl. Auguſto, Septembri.

MELAMPYRUM. Calyx quadrifidus. Corollæ labium ſuperius compreſſum, margine replicato. Capſula bilocularis, obliqua, hinc dehiſcens. Semina duo, gibba.

criſtatum. 1. M. ſpicis quadrangularibus, bracteis cordatis, compactis denticulatis imbricatis. *Sp. pl. p.* 842. *Scop. carn. tom.* 1. *p.* 437.

Melampyrum foliis integerrimis, floribus ſpicatis, bracteis duplicatis, criſtatis. *Hall. hiſt. n.* 311.

In collibus, pratis ſterilibus, montoſis ſylvaticis. Fl. a Iunio ad Septembrim.

arvenſe. 2. M. ſpicis conicis laxis: bracteis dentato-ſetaceis; coloratis. *Sp. pl. p.* 842. *Fl. dan. t.* 911.

Melampyrum foliis integerrimis, bracteis ſemipinnatis. *Hall. hiſt. n.* 310.

Parietaria ſylveſtris tertia. *Cluſ. hiſt. p. XLV.*

Melampyrum purpuraſcente coma. *Bauh. pin.* 234.

Inter ſegetes. Fl. cum præc.

nemoroſum. 3. M. floribus ſecundis lateralibus, bracteis dentatis: ſummis coloratis ſterilibus, calycibus lanatis. *Sp. pl. p.* 843. *Fl. dan. t.* 305. *Scop. carn. tom.* I. *p.* 437.

Melampyrum foliis ovato-lanceolatis, floralibus hamatis, acute dentatis, patulis, calycibus hirſutis. *Hall. hiſt. n.* 309.

Parietaria ſylveſtris prima. *Cluſ. hiſt. p. XLIV.*

Melampyrum coma cœrulea. *Bauh. pin.* 234.

In nemoribus, præcipue ad ſylvarum margines. Fl. Iulio, Auguſto, Septembri.

pratenſe. 4. M. floribus ſecundis lateralibus: conjugationibus remotis, corollis clauſis. *Sp. pl. p.* 843. *Scop. carn. tom.* I. *p.* 437.

Melampyrum foliis imis integerrimis, mediis dentatis, floralibus haſtatis. *Hall. hiſt. n.* 308.

Parietaria ſylveſtris ſecunda. *Cluſ. hiſt. p. XLIV.*

Melampyrum luteum latifolium. *Bauh. pin.* 234.

In ſylvaticis. Fl. æſtate.

ſylvaticum. 5. M. floribus ſecundis lateralibus: conjugationibus remotis, corollis hiantibus. *Sp. pl. p.* 834. *Fl. dan. t.* 145. *Scop. carn. tom.* I. *p.* 438.

Melampyrum foliis integerrimis, floribus hiantibus. *Hall. hiſt. n.* 307.

In ſylvaticis, præſertim ſubalpinis. Fl. æſtate.

LATHRÆA. Calyx quadrifidus. Glandula depreſſa, ad baſin ſuturæ germinis. Capſula unilocularis.

Squamaria. 1. L. caule ſimpliciſſimo, corollis pendulis: labio inferiore trifido. *Sp. pl. p.* 844. *Fl. dan. t.* 136.

Squamaria caule ſimpliciſſimo, floribus nutantibus, labello trifido. *Hall. hiſt. n.* 297.

Squamaria Orobanche. *Scop. carn. tom.* 1. *p.* 438.

Dentaria major, ſive aphyllos. *Clus. hiſt. p.* CXX.

In ſylvis umbroſis ad arborum radices. Fl. primo vere. Per.

TOZZIA. Calyx quinquedentatus. Capſula unilocularis, globoſa, monoſperma.

alpina. 1. Tozzia. *Sp. pl. p.* 844. *Hall. hiſt. n.* 298. *Iacqu. auſtr. t.* 165.

Tozzia alpina lutea, alſines folio, radice ſquamata. *Mich. gen. p.* 20. *t.* 16.

In declivibus alpium ſaxoſis udis. Fl. Iulio, Auguſto. Per.

Radix ſubrotunda, ſquamis tecta. Folia ovata, ſerrata, glabra, oppoſita, ſeſſilia. Pedunculi axillares, ſolitarii, uniflori. Calyx monophyllus, tubuloſus, bilabiatus: labio inferiore tridentato; ſuperiore bidentato. Corolla lutea, *bilabiata: labio inferiore trifido; ſuperiore bifido. Filamenta ſingula gerunt antheras geminas, baſi ſetam emittentes.* Iacqu. l. c.

PEDICULARIS. Calyx quinquefidus. Corollæ labium ſuperius compreſſum. Capſula bilocularis, mucronata, obliqua. Semina tunicata.

paluſtris. 1. P. caule ramoſo, calycibus criſtatis calloſo-punctatis, corollis labio obliquis. *Sp. pl. p.* 845. *Fl. dan. t.* 225. *Scop. carn. tom.* 1. *p.* 439.

Pedicularis caule ramoſo, foliis pinnatis, pinnis pinnatis, dentatis, calyce bifido. *Hall. hiſt. n.* 320.

La Pediculaire des marais. Bulliard herb. t. 129.

In pratis udis. Fl Iunio, Iulio, Auguſto.

roſtrata. 2. P. caule declinato ſubramoſo, corollis galea roſtrato - acuminatis, calycibus criſtatis ſubhirſutis. *Kram. auſtr.* 183. *n.* 3. *Sp. pl. p.* 845. *Iacqu. auſtr. t.* 205. *Scop. carn. tom.* 1. *p.* 440.

Pedicularis caule procumbente, ramoſo, floribus roſtratis, ſparſis. *Hall. hiſt. n.* 322. *t.* 6. *f.* 1.

Alectorolophus alpina tertia, *Clus. hiſt. p.* CCX.

In alpibus. Fl. Iulio, Auguſto.

verticillata. 3. P. caule ſimplici, foliis quaternis. *Kram. auſtr.* 183. *n.* 4. *Sp. pl. p.* 846. *Iacqu. auſtr. t.* 206. *Scop. carn. tom.* 1. *p.* 441.

Pedicularis foliis pinnatis quaternis, floribus ſpicatis. *Hall. hiſt. n.* 318. *t.* 9. *f.* 2.

In alpibus. Fl. Iulio, Auguſto. Per.

recutita. 4. P. caule ſimplici, foliis pinnatifidis ſerratis, ſpica folioſa, calycibus coloratis, corollis obtuſis. *Syſt. veget. p.* 551. *Iacqu. auſtr. t.* 258.

Pedicularis caule ſimplici, foliis pinnatis, floribus ſpicatis, ſtaminibus eminentibus. *Hall. hiſt. n.* 316. *t.* 8. *f.* 2.

Pedicularis obſoleta. *Crantz faſc. p.* 316.

In alpium herbidis humidiuſculis. Fl. Iulio, Auguſto, Septembri. Per.

flammea. 5. P. caule ſimplici, foliis pinnatis retro imbricatis. *Sp. pl. p.* 846. *Hænke in Iacqu. collect. tom.* 2. *p.* 70.

Pedicularis caule ſimplici, foliis ſemipinnatis obtuſis: laciniis imbricatis crenatis. *Fl. lapp. n.* 244. *t.* 4. *f.* 2.

Pedicularis caule ſimplici, foliis pinnatis, pinnis dentatis, denticulis retroverſis. *Hall. hiſt. n.* 315. *t.* 8. *f.* 3.

In ſummis jugis alpium judenburgenſium in Styria ſuperiore. Fl. Iunio, Iulio. Per.

Floræ danicæ figura trigeſima nec forma nec coloribus refert P. flammeam e Lapponia. Retzii ſcand. p. 145.

roſea. 6. P. caule ſimplici, foliis pinnatis: pinnis oblongis pinnatifidis, calycibus hirſutis. *Syſt. veget. p.* 552. *Wulfen in Iacqu. miscell. tom.* 2. *p.* 57. *ic. t.* 115.

In Carinthia in ſummitate alpis villacenſis, & in ſupremis jugis alpis kibeggenſis. Fl. Iulio.

incarnata. 7. P. caule ſimplici, foliis pinnatis ſerratis, calycibus rotundatis glabris, corollis galea uncinatis acutis. *Sp. pl. p.* 847. *Iacqu. auſtr. t.* 140.

Pedicularis roſtrato-ſpicata. *Crantz faſc. p.* 317.

In Auſtria, Styria in alpium declivibus. Fl. Iulio, Auguſto. Per.

folioſa. 8. P. caule ſimplici, ſpica folioſa, corollis galea obtuſiſſima integra, calycibus quinquedentatis. *Syſt. veget. p.* 552. *Iacqu. auſtr. t.* 139.

Pedicularis ſpica folioſa. *Hall. hiſt. n.* 317. *t.* 9. *f.* 1.

Pedicularis comoſa. *Scop. carn. tom.* 1. *p.* 440.

In herbidis alpinis, & ſubalpinis. Fl. Iulio, Auguſto. Per.

tuberoſa. 9. P. caule ſimplici, calycibus crenatis, corollis galea roſtrato-aduncis. *Sp. pl. p.* 847. *Scop. carn. tom.* 1. *p.* 440.

Pedicularis caule ſimplici, floribus ſpicatis, roſtratis, calycibus glabris. *Hall. hiſt. n.* 323. *t.* 10.

In Carniolia in alpium wochinenſium, & tolmienſium herbidis. In Carnia in monte *Croſtis.* Fl. Iulio, Auguſto. Per.

Radix tantum in aliquibus individuis ad caulis exordium incraſſata. In nullo tuberoſa.

acaulis. 10. P. pedunculis radicalibus. *Scop. carn. tom.* 1. *p.* 439. *t.* 31. *Wulfen in Iacqu. collect. tom.* 1. *p.* 207. *t.* 14.

Pedicularis alpina acaulis flore magno. *Segu. ver. ſuppl. p.* 127.

In Carniolia in alpibus tolmienſibus circa *Ottaleſck*, & in paluſtribus labacenſibus. Fl. Majo.

ANTIRRHINUM. Calyx pentaphyllus. Corollæ baſis deorſum prominens nectarifera. Capſula bilocularis.

Cymbalaria. 1. A. foliis cordatis quinquelobis alternis, caulibus procumbentibus. *Sp. pl. p.* 851. *Curtis lond. Scop. carn. tom.* 1. *p.* 443.

Antirrhinum caule repente, foliis reniformibus, quinquelobatis. *Hall. hiſt. n.* 339.

Cymbalaria floſculis purpuraſcentibus. *Bauh. hiſt.* 3. *p.* 685. *Zann. iſt. p.* 77. *t.* 252.

La Cymbalaire. Regn. bot.

In Forojulio, Litorali in muris vetuſtis, maceriis, rupibus. Fl. æſtate. Per.

Elatine. 2. A. foliis haſtatis alternis, caulibus procumbentibus. *Sp. pl. p.* 851. *Curtis lond. Fl. dan. t.* 426. *Scop. carn. tom.* 1. *p.* 444.

Antirrhinum caule procumbente, foliis haſtatis, imis conjugatis, ſuperioribus alternis. *Hall. hiſt. n.* 340.

In arvis, locis arenoſis Auſtriæ, Pannoniæ, Forijulii, Iſtriæ. Fl. Iulio, Auguſto.

Folia petiolata, ima ſubrotunda, oppoſita; proxima dentata, alterna, quæ ſequuntur haſtata. Curtis l. c.

ſpurium. 3. A. foliis ovatis alternis caulibus procumbentibus. *Sp. pl. p.* 851. *Scop. carn. tom.* 1. *p.* 443. *Curtis lond.*

Antirrhinum caule procumbente, foliis villoſis ovatis, imis conjugatis, ſuperioribus alternis. *Hall. hiſt. n.* 341.

La Velvote. Regn. bot.

In arvis. Fl. æſtate.

triphyllum. 4. A. foliis ternis ovatis. *Sp. pl. p.* 852.

Linaria ſicula latifolia triphylla. *Bocc. ſic.* 44. *t.* 22.

Linaria hiſpanica prima. *Clus. hiſt. p.* 320.

Linaria triphylla minor lutea. *Bauh. pin.* 212.

In montanis Iſtriæ, & inſularum venetarum. *I. Hoſt.* Fl. Iunio.

Folia inferiora terna, ſuprema alternatim ſparſa.

alpinum. 5. A. foliis quaternis lineari-lanceolatis, caule diffuſo, floribus racemoſis calcari recto. *Sp. pl. p.* 856. *Iacqu. auſtr. t.* 58. *Scop. carn. tom.* 1. *p.* 442.

Antirrhinum caule procumbente, rariter ſpicato, foliis verticillatis. *Hall. hiſt. n.* 338.

Linaria tertia ſtyriaca. *Clus. hiſt. p.* 322.

In alpium ſaxoſis. Fl. æſtate. Per.

minus. 6. A. foliis plerisque alternis lanceolatis obtuſis, caule ramoſiſſimo diffuſo. *Sp. pl. p.* 852. *Fl. dan. t.* 502. *Curtis lond.*

Antirrhinum viscidum, foliis inferioribus conjugatis, ellipticis, obtuſis, calcare dimidii floris longitudine. *Hall. hiſt. n.* 335.

In cultis, ruderatis, ad vias, & alibi. Fl. æſtate.

geniſti-folium. 7. A. foliis lanceolatis acuminatis, panicula virgata flexuoſa. *Sp. pl. p.* 858. *Iacqu. auſtr. t.* 244.

Linaria pannonica. *Clus. hiſt. p.* 321.

In Auſtriæ, Pannoniæ collibus apricis, & montium declivibus. Fl. æſtate. Per.

Linaria. 8. A. foliis lanceolato-linearibus confertis, caule erecto, ſpicis terminalibus ſeſſilibus, floribus imbricatis. *Sp. pl. p.* 858. *Curtis lond. Fl. dan. t.* 982. *Scop. carn. tom.* 1. *p.* 442.

Antirrhinum foliis linearibus, adſcendentibus, congeſtis, caule erecto, ſpicato. *Hall. hiſt. n.* 336.

La grande Linaire. Regn. bot.

In ruderatis, ad vias, & alibi. Fl. æſtate. Per.

majus. 9. A. corollis ecaudatis, floribus ſpicatis, calycibus rotundatis. *Sp. pl. p.* 859. *Scop. carn. tom.* 1. *p.* 444.

Antirrhinum foliis ellipticis obtuſis, floribus ſpicatis, calycibus obtuſis, calcare breviſſimo. *Hall. hiſt. n.* 333.

Le Muſle de Veaus, ou Muſeaude. Regn. bot.

Occurrit in muris vetuſtis. Copioſiſſime in Croatia in mœniis Zagrabiæ. Fl. æſtate. Per.

Orontium. 10. A. corollis ecaudatis, floribus ſubſpicatis, calycibus corolla longioribus. *Sp. pl. p.* 860. *Curtis lond.*

Antirrhinum foliis ellipticis obtuſis, floribus ſparſis, calycibus ſubulatis, longiſſimis, calcare breviſſimo. *Hall. hiſt. n.* 334.

In agris, arvis, & paſcuis ſiccioribus, alibique. Fl. æſtate.

SCROPHULARIA. Calyx quinquefidus. Corolla ſubgloboſa, reſupinata. Capſula bilocularis.

nodoſa. 1. S. foliis cordatis trinerviis, caule acutangulo. *Sp. pl. p.* 863. *Scop. carn. tom.* 1. *p.* 445.

Scrophularia caule quadrangulo, paniculato, foliis cordatis oblongis. *Hall. hiſt. n.* 326.

La grande Scrophulaire. Regn. bot.

In humidis, ad ſepes. Fl. æſtate. Per.

aquatica. 2. S. foliis cordatis obtuſis petiolatis decurrentibus, caule membranis angulato, racemis terminalibus. *Sp. pl. p.* 864. *Fl. dan. t.* 507. *Curtis lond. Scop. carn. tom.* 1. *p.* 445.

Scrophularia caule quadrangulo, alato, paniculato, foliis ovato-lanceolatis. *Hall. hiſt. n.* 325.

La Scrophulaire, ou l'Herbe du ſiege. Regn. bot.

In rivulis, ad foſſas, fluvios. Fl. æſtate. Per.

Huic radix fibroſa, præcedenti nodoſa.

auriculata. 3. S. foliis cordatis ſubtus tomentoſis baſi appendiculatis, racemis terminalibus. *Syſt. veget. p.* 560. *Scop. carn. tom.* 1. *p.* 446. *t.* 32.

Scrophularia aquatica montana, mollior. *Barr. rar.* 189. *t.* 274.

In Carniolia in declivibus alpium wochinenſium. Fl. Iulio, Auguſto. Per.

Tota planta, præſertim folia ſubtus, villoſa. Caules acute tetragoni, in plerisque individuis parum flexuoſi. Folia ſunt oppoſita, petiolata, cordata, oblonga, dentato-ſerrata: dentibus paucis uno alterove denticulo notatis. Floralia autem alterna lanceolata, acute ſerrata. Pedunculi axillares, ramoſi. Pedicellis ſingulis bractea linearis adhæret. Caulis, ubi flores, & pedunculi obſiti ſunt villis apice nudis, & hisce glanduliferis immixtis. In planta bipedali duo ſola foliorum paria auriculata obſervavi. In aliis auriculas vel nullas, vel harum loco ramulorum rudimenta vidi. Caulis adultus pubem dejicit, folia autem conſervant.

canina. 4. S. foliis pinnatis, racemo terminali nudo, pedunculis bifidis. *Sp. pl. p.* 865. *Scop. carn. tom.* 1. *p.* 446.

Scrophularia foliis pinnatis, pinnis lobatis, circumſerratis, caule paniculato. *Hall. hiſt. n.* 328.

Scrophularia Ruta canina. *Cluſ. hiſt. p.* CCIX.

Scrophularia Ruta canina dicta, vulgaris. *Bauh. pin.* 236. *Zann. iſt. p.* 242. *t.* 221.

In Tyroli, Carinthia, Carniolia, Croatia, Litorali, Forojulio in montium declivibus ſaxoſis, rupeſtribus, in muris. Fl. æſtate. Per.

DIGITALIS. Calyx quinquepartitus. Corolla campanulata, quinquefida, ventricoſa. Capſula ovata, bilocularis.

lutea. 1. D. calycinis foliolis lanceolatis, corollis acutis: labio ſuperiore bifido. *Sp. pl. p.* 867. *Iacqu. hort. tom.* 2. *p.* 47. *t.* 105.

Digitalis foliis calycinis lanceolatis, galea bifida, floribus immaculatis. *Hall. hiſt. n.* 332.

La Digitale jaune. Bulliard herb. t. 65.

In Croatia in ſylvis circa *Merkopail*, & *Raunagora.* Fl. Iulio. Per.

ambigua. 2. D. calycinis foliolis lanceolatis, corollis galea emarginata, foliis ſubtus pubescentibus. *Syſt. veget. p.* 562.

Digitalis foliis calycinis lanceolatis, galea inciſa, faucibus maculoſis. *Hall. hiſt n.* 331.

Digitalis ochroleuca. *Iacqu. auſtr. t.* 57.

In montanis, & ſubalpinis. Fl. a Iunio ad Auguſtum. Per.

ferruginea. 3. D. calycinis foliolis ovatis obtuſis patentibus, corollæ labio inferiore barbato. *Syſt. veget. p.* 562.

In Litorali in ſaxoſis, fruticoſis agri flumineuſis, circa Voloscam, alibique. Fl. Majo, Iunio. Per.

LINDERNIA. Calyx quinquepartitus. Corolla ringens labio ſuperiore breviſſimo. Stamina duo inferiora dente terminali antheraque ſublaterali. Capſula unilocularis.

pyxidaria. 1. L. foliis ovalibus integerrimis ſeſſilibus, pedunculis ſolitariis. *Syſt. veget. p.* 567. *Wulfen in Iacqu. collect. tom.* 1. *p.* 271.

Pyxidaria repens annua, floſculis monopetalis unilabiatis. *Lindern. hort. p.* 269.

Capraria gratioloides. *Sp. pl. p.* 876.

In Carinthia prope Clagenfurtum in ſylvarum ſupra *Keslix* ſtagnis. *Wulfen.* In Carnia in aquis ſtagnantibus non procul a monte *Croſtis. I. Hoſt.* Fl. Iulio, Auguſto.

LIMOSELLA. Calyx quinquefidus. Corolla quinquefida, æqualis. Stamina per paria approximata. Capſula unilocularis, bivalvis, polyſperma.

aquatica. 1. L. foliis lanceolatis. *Syſt. veget. p.* 572. *Fl. dan. t.* 69. *Hall. hiſt. n.* 300.

Plantaginella paluſtris. *Bauh. pin.* 190.

In Auſtriæ, Pannoniæ uliginoſis, paluſtribus. Fl. æſtate.

OROBANCHE. Calyx bifidus. Corolla ringens. Capſula unilocularis, bivalvis, polyſperma. Glandula ſub baſi germinis.

major. 1. O. caule ſimpliciſſimo pubescente, ſtaminibus ſubexertis. *Sp. pl. p.* 882. *Curtis lond. Scop. carn. tom.* 1. *p.* 448.

Orobanche caule ſimplici, ſtipula unica, calyce quadrifido. *Hall. hiſt. n.* 295.

Orobanche prima. *Clus. hiſt. p.* 270.

Orobanche major caryophyllum olens. *Bauh. pin.* 87.

In pratis, collibus, montibus. Fl. æſtate.

purpurea. 2. O. caule ſimplici ramoſoque, corollis quadrifidis, ſtaminibus calcaratis. *Syſt. veget. p.* 573. *Iacqu. auſtr. t.* 276.

In Auſtriæ collibus. Fl. æſtate.

ramoſa. 3. O. caule ramoſo, corollis quinquefidis. *Sp. pl. p.* 882.

Orobanche caule ramoſo, flore quinquepartito. *Hall. hiſt. n.* 296.

Orobanche tertia polyclonos. *Clus. hiſt. p.* 271.

Tergeſti. *I. Hoſt.* In Auſtriæ arvis reſtibilibus circa *Bruck an der Leitha. Indefeſſus Lübeck.* Fl. æſtate.

VITEX. Calyx quinquedentatus. Corollæ limbus ſexfidus. Bacca tetraſperma.

Agnus. 1. V. foliis digitatis ſerratis, ſpicis verticillatis. *Sp. pl. p.* 890. *Scop. carn. tom.* 1. *p.* 448.

L' Agnus caſtus. Regn. bot.

In Litorali ad mare. Fl. Iulio, Auguſto, Septembri. Fr.

ACANTHUS. Calyx foliolis trium parium alternantium, inæqualibus. Corollæ tubus barba clauſus. Labium ſuperius nullum; labium inferius maximum, trilobum. Capſula bilocularis.

mollis. 1. A. foliis ſinuatis inermibus. *Sp. pl. p.* 891.

L' Acanthe, ou la Brancurſine. Regn. bot.

In Litorali inter *Veprinaz* & Lauranam locis ceſpitoſis. In ſimilibus locis in inſulis venetis. Fl. Majo, Iunio. Per.

TETRADYNAMIA.

SILICULOSA.

MYAGRUM. Silicula ſtylo conico terminata: loculo ſubmonoſpermo.

perenne. 1. M. ſiliculis biarticulatis monoſpermis, foliis extrorſum ſinuatis denticulatis. *Sp. pl. p.* 893. *Iacqu. auſtr. t.* 414. *Scop. carn. tom.* 2. *p.* 10.

Myagrum ſiliquæ articulo primo ſtricto, altero globoſo ſtriato. *Hall. hiſt. n.* 525.

Rapiſtrum monoſpermum. *Bauh. pin.* 95. *prodr.* 37.

In Auſtriæ, Pannoniæ, Styriæ arvis, & ad vias. Fl. Iunio, Iulio.

Silicula biarticulata: articulo ſuperiore ovato, ſtylo roſtrato; inferiore oblongo, angulato; vel utroque ſemen gerente, vel unico ſeminifero, altero ſterili. Iacqu. l. c.

rugoſum. 2. M. ſiliculis ſulcatis piloſis rugoſis, foliis oblongis obtuſis dentatis. *Sp. pl. p.* 893. *Scop. carn. tom.* 2. *p.* 11.

Tergeſti ad mare. Fl. Iunio, Iulio.

auſtriacum. 3. M. ſiliculis ſubrotundis glabris, foliis oblongis amplexicaulibus, radice repente. *Syſt. veget. p.* 584. *Iacqu. auſtr. t.* 111.

In Auſtriæ, Pannoniæ uliginoſis, foſſis, & alibi. Fl. Iunio. Per.

Caules ſubanguloſi, fiſtuloſi. Folia oblonga, obtuſa, vel acutiuſcula, radicalia ad baſin attenuata & inciſa, caulina ſeſſilia, amplexicaulia. In planta in aquis creſcente ſunt folia radicalia & caulina ima pinnatifida, ſuperiora inciſa. Silicula globoſa, ſtylo obtuſo craſſoque coronata, obſolete pedicellata, bilocularis, ſemina utrinque continens circiter quina. Iacqu. l. c.

ſativum. 4. M. ſiliculis obovatis, pedunculis polyſpermis. *Sp. pl. p.* 894.

Alyſſum foliis ſagittatis, ſiliquis cordiformibus. *Hall. hiſt. n.* 489.

Alyſſum ſativum. *Scop. carn. tom.* 2. *p.* 9.

Inter ſegetes, ad agrorum margines. Fl. Majo, Iunio.

paniculatum. 5. M. ſiliculis lentiformibus orbiculatis punctato-rugoſis. *Sp. pl. p.* 894. *Fl. dan. t.* 204. *Scop. carn. tom.* 2. *p.* 11.

Rapiſtrum foliis amplexicaulibus auriculatis, acutis. *Hall. hiſt. n.* 522.

Myagro ſimilis ſiliqua rotunda. *Bauh. pin.* 109. *prodr.* 52.

In arvis. Fl. Innio, Iulio.

ſaxatile. 6. M. ſiliculis lentiformibus obovatis glabris, foliis oblongis ſerratis ſcabris, caule paniculato. *Sp. pl. p.* 894. *Iacqu. auſtr. t.* 128. *en. p.* 257.

Alyſſum foliis ſcabris, radicalibus petiolatis, ovatis, dentatis, caule ſubnudo flexuoſo. *Hall. hiſt. n.* 490.

Alyſſum alpinum. *Scop. carn. tom.* 2. *p.* 9.

Thlaſpi petræum myagroides. *Pon. in Cluſ. hiſt. p.* CCCXXXVII.

In rupeſtribus alpinis, & ſubalpinis. Fl. Majo, Iunio. Per.

BUNIAS. Silicula decidua, tetraëdra, angulis uno alterove acumine, non dehiſcens. Semina ſub ſingulo acumine ſingula.

Erucago. 1. B. ſiliculis tetragonis: angulis bicriſtatis. *Sp. pl. p.* 935. *Iacqu. auſtr. t.* 340.

Bunias ſiliqua tetragonia, quatuor alis emarginatis. *Hall. hiſt. n.* 526.

Eruca monſpeliaca, ſiliqua quadrangula echinata. *Bauh. pin.* 99. *prodr.* 41.

In Auſtriæ, Pannoniæ arvis. Fl. Iunio, Iulio.

Cakile. 2. B. ſiliculis ovatis lævibus ancipitibus. *Sp. pl. p.* 936.

Cakile maritima. *Scop. carn. tom.* 2. *p.* 35.

Eruca maritima, anguſtifolia italica, ſiliqua haſtæ cuspidi ſimili. *Bauh. pin.* 99. *Zann. iſt. p.* 45. *t.* 13. 177. 189. 193.

In Litorali ad mare. Fl. æſtate.

ISATIS. Silicula lanceolata, unilocularis, monoſperma, decidua, bivalvis: valvulis navicularibus.

tinctoria. 1. I. foliis radicalibus crenatis; caulinis ſagittati, ſiliculis oblongis. *Sp. pl. p.* 936.

Iſatis foliis radicalibus petiolatis, ovatis, caulinis amplexicaulibus. *Hall. hiſt. n.* 523.

Iſatis vel Glaſtum ſpontanea. *Bauh. hiſt.* 2. *p.* 909.

Le Päſtel, ou la Guede. Regn. bot.

In Auſtriæ, Pannoniæ aſperis ſylvaticis; arvis. Fl. Majo, Iunio.

CRAMBE. Filamenta quatuor, longiora apice bifurca: altero antherifero. Bacca ſicca, globoſa, decidua.

tataria. 1. C. foliis decompoſito-multifidis. *Syſt. veget. p.* 604. *Iacqu. miscell. tom.* 2. *p.* 274. *ic. t.* 129.

Tataria ungarica. *Clus. hiſt. p.* CXCI.

Tataria ungarica edulis, Panacis Heraclei folio, ſemine Libanotidis cachryoferæ. *Bauh. hiſt.* 3. *p.* 163.

In Pannoniæ comitatu heveſſienſi circa *Agriam* (*Erlau*) in agris. In Moravia in vineis ad pagum *Hurtau* circa *Auſſitz.* Fl. Aprili, Majo. Per.

ANASTATICA. Silicula retuſa, margine coronata valvulis diſſepimento duplo longioribus. Stylo intermedio mucronato, obliquo: loculis monoſpermis.

ſyriaca. 1. A. foliis acutis, ſpicis folio longioribus, ſiliculis ovatis roſtratis. *Sp. pl. p.* 895. *Iacqu. auſtr. t.* 6.

Myagrum roſtratum. *Scop. carn. tom.* 2. *p.* 11. *t.* 35.

In Auſtria, Pannonia ad vias, margines agrorum. Fl. Majo.

DRABA. Silicula integra, ovali - oblonga: valvis planiusculis, diſſepimento parallelis. Stylus breviſſimus.

aizoides. 1. D. ſcapo nudo ſimplici, foliis enſiformibus carinatis ciliatis. *Syſt. veget. p.* 585. *Iacqu. auſtr. t.* 192.

Draba cauliculis nudis, foliis linearibus ciliatis. *Hall. hiſt. n.* 498.

Draba alpina. *Scop. carn. tom.* 2. *p.* 5. *Iacqu. en. p.* 254.

Sedum minus duodecimum, alpinum ſextum. *Clus. hiſt. p. LXII.*

In alpium ſaxoſis. Fl. Iulio, Auguſto. Per.

ciliaris. 2. D. caule ſubnudo, foliis linearibus margine carinaque ciliatis, petalis integris. *Syſt. veget. p.* 585. *Scop. carn. tom.* 2. *p.* 6. *t.* 33.

In rupeſtribus in Carniolia in monte *Nanas*. *Scop.* Ego in Croatia in monte *Risnick*. Fl. Iunio. Per.

Folia lanceolata, margine ciliata, facie glabra, carina duobus vel tribus pilis munita: inferiora in roſulam congeſta, baſi attenuata; ſuperiora duo vel tria, alterna, baſi dilatata caulem amplexantia. Silicula oblonga.

verna. 3. D. ſcapis nudis, foliis ſubſerratis. *Syſt. veget. p.* 585. *Fl. dan. t.* 983. *Curtis lond. Scop. carn. tom.* 2. *p.* 8.

Draba cauliculis nudis, foliis ſubhirſutis, tridentatis. *Hall. hiſt. n.* 496.

In cultis, & incultis. Fl. primo vere.

Folia ſunt vel integerrima, vel plus minusve denticulata. Pili modo indiviſi, modo bi- vel trifurcati.

pyrenaica. 4. D. ſcapo nudo, foliis cuneiformibus palmatis trilobis. *Syſt. veget. p.* 585. *Iacqu. auſtr. t.* 228. *en. p.* 255. *Crantz faſc. p.* 13. *t.* 1.

In ſummis alpium jugis. Fl. Iunio, Iulio. Per.

Folia parva, craſſa, nitida, rigidula, plana, baſi in petiolum ſemiamplexantem attenuata, profunde trifida, ad apices & oras ciliata. Iacqu. l. c.

ſtellata. 5. D. foliis hirtis, integerrimis dentatisque, caule ſubunifolio, ſiliculis oblongis rectis. *Iacqu. en. p.* 256. *t.* 4. *f.* 3.

Draba hirta. *Iacqu. auſtr. t.* 432.

Draba auſtriaca. *Crantz faſc. p.* 12. *t.* 1. *f.* 4.

In ſaxoſis alpium auſtriacarum, ſtyriacarum. Fl. Iunio, Iulio. Per.

Folia lanceolata, baſi attenuata, ciliata; utrinque pilis albidis & apice ſtellatis veſtita, integerrima, aut uno alterove denticulo notata. Flores racemoſi. Petala integra. Iacqu. l. c.

Diverſiſſima a Draba hirta Linn. Vid. Dickſon faſc. 2. p. 29.

mollis. 6. D. foliis ſubcarnoſis, glabris: radicalibus petiolatis ovatis denticulatis; caulinis ſeſſilibus lanceolatis integerrimis, ſiliculis oblongis rectis. *Scop. carn. tom.* 2. *p.* 7. *t.* 34.

In Carniolia in alpibus wochinenſibus. Fl. Iulio. Per.

Petala ſunt apice integra.

LEPIDIUM. Silicula emarginata, cordata, polyſperma: valvis carinatis, contrariis.

perfoliatum. 1. L. foliis caulinis pinnato-multifidis, ramiferis cordatis amplexicaulibus integris. *Sp. pl. p.* 897. *Iacqu. auſtr. t.* 346.

In agri vindobonenſis herbidis, arvis. Fl. Majo, Iunio.

Folia radicalia ac caulina inferiora, in individuis ramoſioribus etiam ramea infima, ſunt bipinnatifida. Horum ſubſequa minus diviſa. Superiora omnia indiviſa, integerrima, cordata, amplexicaulia. Iacqu. l. c.

alpinum. 2. L. foliis pinnatis integerrimis, ſcapo ſubradicato, ſiliculis lanceolatis mucronatis. *Sp. pl. p.* 898. *Iacqu. auſtr. t.* 137.

Lepidium foliis pinnatis, pinnis ovatis æqualibus. *Hall. hiſt n.* 516.

Draba Naſturtiolum. *Scop. carn. tom.* 2. *p.* 8.

Cardamine alpina tertia minima. *Clus. hiſt. p. CXXVIII.*

In alpibus. Fl. æſtate. Per.

petræum. 3. L. foliis pinnatis integerrimis, petalis emarginatis calyce minoribus. *Sp. pl. p.* 899. *Iacqu. auſtr. t.* 131.

Naſturtium foliis pinnatis, radicalibus ovato-lanceolatis, caulinis linearibus. *Hall. hiſt. n.* 515.

In Auſtriæ collibus prope *Bertholdsdorf, Medling, Briel*, alibique. Fl. Martio, Aprili.

ruderale. 4. L. floribus diandris apetalis, foliis radicalibus dentato-pinnatis, ramiferis linearibus integerrimis. *Sp. pl. p.* 900. *Fl. dan. t.* 184.

Naſturtium apetalum, foliis pinnatis, radicalibus pinnis ſemipinnatis, caulinis ſimplicibus. *Hall. hiſt.* n. 518.

Naſturtium ruderale. *Scop. carn. tom.* 2. *p.* 13.

In ruderatis, ad vias. Fl. Majo, Iunio.

Iberis. 5. L. floribus diandris tetrapetalis, foliis inferioribus lanceolatis ferratis, ſuperioribus linearibus integerrimis.. *Sp. pl. p.* 900.

Iberis latiore folio. *Bauh. pin.* 97. *Zann. iſt. p.* 147. *t.* 43.

In Litorali ad mare, vias, hortos. Fl. æſtate. Per.

THLASPI. Silicula emarginata, obcordata, polyſperma: valvis navicularibus, marginato-carinatis.

arvenſe. 1. T. ſiliculis orbiculatis, foliis oblongis dentatis glabris. *Sp. pl. p.* 901. *Fl. dan. t.* 793. *Curtis lond. Scop. carn. tom.* 2. *p.* 18. *Hall. hiſt. n.* 511.

In arvis, ad pagos, vias. Fl. Majo, Iunio.

alliaceum. 2. T. ſiliculis ſubovatis ventricoſis, foliis oblongis obtuſis dentatis glabris. *Sp. pl. p.* 901. *Iacqu. miscell. tom.* 2. *p.* 330. *ic. t.* 121.

In Auſtriæ arvis. *Iacqu.* Fl. Iunio.

ſaxatile. 3. T. ſiliculis ſubrotundis, foliis lanceolato-linearibus obtuſis carnoſis. *Sp. pl. p.* 901. *Iacqu. auſtr. t.* 236.

Naſturtium foliis pulpoſis, ellipticis, ſiliculis alatis emarginatis. *Hall. hiſt. n.* 512.

Thlaspi peregrinum. *Scop. carn. tom.* 2. *p* 17.

Thlaspi parvum ſaxatile, flore rubente. *Bauh. pin.* 107.

In rupeſtribus Litoralis, Carnioliæ calidioris. Fl. a Martio ad Majum. Per.

Folia ſparſa, glauca, carnoſa, adulta rubescentia, oblonga vel lanceolata, infima etiam obverſe ovata, integerrima. Petala alba, lineis roſeis variegata, dum marceſcunt, rubescentia. Siliculæ ſummæ plerumque convexæ & ſubtus concavæ, reliquæ ex adverſo ſuperne concavæ & dorſo convexæ, margine membranaceo amplo & emarginato donantur. Iacqu. l. c.

campeſtre. 4. T. ſiliculis ſubrotundis, foliis ſagitatis dentatis incanis. *Sp. pl. p.* 902. *Curtis lond. Scop. carn. tom.* 2. *p.* 17.

Naſturtium foliis imis petiolatis, ovatis, caulinis ſagittatis, dentatis. *Hall. hiſt. n.* 509.

In cultis, ad agrorum margines, vias. Fl. Majo.

Folia radicalia longe petiolata, oblonga, obtuſa, integra, vel pinnatifida. Caulina ſagittata, dentata, alternatim ſparſa, amplexicaulia.

montanum. 5. T. ſiliculis obcordatis, foliis glabris: radicalibus ſubcarnoſis obovatis ſubintegerrimis, caulinis amplexicaulibus, corollis calyce majoribus. *Sp. pl. p.* 902. *Iacqu. auſtr. t.* 237. *Scop. carn. tom.* 2. *p.* 18.

Lepidium caule erecto, foliis radicalibus petiolatis, caulinis amplexicaulibus ſagittatis. *Hall. hiſt. n.* 518.

Thlaspi alpinum. *Iacqu. auſtr. t.* 238.

Thlaspi præcox. *Wulfen in Iacqu. collect. tom.* 2. *p.* 124. *t.* 9.

Thlaspi montanum ſecundum ſeu badenſe. *Cluſ. hiſt. p. CXXXI.*

In montanis, & alpinis. Fl. pro diverſa elevatione a Martio ad Auguſtum. Per.

Radix per ætatem ſuperne multiceps. Folia ad exortum caulium ſunt plura, in orbem humiſtrata, plana, facie ſaturo & quaſi cæſio virore nitentia, in poſtica parte utplurimum livide purpurea, obovata, aut lanceolata, in petiolum attenuata, integerrima, aut denticulo uno alterove notata, obtuſiſſima, rarius emarginata,

craſſa & firma. Caulina alterna, amplexicaulia. Flores corymboſi. Calyx aut viret, aut purpuraſcit, aut flavet, conſtatque foliolis oblongis, obtuſis concavis. Petala nivea, æqualia, obverſe ovata. Silicula obverſe cordata, ſubtus convexa, ſupra magis minusve concava, glabra, ſtylo perſiſtente munita. Iacqu. l. c.

Cl. Murray in decima quarta Syſt. veget. editione omnia tribuit T. alpino, quæ Linnæus T. montano, foliis caulinis exceptis, quæ in illo ſimpliciter ſeſſilia eſſe dicit, dum utrique ſeſſilia & amplexicaulia ſunt.

perfoliatum. 6. T. foliis imis ovatis, reliquis amplexicaulibus cordatis. *Hall. hiſt. n.* 510. ſub Naſturio. *Sp. pl. p.* 902. *Iacqu. auſtr. t.* 337.

Thlaspi alpeſtre. *Sp. pl. p.* 903.

Thlaspi tertium pumilum. *Clus. hiſt. p. CXXXI.*

In Auſtriæ, Pannoniæ cultis, collibus, ad ſepes, alibique. Fl. Martio, Aprili.

Cl. Linnæus in ſpeciebus Plantarum ut T, alpeſtre a perfoliato diſtingueret, huic caulem ramoſum, ſimplicem illi eſſe dicit, citatque ſub alpeſtri Thlaspi pumilum tertium Cluſii, quod eſt perfoliati varietas minor, in collibus apricis aridisque Auſtriæ creſcens. Citat quoque Thlaspi perfoliatum minus Caſpari Bauhini, nempe pumilum tertium Cluſii, tam ſub montano, quam ſub alpeſtri.

Burſa Paſtoris. 7. T. ſiliculis obcordatis, foliis radicalibus pinnatifidis. *Sp. pl. p.* 903. *Curtis lond. Fl. dan. t.* 729. *Scop. carn. tom.* 2. *p.* 17.

Naſturtium ſiliquis triangularibus. *Hall. hiſt. n.* 514.

Le Tabouret, ou la Bourſe a Paſteur. Regn. bot.

In cultis, & incultis. Fl. a primo vere ad hyemem.

Folia radicalia in aliis plantis pinnatifida, caulina dentata; in aliis omnia integra dentataque.

COCHLEARIA. Silicula emarginata, turgida, ſcabra: valvis gibbis, obtuſis.

Coronopus, 1, C. foliis pinnatifidis, caule depreſſo. *Syſt. veget. p.* 588. *Fl. dan. t.* 202. *Scop. carn. tom.* 2. *p.* 12. *Hall. hiſt. n.* 502.

Naſturtium ſylveſtre capſulis criſtatis. *Tourn. inſt.* 214. *Zann. iſt. p.* 192. *t.* 115.

In Auſtriæ, Pannoniæ, Croatiæ, Litoralis uliginoſis, cultis, ad vias, foſſas, Fl. a Majo ad Iulium.

Armoracia, 2, C. foliis radicalibus lanceolatis crenatis, caulinis inciſis. *Sp. pl. p.* 904. *Hall. hiſt. n.* 504.

Raphanus ruſticanus. *Bauh. pin.* 96.

Le Raifort ſauvage. Regn. bot.

In Auſtria ad pagos, vias, foſſas, Fl. Majo, Iunio. Per.

Draba, 3, C. foliis caulinis lanceolatis dentatis amplexicaulibus. *Sp. pl. p.* 904. *Iacqu. auſtr. t.* 315.

Draba prima vulgaris. *Clus. hiſt. p.* CXXIV.

In Auſtria, Pannonia in ruderatis, cultis, ad vias, Fl. a Majo ad Iulium. Per.

IBERIS. Corolla irregularis: petalis duobus exterioribus majoribus. Silicula polyſperma, emarginata.

rotundifolia, 1, I. herbacea, foliis ovatis: caulinis amplexicaulibus lævibus ſuccoſis. *Sp. pl. p.* 905. *Scop. carn. tom.* 2. *p.* 15. *t.* 37.

Lepidium caule repente, foliis ovatis, amplexicaulibus. *Hall. hiſt. n.* 517.

In ſaxoſis alpium carniolicarum, carinthiacarum, tyrolenſium. Fl. Iulio, Auguſto. Per.

cepeæfolia. 2. I. herbacea, foliis ovatis: caulinis ſeſſilibus lævibus ſuccoſis. *Syſt. veget. p.* 589. *Wulfen in Iacqu. miſcell. tom.* 2. *p.* 28. *t.* 1.

Thlaspi montanum, ſerrato Cepeæ folio, flore purpurascente umbellato. *Barr. rat. p.* 38. *n.* 365. *ic.* 848.

Thlaspi alpinum, folio rotundiore, carnoſo, purpurascente flore. *Tourn. inſt.* 212.

In Carinthia in valle *Rabl* ad fodinas calaminares *unter der Vitriolwand. Wulfen.* Fl. Majo.

umbellata. 3. I. herbacea, foliis lanceolatis acuminatis; inferioribus ſerratis; ſuperioribus integerrimis. *Sp. pl. p.* 906. *Curtis magaz.* 106.

Le Thlaspi de Crete. Regn. bot.

In Litorali prope Duinum. Fl. æſtate.

Fructus hujus corymboſus hemisphæricus perſiſtit, I. amaræ racemoſus evadit. Linn. l. c.

amara. 4. I. herbacea, foliis lanceolatis acutis ſubdentatis, floribus racemoſis. *Sp. pl. p.* 906. *Scop. carn. tom.* 2. *p.* 16.

In Carniolia circa *Heidenſchaft. Scop.* Fl. Iunio, Iulio.

pinnata. 5. I. herbacea, foliis linearibus pinnatifidis. *Sp. pl. p.* 907.

Thlaspi alterum minus umbellatum, naſturtii hortenſis folio, narbonenſe. *Lob. ic.* 218.

Tergeſti *alle Saule. I. Hoſt.* Fl. Iunio, Iulio.

Tota planta pilis inſperſa. Folia inferiora pinnatifida, ſuperiora trifida vel bifida, ſumma integra.

ALYSSUM. Filamenta quædam introrſum denticulo notata. Silicula emarginata.

monta-num. 1. A. ramulis ſuffruticoſis diffuſis, foliis punctato-echinatis. *Sp. pl. p.* 907. *Iacqu. auſtr. t.* 37. *Scop. carn. tom.* 2. *p.* 14.

Alyſſum ſubfruticoſum foliis ellipticis ſcabris. *Hall. hiſt. n.* 492.

Alyſſum alpeſtre. *Wulfen in Iacqu. collect. tom.* 4. *p.* 227. *t.* 4. *f.* 1.

Thlaſpi montanum incanum luteum, *Bauh. pin.* 107, *prodr.* 49.

In collibus aridis, montoſis apricis, & in alpibus. Fl. æſtate. Per.

Tota planta, corollis exceptis, veſtita eſt tomento albido, quod conſtat ex ſtellulis plurimis. Folia ſunt obverſe lanceolata, acutiuſcula, aut obtuſa. Siliculæ aliæ ovatæ, quædam obovatæ, multæ orbiculatæ, nonnullæ apice parum emarginatæ.

ſaxatile. 2. A. caulibus frutescentibus paniculatis, foliis lanceolatis molliſſimis undulatis integris. *Sp. pl. p.* 908. *Mill. dict. t.* 20. *f.* 1.

In Auſtria in ſaxoſis prope *Krems. Iacqu.* Fl. æſtate. Per.

incanum. 3. A. caule herbaceo erecto, foliis lanceolatis incanis integerrimis, floribus corymboſis, petalis bifidis. *Sp. pl. p.* 908.

Thlaſpi incanum machlinenſe. *Clus. hiſt. p.* CXXXII.

In pratis, ruderatis, collibus, ſecus vias, alibique. Fl. æſtate.

calyci-num. 4. A. caulibus herbaceis, ſtaminibus omnibus dentatis, calycibus perſiſtentibus. *Sp. pl. p.* 908. *Iacqu. auſtr. t.* 338.

Alyſſum foliis ellipticis ſcabris, calycibus perſiſtentibus. *Hall. hiſt. n.* 494.

Adyſeton calycinum. *Scop. carn. tom.* 2. *p.* 13.

Alyſſon minimum. *Cluſ. hiſt. p.* CXXXIII.

In campis, arvis, collibus, ad vias. Fl. æſtate.

gemonenſe. 5. A. caule herbaceo, ramis divaricatis, foliis radicalibus obovatis ſubtomentoſis, ſiliculis inflatis. *Syſt. veget. p.* 591. *Wulfen in Iacqu. collect. tom.* 2. *p.* 159. *ic. t.* 503.

Alyſſum petræum, caulibus ſuperne ramoſis, ramis divaricatis flexuoſis; ſiliculis ſubinflatis, tetraſpermis. *Arduin. animadv.* 2. *p.* 30. *t.* 14.

In Carnia prope Gemonam. In Carniolia *in der Wochein*, prope *Neumarktl*, in monte *Nanas*, ubique locis ſaxoſis. Fl. æſtate. Per.

CLYPEOLA. Silicula emarginata, orbiculata, compreſſo-plana, decidua.

Ionthlaſpi. 1. C. ſiliculis orbiculatis unilocularibus monoſpermis. *Syſt. veget. p.* 591.

Thlaspi clypeatum, ſerpylli folio. *Bauh. pin.* 107. *Zann. iſt. p.* 148. *t.* 274.

Ionthlaspi minimum ſpicatum lunatum. *Col. ecphr.* 1. *p.* 281. *t.* 284.

In arenoſis Litoralis, inſularumque venetarum. Fl. primo vere.

maritima. 2. C. perennis, ſiliculis bilocularibus ovatis diſpermis. *Syſt. veget. p.* 426.

Alyſſon caulibus diffuſis, foliis linearibus. *Ger. prov. p.* 350.

Thlaspi montanum perenne flore albo. *Barr. rar.* 844.

In aſperis Litoralis ad mare. Fl. Iunio, Iulio. Per.

PELTARIA. Silicula integra, ſuborbiculata, compreſſo-plana, non dehiſcens.

alliacea. 1. P. foliis amplexicaulibus oblongis indiviſis. *Syſt. veget. p.* 591. *Iacqu. auſtr. t.* 123. *Scop. carn. tom.* 2. *p.* 4.

Thlaſpi montanum primum peltatum. *Clus. hiſt. p.* CXXX.

In Auſtria in rupeſtribus ad arcem *Stückſenſtein.* In agro fluminenſi in declivibus montis Terſacti, alibique. Fl. Aprili, Majo. Per.

Tota planta allium redolet. Folia primi anni longe petiolata, ovata vel magis ſubrotunda, obtuſa, integra, glabra. Folia ſecundi anni caulina ovata vel magis lanceolata, acutiuſcula, amplexicaulia, alterna, infima quandoque in petiolum attenuata. Iacqu. l. c.

BISCUTELLA. Calycis foliola baſi gibba. Silicula compreſſo-plana, rotundata, ſupra infraque biloba.

lævigata. 1. B. ſiliculis glabris, foliis lanceolatis ſerratis. *Syſt. veget. p.* 592. *Iacqu. auſtr. t.* 339.

Biscutella ſilicula lævi, foliis dentatis aſperis. *Hall. hiſt. n.* 501.

Biscutella didyma. *Scop. carn. tom.* 2. *p.* 14.

Thlaſpi clypeatum. *Clus. hiſt. p.* CXXXIII.

In montanis ſylvaticis, arenoſis, in alpinis, & ſubalpinis. Fl. æſtate. Per.

LUNARIA. Calyx tetraphyllus: foliolis duobus alternis baſi gibbis ſaccatisque. Silicula compreſſo-plana, integra, pedicellata: valvis diſſepimento æqualibus, parallelis, planis.

rediviva. 1. L. ſiliculis lanceolatis, radice perenni. *Retzii ſcand. p.* 154. *obs. faſc.* 1. *p.* 22. *Sp. pl. p.* 911. *Scop. carn. tom.* 2. *p.* 5.

Lunaria perennis, ſiliquis ovato-lanceolatis. *Hall. hiſt. n.* 500.

Viola latifolia, lunaria odorata. *Cluſ. hiſt. p.* 297.

In alpinis, & ſubalpinis. Fl. Majo, Iunio, Iulio. Per.

Differt a Lunaria annua, cui radix biennis, ſiliculæ ellipticæ. Retz. l. c.

SILIQUOSA.

DENTARIA. Calyx longitudinaliter connivens. Siliqua elongata, teres, emarginata, elaſtice revolvens valvulas. Semina in foveis ſepti fungoſi nidulantia.

enneaphylla. 1. D. foliis ternis ternatis. *Sp. pl. p.* 912. *Iacqu. auſtr. t.* 316. *Scop. carn. tom.* 2. *p.* 19.

Dentaria quinta triphyllos. *Cluſ. hiſt. p.* CXXI.

In ſylvis montanis, & ſubalpinis. Fl. ab Aprili ad Iunium. Per.

bulbifera. 2. D. foliis inferioribus pinnatis, ſummis ſimplicibus. *Sp. pl. p.* 912. *Fl. dan. t.* 361. *Scop. carn. tom.* 2. *p.* 19. *Hall. hiſt. n.* 470.

Dentaria quarta baccifera. *Cluſ. hiſt. p.* CXXI.

Cum præc. Fl. Aprili, Majo. Per.

pentaphyllos. 3. D. foliis ſummis digitatis. *Sp. pl. p.* 912. *Scop. carn. tom.* 2. *p.* 20.

Dentaria foliis quinis digitatis. *Hall. hiſt. n.* 469.

Dentaria ſexta, pentaphyllos prima. *Clus. hiſt. p.* CXXII.

Dentaria ſeptima, pentaphyllos ſecunda. *Clus. hiſt. p.* CXXII.

Dentaria octava, heptaphyllos. *Clus. hiſt. p.* CXXIII.

In Carnioliæ, Croatiæ, Litoralis ſylvis montanis, & ſubalpinis. Fl. Aprili, Majo. Per.

Variat foliis pinnatis, digitatis; foliolis æqualibus, inæqualibus, ſimpliciter vel duplicato-ſerratis.

CARDAMINE. Calyx ſubhians. Stigma integrum. Siliqua elaſtice diſſiliens valvulis vel revolutis, vel a baſi tantum dehiſcentibus.

hetero-phylla. 1. C. foliis radicalibus ovatis, caulinis integris lobatis pinnatifidisve: petiolis longis.

Cardamine foliis radicalibus petiolatis, ovatis, caulinis ovatis, aut pinnatis, obtuſis. *Hall. hiſt. n.* 476.

Cardamine reſedifolia. *Sp. pl. p.* 913.

Cardamine bellidifolia. *Wulfen in Iacqu. miscell. tom.* 1. *p.* 148.

Naſturtium alpinum minus, reſedæ folio. *Bauh. pin.* 104. *prodr.* 45.

In alpibus Styriæ, Carinthiæ, Tyrolis. Fl. Iulio, Auguſto, Septembri. Per.

In aliis caulibus folia omnia integra; in aliis pauca obſolete lobata; in nonnullis radicalia caulinaque inferiora integra, ſuperiora autem lobata, pinnatifidaque: lobis & pinnis integris, vel diviſis. Obſervavi hanc foliorum diverſitatem non tantum in caulibus ſingulis e ſingulis radicibus, ſed in pluribus quoque ex eadem radice, ortis. An C. bellidifolia Linn. huc pertineat, aſſerere non audeo.

petræa. 2. C. foliis ſimplicibus oblongis dentatis. *Sp. pl. p.* 913. *Fl. dan. t.* 386. *Hænke in Iacqu. collect. tom.* 2. *p.* 24.

Arabis *hispida*, foliis hispidis: radicalibus runcinatis; caulinis lanceolatis. *Linn. ſuppl. p.* 298. *Syſt. veget. ed.* 13. *p.* 501. *ed.* 14. *p.* 600.

Cardamine petræa cambrica, naſturtii facie. *Dill. elth.* 70. *t.* 61. *f.* 71.

In rupeſtribus Auſtriæ, Styriæ. Fl. a primo vere ad Autumnum. Per.

Folia glabra, aut hirſuta: radicalia pinnatifida, vel dentata; caulina linearia.

trifolia. 3. C. foliis ternatis obtuſis, caule ſubnudo. *Sp. pl. p.* 913. *Iacqu. auſtr. t.* 27. *Scop. carn. tom.* 2. *p.* 20. *Hall. hiſt. n.* 475.

Cardamine alpina prima trifolia. *Clus. hiſt. p.* CXXVII.

In ſylvis umbroſis ſubalpinis. Fl. Iunio. Per.

chelidonia. 4. C. foliis pinnatis: foliolis quinis inciſis. *Sp. pl. p.* 914.

Cardamine glabra, chelidonii folio. *Tourn. inſt.* 225. *Barr. ic.* 156.

Naſturtium pyrenæorum aquaticum latifolium, purpurascente flore. *Herm. par.* 203. *t.* 204.

Naſturtium alpinum, Halicacabi peregrini foliis. *Ambros.* 374.

In Litorali in udis montis majoris. *I. Hoſt.* Fl. Iunio, Iulio. Per.

Caulis tetragonus. Folia longe petiolata, pinnata: pinnulis ciliatis, lobatis, quandoque una alterave auricula auctis.

impatiens. 5. C. foliis pinnatis inciſis ſtipulatis, floribus apetalis, *Sp. pl. p.* 914. *Scop. carn. tom.* 2. *p.* 21.

Cardamine foliis pinnatis, imis obtuſe lobatis, caulinis acute dentatis, floribus apetalis. *Hall. hiſt. n.* 471.

In ſylvis umbroſis, alibique. Fl. ab Aprili ad Iunium.

hirſuta. 6. C. foliis pinnatis, floribus tetrandris hexandrisque. *Sp. pl. p.* 915. *Curtis lond. Scop. carn. tom.* 2. *p.* 21. *t.* 38.

Cardamine caule foliisque pinnatis hirſutis, ſubrotundis. *Hall. hiſt. n.* 472.

Naſturtium aquaticum minus. *Bauh. pin.* 104. *Zann. iſt. p.* 49. *t.* 225.

Ad fluvios, torrentes, ſepes, in cultis, ſylvis ſubalpinis, alibique. Fl. primo vere.

pratenſis. 7. C. foliis pinnatis: foliolis radicalibus ſubrotundis: caulinis lanceolatis. *Sp. pl. p.* 915. *Curtis lond. Scop. carn. tom.* 2. *p.* 23.

Cardamine foliis pinnatis, radicalibus ſubrotundis, caulinis linearibus. *Hall. hiſt. n.* 473.

Cardamine altera ſimplici flore. *Clus. hiſt. p.* CXXVIII.

Cardamine altera pleno flore. *Clus. hiſt. p.* CXXIX.

In pratis udis, ad rivulos, fluvios, foſſas. Varietatem flore pleno legi in pratis Auſtriæ paluſtribus prope *Himberg*. Fl. Aprili, Majo. Per.

In ſummis alpium judenburgenſium jugis inveni varietatem nanam, folio radicali uno alterove indiviſo, C. reſedifoliæ wulfenianæ, quam t. 21. *Flore auſtriacæ app. ſiſtit, ſimillimam.*

amara. 8. C. foliis pinnatis, axillis ſtoloniferis. *Sp. pl. p.* 915. *Curtis lond.*

Cardamine foliis pinnatis, ſubrotundis, anguloſis. *Hall. hiſt. n.* 474.

Naſturtium aquaticum majus & amarum. *Bauh. pin.* 104. *prodr.* 45.

Ad fluvios, rivulos, & in aliis udis locis demiſſis, ſubalpinis, & alpinis. Fl. Majo, Iunio. Per.

Huic ſtylus obliquus, præcedenti erectus.

SISYMBRIUM. Siliqua dehiſcens valvulis rectiuſculis. Calyx & corolla patens.

Naſturtium. 1. S. ſiliquis declinatis, foliis pinnatis: foliolis ſubcordatis. *Sp. pl. p.* 916. *Curtis lond. Fl. dan. t.* 690. *Scop. carn. tom.* 2. *p.* 24.

Siſymbrium foliis pinnatis, pinnis ſubrotundis, ſpicis ſeſſilibus. *Hall. hiſt. n.* 482.

Le Creſſon de fontaine. Regn. bot.

In aquis ſtagnantibus, vel lente fluentibus. Fl. Majo, Iunio. Per.

Sylveſtre. 2. S. ſiliquis declinatis, foliis pinnatis: foliolis lanceolatis ſerratis. *Sp. pl. p.* 916. *Curtis lond.*

Siſymbrium foliis pinnatis, pinnis dentatis diſſitis. *Hall. hiſt. n.* 485.

Naſturtium islandicum. *Fl. dan. t.* 409.

In ruderatis, ad vias, foſſas. Fl. æſtate. Per.

Folia radicalia pinnatifida: pinnis ſubovatis dentato-ſerratis. Caulina alterna, ſubpinnatifida: pinnis lanceolatis, ſerratis integerrimisque. Curtis. l. c.

amphibium. 3. S. ſiliquis declinatis oblongo-ovatis, foliis pinnatifidis ſerratis. *Sp. pl. p.* 917. *Fl. dan. t.* 931. 984.

Radicula foliis integris & pinnatifidis, petalis calyce longioribus. *Hall. hiſt. n.* 486.

In aquis ſtagnantibus, paluſtribus, & ſiccis. Fl. æſtate. Per.

In aquis creſcens habet folia ſimplicia, dentata, ſerrata; in paluſtribus pinnatifida: pinnis ſerratis; in ſiccis inferiora ſimplicia, ſuperiora pinnata.

tenuifolium. 4. S. foliis integerrimis: infimis tripinnatifidis; ſupremis integris. *Sp. pl. p.* 917.

La Roquette ſauvage. Regn. bot.

In muris, cultis, ruderatis, ad vias. Fl. æſtate. Per.

murale. 5. S. ſubacaule, foliis lanceolatis ſinuato - ſerratis læviuſculis, ſcapis ſubſcabris adſcendentibus. *Sp. pl. p.* 918.

Eruca ſylveſtris major caule aſpero. *Bauh. pin.* 98. *Zann. iſt. p.* 95. *t.* 178.

Habitat & floret cum præc.

arenoſum. 6. S. caule ſubfolioſo ramoſo, foliis lyratis rectangulo - dentatis hiſpidis: pilis ramoſis. *Sp. pl. p.* 919.

Eruca foliis ſcabris, ſemipinnatis, pinnis rectangulis, rariter dentatis. *Hall. hiſt. n.* 462.

Arabis arenoſa. *Scop. carn. tom.* 2. *p.* 52. *t.* 40.

Eruca cœrulea in arenoſis creſcens. *Bauh. pin.* 99. *prodr.* 40.

In arenoſis ad fluvios, torrentes, ad pedes montium, in rupeſtribus convallium alpinarum, & aſcendit ad ipſas alpes. Fl. æſtate. Per.

Caules modo ſimplices, modo ramoſi. Corollæ in aliis individuis albæ, violaceæ in aliis. In Tyroli in Sellrainerthal, *& in Styria in alpibus judenburgenſibus inveni varietatem caule foliisque indiviſis glabris, ab Arabi ovirenſi Wulfenii non multum abſimilem.*

Sophia. 7. S. petalis calyce minoribus, foliis decompoſito-pinnatis. *Sp. pl. p.* 920. *Fl. dan. t.* 528. *Scop. carn. tom.* 2. *p.* 24.

Siſymbrium pinnis & pinnulis pinnatis, laciniis linearibus ſubhirſutis. *Hall. hiſt. n.* 484.

Sophia chirurgorum. *Lob. ic. p.* 738.

Le Siſimbre parviflore. Bulliard herb. t. 271.

In cultis, ruderatis, ad vias. Fl. Aprili, Majo.

Irio 8. S. foliis runcinatis dentatis nudis, caule lævi, ſiliquis erectis. *Sp. pl. p.* 921. *Iacqu. auſtr. t.* 322.

In cultis, ad agrorum margines. Fl. a Majo ad Iulium.

Columnæ. 9. S. caule foliisque runcinatis dentatis villoſis ſubincanis, ſiliquis erectis. *Syſt. veget. p.* 596. *Iacqu. auſtr. t.* 323.

In ſterilibus ſiccis, ad vias. Fl. æſtate.

Læſelii. 10. S. foliis runcinatis acutis hirtis, caule hiſpido. *Sp. pl. p.* 920. *Iacqu. auſtr. t.* 324.

Eryſimum hirſutum, ſiliqua Erucæ. *Læs. pruſſ.* 69. *t.* 4.

In Auſtria, Pannonia ad aggeres, vias, in cultis, ruderatis, paluſtribus. Fl. a Majo ad Iulium.

ſtrictiſſimum. 11. S. foliis lanceolatis dentato-ſerratis caulinis. *Sp. pl. p.* 922. *Iacqu. auſtr. t.* 194.

Eruca caule paniculato, foliis nervoſis, ovato-lanceolatis. *Hall. hiſt. n.* 458.

Draba lutea, ſiliquis ſtrictiſſimis. *Bauh. pin.* 110.

Heſperis lutea, ſiliquis ſtrictiſſimis. *Tourn. inſt.* 222.

Ad ſepes, in ſylvaticis Auſtriæ, Styriæ, Pannoniæ. Fl. a Majo ad Iulium. Per.

ERYSIMUM. Siliqua columnaris, exacte tetraëdra. Calyx clauſus.

officinale. 1. E. ſiliquis ſpicæ adpreſſis, foliis runcinatis. *Sp. pl. p.* 923. *Curtis lond. Fl. dan. t.* 560.

Eryſimum foliis pinnatis, pinnis rectangulis acutis, extrema triangulari maxima, ſiliquis adpreſſis. *Hall. hiſt. n.* 478.

Siſymbrium officinale. *Scop. carn. tom.* 2. *p.* 26.

Le Velar, ou la Tortelle. Regn. bot.

Ad vias, pagos, in ruderatis. Fl. æſtate.

Barbarea. 2. E. foliis lyratis: extimo ſubrotundo. *Sp. pl. p.* 922. *Fl. dan. t.* 985.

Eryſimum foliis pinnatis, pinna extrema maxima, ſubrotunda. *Hall. hiſt. n.* 479.

Siſymbrium Barbarea. *Scop. carn. tom.* 2. *p.* 26.

L' Herbe de Sainte Barbe. Regn. bot.

In inundatis, ad fluvios, foſſas, alibique. Fl. Majo, Iunio. Per.

Alliaria. 3. E. foliis cordatis. *Sp. pl. p.* 922. *Curtis lond. Fl. dan. t.* 935. *Hall. hiſt. n.* 480.

Siſymbrium Alliaria. *Scop. carn. tom.* 2. *p.* 26.

L' Alliaire. Regn. bot.

In cultis, ad ſepes, in ſylvis. Fl. Majo.

repandum. 4. E. foliis lanceolatis dentatis, racemis oppoſitifoliis, ſiliquis racemoſis ſubſeſſilibus, corollis minutis. *Sp. pl. p.* 923. *Iacqu. auſtr. t.* 22.

In cultis, ruderatis, ad vias. Fl. a Majo ad Iunii finem.

cheiranthoides. 5. E. foliis lanceolatis integerrimis, ſiliquis patulis. *Syſt. veget. p.* 597. *Iacqu. auſtr. t.* 23. *Fl. dan t.* 731.

In cultis, ruderatis, campeſtribus Auſtriæ. Fl. Iunio, Iulio.

hieracifolium. 6. E. foliis lanceolatis ſerratis. *Sp. pl. p.* 923. *Iacqu. auſtr. t.* 73.

In ſcrobibus glareoſis, ad latera viarum, in agris lapidoſis. Fl. æſtate.

Folia oblongo-lanceolata, baſi attenuata, ſubſeſſilia, acuta, raris denticulis ad oras munita, interdum profunde ſinuata. Flores odorem ſuavem ſpirant. Siliquæ exacte tetragonæ. Racemi valde producuntur, denſeque tandem toti teguntur fructibus; qua ſola nota facile ab affinibus diſtinguitur. Tum præterea ab Eryſimo cheiranthoide in

primis florum odore & magnitudine, petalis magis ſubrotundis & foliis verſus apicem minus attenuatis. A Cheirantho eryſimoide ſiliquarum coſtis omnibus rectangulis, radiceque annua. A Cheirantho alpino caulibus magis argute angulatis, flore paullo majore, foliis non incanis. Iacqu. l. c.

CHEIRANTHUS. Germen utrinque denticulo glandulato. Calyx clauſus: foliolis duobus baſi gibbis. Semina plana.

eryſimoides. 1. C. foliis lanceolatis dentatis nudis, caule erecto ſimpliciſſimo, ſiliquis tetragonis. *Sp. pl. p.* 923. *Iacqu. auſtr. t.* 74. *Fl. dan. t.* 229.

Leucojum ſylveſtre. *Cluſ. hiſt. p.* 299.

In apricis aſperis. Fl. Majo.

Folia ſunt anguſta, acutiuſcula, ſeſſilia, obſcure virentia, plerumque parum conniventia; in aliis ferme linearia, in aliis ex oblongo lanceolata, ſæpius integerrima, inferiora tamen in quibusdam individuis denticulata. Stylus nullus. Stigma craſſum immediate germini imponitur. Iacqu. l. c.

helveticus. 2. C. foliis lanceolatis dentatis nudis, caule erecto, ſiliqua tetragona, ſtylo acuminata. *Syſt. veget. p.* 597. *Iacqu. hort. tom.* 3. *p.* 9. *t.* 9.

In alpibus tyrolenſibus. Fl. Iulio. Per.

Folia ſunt anguſte lanceolata, integerrima, vel raro obiter dentata. Stigma per ſtylum perpetuo intermedium diſtat a germine. Iacqu. l. c.

alpinus. 3. C. foliis linearibus integris ſubtomentoſis, caule ramoſo. *Syſt. veget. p.* 597. *Iacqu. auſtr. t.* 75.

In aggeribus, apricis aſperis Auſtriæ, Pannoniæ. Fl. Iunio, Iulio.

triſtis. 4. C. foliis linearibus ſubſinuatis, floribus ſeſſilibus, petalis undatis, caule ſuffruticoſo. *Sp. pl. p.* 925.

Leucojum minus, lavendulæ folio, obſoleto flore. *Bocc. muſ.* 148. *t.* 111. *Barr. ic.* 83.

In Carnia in arenoſis *al Tagliamento. I. Hoſt.* Fl. Iulio, Auguſto. Per.

HESPERIS. Petala oblique flexa. Glandula intra ſtamina breviora. Siliqua ſtricta. Stigma baſi bifurca apice connivente. Calyx clauſus.

triſtis. 1. H. caule hiſpido ramoſo patente. *Sp. pl. p.* 927. *Jacqu. auſtr. t.* 122.

Heſperis prima, *Cluſ. hiſt. p.* 296.

In Auſtria, Pannonia ad margines ſylvarum, agrorum, alibique. Fl. Majo.

ſylveſtris. 2. H. caule ſimplici erecto, foliis ſubhaſtatis dentatis, petalis obtuſis. *Crantz faſc. p.* 32.

Heſperis caule hirſuto, foliis ovato-lanceolatis, ſerratis, petiolatis, ſubaſperis, *Hall. hiſt. n.* 448.

Heſperis inodora. *Sp. pl. p.* 927. *Iacqu. auſtr. t.* 347.

Heſperis tertia. *Cluſ. hiſt. p.* 297.

In agro vindobonenſi in ſylvaticis, ad montium radices, vineas. Fl. Majo, Iunio.

Folia acuminata, utrinque aſperiuſcula & villoſa, argute dentato-ſerrata, late aut anguſte lanceolata, conferta, infima plerumque petiolata, interdum omnia; in aliis ex adverſo ſeſſilia omnia, vel ſemiamplexicaulia, hinc baſi aut rotundata, aut ſubhaſtata. Flores, præſertim veſpere, ſuavi odore pollentes. Iacqu. l. c.

ARABIS. Glandulæ nectariferæ quatuor, ſingula intra calycis foliolum ſquamæ inſtar reflexæ. Stigma ſimplex. Siliqua linearis, toroſa.

alpina. 1. A. foliis amplexicaulibus dentatis. *Sp. pl. p.* 928. *Fl. dan. t.* 62. *Scop. carn. tom.* 2. *p.* 29.

Arabis caule ramoſo, folioſo, diffuſo, foliis amplexicaulibus dentatis, incanis. *Hall. hiſt. n.* 451.

Draba tertia ſucculenta. *Cluſ. hiſt. p.* CXXV.

In alpinis, & ſubalpinis. Fl. æſtate. Per.

thaliana. 2. A. foliis petiolatis lanceolatis ſubintegerrimis. *Sp. pl. p.* 929. *Curtis lond. Retzii obs. faſc.* 1. *p.* 22.

Arabis foliis radicalibus petiolatis, ovatis, dentatis, caule ſubnudo, ramoſo. *Hall. hiſt. n.* 452.

In arvis, collibus herbidis, in ſylvaticis. Fl. primo vere, in Tyroli prope Oenopontum quandoque & Autumno.

Pars inferior caulis hirta eſt, ſuperior glabra. Folia radicalia oblongo-ovata, petiolata, caulina ſeſſilia: omnia integerrima, aut dentata, utrinque punctis prominulis ſcabra, plus vel minus hirſuta: pilis ſimplicibus, aut ramoſis.

bellidifolia. 3. A. foliis ſubdentatis: radicalibus obovatis; caulinis lanceolatis. *Syſt. veget. p.* 599. *Iacqu. auſtr. t.* 280. 281.

Leucojum foliis læviſſimis, rariter dentatis, imis petiolatis ovatis, caulinis amplexicaulibus. *Hall. hiſt. n.* 446.

Leucojum foliis radicalibus ſubrotundis, ſcabris, dentatis, caulinis amplexicaulibus hirſutis. *Hall. hiſt. n.* 447.

Arabis cœrulea. *Hœnke in Iacqu. collect. tom.* 2. *p.* 56.

Turritis bellidifolia. *Allion. pedem. tom.* 1. *p.* 270. *t.* 40. *f.* 1.

Turritis cœrulea. *Allion. pedem. tom.* 1. *p.* 270. *t.* 40. *f.* 2.

Cardamine bellidifolia. *Allion. pedem. tom.* 1. *p.* 260. *t.* 18. *f.* 3.

Plantula cardamines alterius æmula. *Cluſ. hiſt. p.* CXXIX.

In alpinis, & ſubalpinis. Fl. æſtate. Per.

Folia radicalia ſunt lanceolata, aut obverſe ovata. Caulina ſeſſilia, aut amplexicaulia. Omnia ciliata, apice integerrima, vel dentata: in planta in humidis creſcente, ut & caulis, glabra, aut raris pilis inſperſa; in aſperis vero habitante punctis elevatis in pilos deſinentibus ſcabra. Pili ſunt ſimplices, vel ramoſi. Flores albi, aut cærulei.

Turrita. 4. A. foliis amplexicaulibus, ſiliquis decurvis planis linearibus, calycibus ſubrugoſis. *Sp. pl. p.* 930. *Iacqu. auſtr. t.* 11.

Turrita major. *Cluſ. hiſt. p.* CXXVI.

In rupeſtribus umbroſis, ſylvaticis. Fl. Aprili, Majo.

In A. Turrita caulis tenuiter villoſus, in A. pendula hiſpidus. Folia in pendula ſerrata, in turrita dentata. Turrita gerit ſiliquas minime pendulas; ſed pedicelli erecti ſunt, & ſiliquæ curvantur; ſunt compreſſæ, margine inæquali & fere toruloſæ. Semina triplo majora, fuſco nigricantia, reniformia, marginata. In pendula ſiliquæ ſunt pendulæ, lineares, margine æquales. Semina minuta, compreſſiſſima, oblonga, minime emarginata, flava. Retzii obs. fasc. 3. p. 36.

TURRITIS. Siliqua longiſſima, angulata. Calyx connivens, erectus. Corolla erecta.

glabra. 1. T. foliis radicalibus dentatis hiſpidis; caulinis integerrimis ampelxicaulibus glabris. *Sp. pl. p.* 930. *Fl. dan. t.* 809. *Curtis lond. Hall. hiſt. n.* 455.

In ſylvaticis, & cultis. Fl. Majo, Iunio.

hirſuta. 2. T. foliis omnibus hiſpidis, caulinis amplexicaulibus. *Sp. pl. p.* 930. *Iacqu. collect. tom.* 1. *p.* 70. *ic. t.* 126. *Hall. hiſt. n.* 456.

Turrita vulgatior. *Cluſ. hiſt. p.* CXXVI.

Eryſimo ſimilis hirſuta, non laciniata alba. *Bauh. pin.* 101. *prodr.* 42.

In paſcuis montanis, agris, alibique. Fl. cum præc.

BRASSICA. Calyx erectus, connivens. Semina globoſa. Glandula inter ſtamina breviora & piſtillum, interque longiora & calycem.

orientalis. 1. B. foliis glabris: radicalibus petiolatis; caulinis amplexicaulibus, ſiliquis tetragonis. *Iacqu. auſtr. t.* 282.

Braſſica campeſtris prima. *Cluſ. hiſt. p.* CXXVII.

In agri vindobonenſis cultis. Fl. Majo.

SINAPIS. Calyx patens. Corollæ ungues recti. Glandula inter ſtamina breviora & piſtillum, interque longiora & calycem.

arvenſis. 1. S. ſiliquis multangulis toroſo-turgidis lævibus roſtro ancipiti longioribus. *Syſt. veget. p.* 602. *Curtis lond. Fl. dan. t.* 753. *Scop. carn. tom.* 2. *p.* 34. *Hall. hiſt. n.* 467.

In agris, inter ſegetes. Fl. Iunio.

alba. 2. S. ſiliquis hiſpidis: roſtro obliquo longiſſimo enſiformi. *Syſt. veget. p.* 602. *Curtis lond. Scop. carn. tom.* 2. *p.* 35. *Hall. hiſt. n.* 466.

In cultis, ruderatis. Fl. Iunio.

nigra. 3. S. ſiliquis glabris racemo appreſſis. *Syſt. veget. p.* 602.

Sinapis ſiliquis tetragonis glabris. *Hall. hiſt. n.* 465.

La Moutarde. Regn. bot.

In ruderatis, cultis. Fl. Iulio.

RAPHANUS. Calyx clauſus. Siliqua toroſa, ſubarticulata, teres. Glandulæ melliferæ duæ inter ſtamina breviora & piſtillum, totidem inter ſtamina longiora & calycem.

Raphaniſtrum. 1. R. ſiliquis teretibus articulatis lævibus unilocularibus. *Sp. pl. p.* 935. *Curtis lond. Hall. hiſt. n.* 678.

Sinapis arvenſis. *Fl. dan. t.* 678.

In arvis, inter ſegetes. Fl. æſtate.

MONADELPHIA.

PENTANDRIA.

ERODIUM. Calyx pentaphyllus. Corolla pentapetala. Squamulæ quinque cum filamentis alternantes; & glandulæ nectariferæ baſi ſtaminum inſidentes. Fructus pentacoccus, roſtratus; roſtris ſpiralibus introrſum barbatis.

cicutarium. 1. E. pedunculis multifloris, foliis pinnatis, foliolis multifidis ſeſſilibus. *L' Herit. ger. n.* 13.

Geranium caule procumbente, petiolis multifloris, foliis duplicato-pinnatis, pinnis acute inciſis. *Hall. hiſt. n.* 944.

Geranium cicutarium. *Sp. pl. p.* 951. *Fl. dan. t.* 986. *Curtis lond. Cavan. diſſ. p.* 226. *t.* 93. *f.* 1.

In aggeribus, campis, ruderatis, cultis, & alibi. Fl. Aprili, Majo.

ciconium. 2. E. pedunculis multifloris biflorisve, foliis interrupte bipinnatifidis laciniatis, caule ſtrigoſo. *L' Herit. ger. n.* 16.

Geranium petiolis multifloris, foliis pinnatis, pinnis latescentibus, dentatis, alterne maximis. *Hall. hiſt. n.* 946.

Geranium ciconium. *Sp. pl. p.* 952. *Iacqu. hort. tom.* 1. *p.* 7. *t.* 18.

In herbidis Litoralis, Iſtriæ. Fl. Majo, Iunio.

DECANDRIA.

GERANIUM. Calyx pentaphyllus. Corolla pentapetala, regularis. Glandulæ quinque melliferæ, baſi longiorum filamentorum adnatæ. Fructus pentacoccus, roſtratus: roſtris ſimplicibus, nudis.

macrorrhizum. 1. G. pedunculis bifloris, calycibus inflatis, piſtillo longiſſimo. *Sp. pl. p.* 953. *Wulfen in Iacqu. collect. tom.* 1. *p.* 258. *ic. t.* 134. *Cavan. diſſ. p.* 212. *t.* 85. *f.* 1.

In montanis rupeſtribus Forijulii, Croatiæ, Carnioliæ calidioris, Litoralis. Fl. ab Aprili ad Iunium. Per.

phæum. 2. G. pedunculis bifloris foliiſque alternis, calycibus ſubariſtatis, caule erecto, petalis undulatis. *Sp. pl. p.* 953. *Fl. dan. t.* 987. *Cavan. diſſ. p.* 210. *t.* 89. *f.* 2.

Geranium foliis rugoſis hirſutis, ſemiſeptilobis, lobis dentatis, petalis ſerratis acuminatis. *Hall. hiſt. n.* 934.

Geranium primum, pullo flore. *Clus. hiſt. p.* XCIX.

In alpinis, & ſubalpinis. Fl. æſtate. Per.

nodoſum. 3. G. pedunculis bifloris, foliis caulinis trilobis integris ſerratis: ſummis ſubſeſſilibus. *Sp. pl. p.* 953. *Cavan. diſſ. p.* 208. *t.* 80. *f.* 1.

Geranium quintum, ſextum Plateau. *Clus. hiſt. p.* CI.

In Carnioliæ calidioris, Croatiæ, Litoralis, Forijulii ſylvaticis aſperis. Fl. æſtate. Per.

ſylvaticum. 4. G. pedunculis bifloris, foliis ſubpeltatis quinquelobis inciſo-ſerratis, caule erecto, petalis emarginatis. *Sp. pl. p.* 954. *Fl. dan. t.* 124.

Geranium caule erecto, foliis rugoſis ſeptilobis, lobis acute ſerratis, petalis emarginatis. *Hall. hiſt. n.* 932.

Geranium ſecundum batrachioides minus. *Cluſ. hiſt. p.* XCIX.

In Auſtriæ, Styriæ alpinis, & ſubalpinis. Fl. Iulio, Auguſto. Per.

paluſtre. 5. G. pedunculis bifloris longiſſimis declinatis, foliis quinquelobis inciſis, petalis integris. *Sp. pl. p.* 954. *Cavan. diſſ. p.* 211. *t.* 87. *f.* 2.

Geranium caule diffuſo, foliis rugoſis ſemiquinquelobis, lobis trifidis, floribus ſparſis, petalis integerrimis. *Hall. hiſt. n.* 933.

Ad ſepes humidas, & in aliis udis locis Carnioliæ, Carinthiæ, Styriæ, Tyrolis. Fl. æſtate. Per.

pratenſe. 6. G. pedunculis bifloris, foliis ſubpeltatis multipartitis pinnato-laciniatis rugoſis acutis, petalis integris. *Sp. pl. p.* 954. *Curtis lond. Cavan. diſſ. p.* 210. *t.* 87. *f.* 1.

Geranium caule erecto, foliis rugoſis, hirſutis, multilobis, lobis trifidis, lobulis ſemipinnatis, floribus umbellatis. *Hall. hiſt. n.* 931.

Geranium tertium batrachioides majus. *Cluſ. hiſt. p.* C.

In pratenſibus. Fl. æſtate. Per.

argenteum. 7. G. pedunculis bifloris, foliis ſubpeltatis ſeptempartitis trifidis tomentoſo-ſericeis, petalis emarginatis. *Sp. pl. p.* 954. *Wulfen in Iacqu. collect. tom.* 2. *p.* 143. *ic. t.* 546. *Cavan. diſſ. p.* 205. *t.* 77. *f.* 3.

Geranium alpinum longius radicatum. *Pon. in Cluſ. hiſt. p.* CCCXLII. *Segu. ver.* I. *p.* 471. *t.* 10.

In Carniolia in alpibus wochinenſibus in ſummis jugis montis *Liſez*, *Cserna Perſt.* Fl. Iulio, Auguſto. Per.

robertianum. 8. G. foliis quinatis ternatisque inciſis, calycibus decemangulatis. *L' Herit. ger. n.* 42. *Sp. pl. p.* 955. *Curtis lond. Fl. dan. t.* 694. *Cavan. diſſ. p.* 215. *t.* 86. *f.* 1.

Geranium foliis duplicato-pinnatis, pinnis ultimis confluentibus, calycibus ſtriatis, hirſutis. *Hall. hiſt. n.* 943.

L' Herbe à Robert. Regn. bot.

In cultis, ſylvaticis, ad ſepes. Fl. æſtate.

lucidum. 9. G. pedunculis bifloris, calycibus pyramidatis angulatis elevato-rugoſis, foliis quinquelobis rotundatis. *Sp. pl. p.* 955. *Fl. dan. t.* 218. *Cavan. diſſ. p.* 214. *t.* 80. *f.* 2.

Geranium foliis ſubrotundis, ſemiquinquelobis, lobis obtuſis, calycibus transverſim rugoſis. *Hall. hiſt. n.* 942.

In rupeſtribus umbroſis montanis, & ſubalpinis. Fl. Majo, Iunio.

molle. 10. G. pedunculis bifloris, foliisque floralibus alternis, petalis bifidis, calycibus muticis, caule erectiusculo. *Syſt. veget. p.* 617. *Curtis lond. Fl. dan. t.* 679. *Cavan. diſſ. p.* 203. *t.* 83. *f.* 3.

Geranium foliis molliſſimis, hirſutis, reniformibus, ſemiquinquefidis, lobis ſemitripartitis obtuſis. *Hall. hiſt. n.* 939.

In cultis, ruderatis, ad vias. Fl. Majo, Iunio.

columbinum. 11. G. pedunculis folio longioribus, foliis quinquepartito-multifidis, calycibus pentagonis, arillis glabris. *L' Herit. ger. n.* 35. *Sp. pl. p.* 956. *Cavan. diſſ. p.* 200. *t.* 82. *f.* 1.

Geranium caule procumbente, foliis quinquepartitis, lobis trilobis, lobulis trifidis. *Hall. hiſt. n.* 939.

In cultis, paſcuis, ad ſepes, ceſpites. Fl. Iunio, Iulio.

diſſectum. 12. G. pedunculis folio brevioribus, foliis quinquepartitis trifidis, petalis emarginatis, arillis villoſis. *L' Herit. ger. n.* 35. *Sp. pl. p.* 956. *Fl. dan. t.* 936. *Curtis lond. Retzii obſ. faſc.* 1. *p.* 22. *Cavan. diſſ. p.* 199. *t.* 78. *f.* 2.

Geranium caule erecto, foliis quinquelobis, lobis trifidis acutis. *Hall. hiſt. n.* 937.

In Tyroli circa Oenopontum, copioſiſſime in *Giggelberg*; circa Halam ad Oenum, alibique. Fl. æſtate.

rotundifolium. 13. G. petalis ſubintegris calyci ſubariſtato æqualibus, foliis reniformibus lobatis inciſis: ſinubus glanduloſis. *L' Herit. ger. n.* 33. *Sp. pl. p.* 957. *Cavan. diſſ. p.* 214. *t.* 93. *f.* 2.

Geranium viscidum, caule decumbente, foliis ſemiquinquelobis, lobis tridentatis & quinquelobatis. *Hall. hiſt. n.* 941.

In cultis, ruderatis, ad vias. Fl. Iunio, Iulio.

ſanguineum. 14. G. pedunculis unifloris, foliis quinquepartitis trifidis orbiculatis. *Sp. pl. p.* 958. *Cavan. diſſ. p.* 195. *t.* 76. *f.* 1.

Geranium pedunculis unifloris, foliis ſeptilobis, petalis bifidis. *Hall. hiſt. n.* 930.

Geranium ſeptimum hæmatodes. *Clus. hiſt. p.* CII.

Le Bec de Grue ſanguin. Bulliard herb. t. 12.

In ſylvaticis, pratis ſicciorbus, ad agrorum margines. Fl. æſtate. Per.

POLYANDRIA.

ALTHÆA. Calyx duplex: exterior novemfidus. Arilli plurimi, monoſpermi.

officinalis. 1. A. foliis ſimplicibus tomentoſis. *Sp. pl. p.* 966. *Fl. dan. t.* 530. *Scop. carn. tom.* 2. *p.* 41.

Althæa tomentoſa, herbacea, caule erecto, foliis cordato-lanceolatis obſolete trifidis. *Hall. hiſt. n.* 1074.

Althæa vulgaris. *Clus. hiſt. p. XXIV.*

La Guimauve. Regn. bot.

In pratis humidis, ad ſepes humidas. Fl. æſtate. Per.

cannabi-na. 2. A. foliis inferioribus palmatis, ſuperioribus digitatis. *Sp. pl. p.* 966. *Iacqu. auſtr. t.* 101. *Scop. carn. tom.* 2. *p.* 42.

Alcea fruticoſa cannabino folio. *Cluſ. hiſt. p. XXV.*

Ad ſepes Auſtriæ, Pannoniæ, Croatiæ, Litoralis, Forijulii. Fl. a Iulio ad Septembrim. Per.

hirſuta. 3. A. foliis trifidis piloſo-hispidis, ſupra glabris, pedunculis ſolitariis unifloris. *Syſt. veget. p.* 624. *Iacqu. auſtr. t.* 170. *Scop. carn. tom.* 2. *p.* 42.

Althæa foliis radicalibus reniformibus, caulinis palmatis, trilobis & quinquelobis, calycibus hispidis. *Hall. hiſt. n.* 1073.

In aſperis Auſtriæ, Pannoniæ, Forijulii, Litoralis. Fl. æſtate.

MALVA. Calyx duplex: exterior triphyllus. Arilli plurimi, monoſpermi.

rotundi-folia. 1. M. caule proſtrato, foliis cordato-orbiculatis obſolete quinquelobis, pedunculis fructiferis declinatis. *Sp. pl. p.* 969. *Curtis lond. Scop. carn. tom.* 2. *p.* 43. *Hall. hiſt. n.* 1070.

Malva ſylveſtris, folio ſubrotundo. *Bauh. pin.* 314.

La Mauve a feuilles rondes. Bulliard herb. t. 161.

In cultis, ruderatis, ad vias. Fl. æſtate.

ſylveſtris. 2. M. caule erecto herbaceo, foliis ſeptemlobatis acutis, pedunculis petiolisque piloſis. *Sp. pl. p.* 969. *Curtis lond. Fl. dan. t.* 721. *Scop. carn. tom.* 2. *p.* 43.

Malva caule erecto, foliis lobatis, lobis ſerratis, quinis & ſeptenis. *Hall. hiſt. n.* 1069.

Malva ſylveſtris, folio ſinuato. *Bauh. pin.* 314.

Habitat & floret cum præc.

Alcea. 3. M. caule erecto pilis faſciculatis ſcabro, foliis multipartitis ſcabriuſculis. *Retzii ſcand. p.* 163. *Sp. pl. p.* 971. *Scop. carn. tom.* 2. *p.* 43. *Mœnch. meth. p.* 611.

Malva caule erecto, foliis ſcabris, trilobatis, lobis lateralibus bipartitis, medio tripartito & quinquepartito. *Hall. hiſt. n.* 1071.

Ad vias, radices montium, lapidum acervos, agrorum margines Styriæ, Carinthiæ, Carnioliæ. Fl. æſtate. Per.

Calycinis foliolis ovatis; annulo ad baſin calycis protuberante diverſa a M. moſchata, cui foliola calycis lanceolata, aut linearia; annullus nullus. Curtis lond. ſub Malva moſchata.

LAVATERA. Calyx duplex: exterior trifidus. Arilli plurimi, monoſpermi.

thuringiaca. 1. L. caule herbaceo, fructibus denudatis, calycibus inciſis. *Sp. pl. p.* 973. *Iacqu. auſtr. t.* 311.

Althæa pannonica ſecunda. *Clus. hiſt. p. XXIII.*

In Auſtria, Pannonia ad ſylvarum margines, ad ſepes, vineas, alibique. Fl. Iulio, Auguſto. Per.

HIBISCUS. Calyx duplex: exterior polyphyllus. Capſula quinquelocularis, polyſperma.

Pentacarpos. 1. H. foliis inferioribus cordatis angulatis, ſuperioribus ſubhaſtatis, floribus ſubnutantibus, piſtillo cernuo. *Sp. pl. p.* 981. *Iacqu. collect. tom.* 1. *p.* 150. *ic. t.* 143.

Ketmia paluſtris minor, folio anguſto, flore parvo purpuraſcente, fructu depreſſo pentagono. *Zann. iſt. p.* 155. *t.* 91.

In Carniæ udis, *nella valle del Chiadin.* Fl. Iulio Per.

Trionum. 2. H. foliis tripartitis inciſis, calycibus inflatis. *Sp. pl. p.* 981. *Curtis magaz.* 209.

Alcea veſicaria. *Bauh. pin.* 317.

Ad vias, in cultis. Fl. æſtate.

DIADELPHIA.

HEXANDRIA.

FUMARIA. Calyx diphyllus, minimus. Corolla ringens. Filamenta duo, membranacea. Antheræ tres in ſingulo filamento, terminales.

bulboſa. 1. F. caule ſimplici, bracteis longitudine florum. *Sp. pl. p.* 933. *Fl. dan. t.* 605. *Scop. carn. tom.* 2. *p.* 46.

Radix cava major. *Cluſ. hiſt. p.* 271.

La Fumeterre bulbeuſe. Regn. bot.

In ſylvaticis. Fl. Aprili. Per.

Variat flore albo, rubro, flaveſcente.

capnoides. 2. F. ſiliquis linearibus tetragonis, caulibus diffuſis acutangulis. *Sp. pl. p.* 984. *Scop. carn. tom.* 2. *p.* 47.

Fumaria foliis pinnatis, pinnis pinnatis, pinnulis lobatis, ovatis, ſiliquis brevibus incriſpatis. *Hall. hiſt. n.* 347.

Pſeudo-fumaria flore luteo. *Riv. tetr. t.* 74.

In rupeſtribus Litoralis, copioſiſſime in agro fluminenſi *ai molini.* Fl. a primo vere ad hyemem. Per.

officinalis. 3. F. pericarpiis monoſpermis racemoſis, caule diffuſo. *Sp. pl. p.* 984. *Curtis lond. Fl. dan. t.* 940. *Scop. carn. tom.* 2. *p.* 47.

Fumaria foliis multifidis, ſubrotunde lanceolatis, fructibus monoſpermis. *Hall. hiſt. n.* 346.

La Fumeterre. Regn. bot.

In cultis, ad vias. Fl. æſtate.

capreolata. 4. F. pericarpiis monoſpermis racemoſis, foliis ſcandentibus ſubcirrhoſis. *Sp. pl. p.* 985. *Fl. dan. t.* 340.

Fumaria viticulis & capreolis plantis vicinis adhærens. *Bauh. pin.* 143.

In Iſtria veneta prope portum *Lemme. I. Hoſt.* Fl. æſtate.

OCTANDRIA.

POLYGALA. Calyx pentaphyllus: foliolis duobus alæformibus, coloratis. Legumen obcordatum, biloculare.

amara. 1. P. floribus criſtatis racemoſis, caulibus erectiusculis, foliis radicalibus obovatis majoribus. *Sp. pl. p.* 987. *Iacqu. auſtr. t.* 412.

Polygala procumbens, foliis imis ſubrotundis, ſupremis linearibus. *Hall. hiſt. n.* 343.

In Auſtriæ pascuis, montanis ſylvaticis. Fl. primo vere. Per.

vulgaris. 2. P. floribus criſtatis racemoſis, caulibus herbaceis ſimplicibus procumbentibus, foliis lineari-lanceolatis. *Sp. pl. p.* 986. *Fl. dan. t.* 516. *Scop. carn. tom.* 2. *p.* 48.

Le Laitier commun. Bulliard. herb. t. 177.

In pratis, collibus, alibique. Fl. Majo, Iunio. Per.

Variat floris colore cœruleo, albo, purpureo.

major. 3. P. floribus criſtatis racemoſis, caulibus herbaceis ſimplicibus ſuberectis, foliis lineari-lanceolatis. *Syſt. veget. p.* 638. *Iacqu. auſtr. t.* 413.

Polygala vulgaris major prima. *Clus. hiſt. p.* 324.

In collibus Auſtriæ, Pannoniæ. Fl. Majo, Iunio. Per.

Chamæbuxus. 4. P. floribus imberbibus ſparſis: carinæ apice ſubrotundo, caule fruticoſo, foliis lanceolatis. *Sp. pl. p.* 989. *Iacqu. auſtr. t.* 233. *Scop. carn. tom.* 2. *p.* 49.

Polygaloides procumbens, foliis duris, ovatis, nervo ariſtato. *Hall. hiſt n.* 345.

Anonymos flore colutex. *Clus. hiſt. p.* 105.

In collibus, montoſis ſylvaticis. Fl. primo vere.

Flos ochroleucus. Dantur tamen individua, in quibus rubro-purpureus eſt. In aliis vero bina calycis foliola, alarum inſtar ſurſum expanſa, rubent, tertium autem, in tubulum convolutum, flaveſcit, ejusque extrema pars aurea eſt. Iacqu. l. c.

DECANDRIA.

SPARTIUM. Calyx deorſum productus. Filamenta germini adhærentia. Stigma longitudinale, ſupra villoſum.

junceum. 1. S. ramis oppoſitis teretibus apice floriferis, foliis ſimplicibus lanceolatis. *Sp. pl. p.* 995. *Curtis magaz.* 85.

Geniſta juncea. *Scop. carn. tom.* 2. *p.* 50.

Spartium Dioscoridis, Geniſta Plinii. *Clus. hiſt. p.* 102.

Le Genet d' Espagne. Regn. bot.

In aſperis agri tergeſtini, fluminenſis. Fl. Majo, Iunio. Fr.

decumbens. 2. S. caule decumbente ramoſo, foliis ſimplicibus ovato-lanceolatis, floribus longe pedunculatis. *Hall. hiſt. n.* 355. *Ait. Kew. tom.* 3. *p.* 11. *Iacqu. ic. t.* 555.

Geniſta pedunculata. *L' Herit. ſtirp. nov. t.* 89.

Geniſta humifuſa. *Wulfen in Iacqu. collect. tom.* 2. *p.* 169.

Le Genet de Haller. Reynier mem. pour l' hiſt. nat. de la Suiſſe, 1. *p.* 211.

In apricis declivibus agri tergeſtini circa *Hundsberg*. *Wulfen*. Ego in Styria in declivibus montis *Gamowitz*. Fl. Majo, Iunio. Fr.

Caules plerique decumbentes, pauci erecti. Pedunculi, qui primo breves, dein elongantur.

ſcoparium. 3. S. foliis ſimplicibus ternatisque, ramis acutangulis, floribus axillaribus ſolitariis, leguminibus margine villoſis. *Ait. Kew. tom.* 3. *p.* 12. *Sp. pl. p.* 996. *Curtis lond. Fl. dan. t.* 313.

Spartium foliis inferioribus ternatis, hirſutis, ſuperioribus ſimplicibus. *Hall. hiſt. n.* 354.

In Iſtria veneta verſu *Lonche*. *I. Hoſt*. Fl. Majo, Iunio. Fr.

radiatum. 4. S. foliis ternatis linearibus ſeſſilibus, petiolis perſiſtentibus, ramis oppoſitis angulatis. *Sp. pl. p.* 996.

Geniſta radiata. *Scop. carn. tom.* 2. *p.* 51. *Miller ic.* 173. *t.* 259. *f.* 1.

Geniſta radiata ſeu ſtellaris. *Bauh. hiſt.* 1. *p.* 399.

In Carniolia in alpinis, ſubalpinis, & in alpium convallibus. Fl. Iunio, Iulio. Fr.

GENISTA. Calyx bilabiatus. Vexillum oblongum a piſtillo, ſtaminibus carinam deprimentibus, deorſum reflexum.

ſagittalis. 1. G. ramis ancipitibus membranaceis articulatis, foliis ovato-lanceolatis. *Syſt. veget. p.* 645. *Jacqu. auſtr. t.* 209. *Scop. carn. tom.* 2. *p.* 51.

Geniſta foliis adpreſſis, ellipticis, ſine lanceolato patente. *Hall. hiſt. n.* 353.

In Auſtria in collibus herbidis circa *Fridau*. In ſimilibus locis, & in montanis, ſubalpinis Styriæ, Carinthiæ, Carnioliæ, Croatiæ, Pannoniæ. Fl. a Majo ad Iulium. Per.

tinctoria. 2. G. foliis lanceolatis glabris, ramis ſtriatis teretibus erectis. *Sp. pl. p.* 998. *Fl. dan. t.* 526. *Scop. carn. tom.* 2. *p.* 52.

Geniſta inermis, foliis glabris, confertis, ellipticis, lanceolatis, floribus ſeſſilibus, ſpicatis. *Hall. hiſt. n.* 350.

Geniſta tinctoria vulgaris. *Clus. hiſt. p.* 101.

In ſylvaticis. Fl. æſtate. Per.

ſericea. 3. G. foliis lanceolatis ſubtus villoſis, ramis teretibus ſtriatis erectis, racemis paucifloris ſecundis, bracteis calyce longioribus. *Wulfen in Iacqu. collect. tom.* 2. *p.* 167. *ic. t.* 556.

In Litorali inter Tergeſtum & Obzinam, *Wulfen*. Ego inter *Moſchienizze* & *Fianona*. Fl. Iunio. Fr.

piloſa. 4. G. foliis lanceolatis obtuſis, caule tuberculato decumbente. *Sp. pl. p.* 999. *Iacqu. auſtr. t.* 208. *Scop. carn. tom.* 2. *p.* 52.

Chamægeniſta prima. *Clus. hiſt. p.* 103.

In pratis ſiccioribus, collibus apricis, montanis herbidis, aſperis. Fl. Aprili, Majo. Fr.

germanica. 5. G. ſpinis compoſitis, ramis floriferis inermibus, foliis lanceolatis. *Sp. pl. p.* 999. *Scop. carn. tom.* 2. *p.* 54.

Geniſta foliis ovato-lanceolatis, hirſutis, ſpinis ramoſis, in ramis ſeneſcentibus. *Hall. hiſt. n.* 352.

In ſylvaticis, pratis montanis. Fl. Majo, Iunio. Fr.

hiſpanica. 6. G. ſpinis decompoſitis, ramis floriferis inermibus, foliis linearibus piloſis. *Sp. pl. p.* 999. *Wulfen in Iacqu. collect. tom.* 2. *p.* 165. *ic. t.* 557.

Geniſta ſylveſtris. *Scop. carn. tom.* 2. *p.* 53.

Geniſtella montis ventoſi ſpinoſa. *Bauh. hiſt.* I. *p.* 400.

Geniſtella monſpeliaca ſpinoſa. *Bauh. prodr.* 157.

In herbidis, montanis Litoralis, Croatiæ, Carnioliæ calidioris. Fl. Majo, Iunio. Fr.

ONONIS. Calyx quiqnepartitus: laciniis linearibus. Vexillum ſtriatum. Legumen turgidum, ſeſſile. Filamenta connata absque fiſſura.

ſpinoſa. 1. O. floribus axillaribus geminatis, foliis ternatis: ſuperioribus ſolitariis, ramisque villoſis. *Ait. Kew. tom.* 3. *p.* 21. *Retzii ſcand. p.* 169. *obs. fasc.* 2. *p.* 68. *Fl. dan. t.* 783. *Syſt. veget. p.* 651.

Ononis floribus ſubſeſſilibus lateralibus, caùle ſpinoſo. *Hort. cliff.* 359.

Anonis caule decumbente, ramis ſenescentibus ſpinoſis. *Hall. hiſt. n.* 356.

Anonis ſpinoſa. *Scop. carn. tom.* 2. *p.* 54.

La Bugrane des Champs. Bulliard. herb. t. 105.

In pratis ſiccioribus, campis, montanis ſylvaticis. Fl. Iunio, Iulio. Per.

hircina. 2. O. floribus ſubſpicatis geminatis, foliis inferioribus ternatis; ſuperioribus ſolitariis villoſiusculis, ramis piloſo-villoſis. *Ait. Kew. tom.* 3. *p.* 21. *Iacqu. hort. tom.* 1. *p.* 40. *t.* 93. *Syſt. veget. p.* 651.

Ononis arvenſis. *Syſt. veget. p.* 651. *Retzii ſcand. p.* 169.

Ononis ſpinoſa var. mitis. *Sp. pl. p.* 1006.

Ononis fœtens. *Allion. pedem. tom.* 1. *p.* 317. *t.* 41. *f.* 1.

Ononis altiſſima. *De Lamarck. encycl.* 1. *p.* 506.

Anonis mitior prima purpureo flore. *Clus. hiſt. p.* 99.

Anonis non ſpinoſa, flore thyrſoide carneo, polonica. *Ban. ic.* 1214.

Ononis mitior flore purpureo. *Besl. eyſt. æſtiv.* 10. *t.* 2. *f.* 2.

In Tyrolis, Auſtriæ, Styriæ, Pannoniæ pratis, convallibus alpinis. Fl. æſtate. Per.

Tota planta viſcido, & hircinum virus olente humore obducta eſt. Clus. l. c.

repens. 3. O. caulibus diffuſis, ramis erectis: foliis ſuperioribus ſolitariis, ſtipulis ovatis. *Sp. pl. p.* 1006.

Anonis maritima procumbens, foliis hirſutie pubescentibus. *Pluk. alm.* 33. *Dill. elth.* 29. *t.* 25. *f.* 28.

In Tyroli ad pedes montium inter *Sterzing* & *Mittewalde.* Fl. æſtate. Per.

minutiſſima. 4. O. floribus ſubſeſſilibus lateralibus, foliis ternatis glabris, ſtipulis enſiformibus, calycibus ariſtis corolla longioribus. *Sp. pl. p.* 1007. *Iacqu. auſtr. t.* 240.

Anonis foliis ternatis, ovatis, ſerratis, hirſutis; floribus ſeſſilibus, calycibus hiantibus. *Hall. hiſt. n.* 359.

In Auſtria in montium declivibus apricis inter *Baaden* & *Gumpoldskirchen.* Fl. æſtate. Per.

pinguis. 5. O. pedunculis unifloris ariſtatis, foliis ternatis lanceolatis: ſtipulis integerrimis. *Sp. pl. p.* 1009.

Anonis lutea, non ſpinoſa, Dalechampii. *Bauh. hiſt.* 2. *p.* 393. *Zann. iſt. p.* 18. *t.* 275.

In Carnia *nella valle del Chiadin. I. Hoſt.* Per.

Natrix. 6. O. pedunculis unifloris ariſtatis, foliis ternatis viſcoſis, ſtipulis integerrimis. *Sp. pl. p.* 1008.

Anonis viſcida, foliis petiolatis ternatis, petiolis unifloris. *Hall. hiſt. n.* 358.

In Tyroli locis riguis, ad rivulos inter *Sterzing* & *Mittewalde.* Fl. æſtate. Per.

rotundifolia. 7. O. fruticoſa, foliis ternatis ovatis, dentatis, calycibus triphyllo-bracteatis, pedunculis ſubtrifloris. *Syſt. veget. p.* 653. *Iacqu. auſtr. app. t.* 49.

Anonis foliis ſubrotundis, ſerratis, ternatis, petiolis multifloris. *Hall. hiſt. n.* 357.

In Tyroli in ſubalpinis leontinis. Fl. a Majo ad Iulium. Fr.

ANTHYLLIS. Calyx ventricoſus. Legumen ſubrotundum, tectum.

Vulneraria. 1. A. herbacea, foliis pinnatis inæqualibus, capitulo duplicato. *Sp. pl. p.* 1012. *Fl. dan. t.* 988.

Vulneraria foliis radicalibus ovatis, caulinis pinnatis, umbellis transverſis. *Hall. hiſt. n.* 398.

Vulneraria Anthyllis. *Scop. carn. tom.* 2. *p.* 55.

In pratis, collibus, alibique. Fl. æſtate. Per.

montana. 2. A. herbacea, foliis pinnatis æqualibus, capitulo terminali ſecundo, floribus obliquatis. *Sp. pl. p.* 1012. *Iacqu. auſtr. t.* 334.

Vulneraria caule procumbente, foliis pinnatis ſericeis, ovato-lanceolatis, umbellis globoſis. *Hall. hiſt. n.* 397.

Vulneraria montana. *Scop. carn. tom.* 2. *p.* 56.

In Auſtriæ, Carnioliæ, Croatiæ, Litoralis montanis. Fl. Majo, Iunio. Per.

OROBUS. Calyx baſi obtuſus, ore quinquedentatus: dentibus tribus inferioribus acutioribus; duobus ſuperioribus brevioribus, profundius & obtuſe diviſis. Stigma lineare, latere interiore a medietate ad apicem ſtyli pubescens.

luteus. 1. O. foliis pinnatis: foliolis ovato-oblongis, ſtipulis rotundato-lunatis dentatis, caule ſimplici. *Sp. pl. p.* 1028.

Orobus caule ramoſo, erecto, foliis ovato-lanceolatis. *Hall. hiſt. n.* 419.

Orobus montanus. *Scop. carn. tom.* 2. *p.* 60. *t.* 41.

In Carnioliæ, Croatiæ ſubalpinis, & alpinis. Fl. Iunio, Iulio. Per.

vernus. 2. O. foliis pinnatis: foliolis ovatis, ſtipulis ſemiſagittatis integerrimis, caule ſimplici. *Sp. pl. p.* 1028. *Scop. carn. tom.* 2. *p.* 58.

Orobus caule ſimpliciſſimo, foliis ſenis, ovatis, acutis. *Hall. hiſt. n.* 416.

Orobus pannonicus primus. *Cluſ. hiſt. p.* CCXXX.

L' Orobe. Regn. bot.

In ſylvis. Fl. primo vere. Per.

tuberoſus. 3. O. foliis pinnatis: foliolis lanceolatis, ſtipulis ſemiſagittatis integerrimis, caule ſimplici. *Sp. pl. p.* 1028. *Fl. dan. t.* 781. *Curtis lond. Scop. carn. tom.* 2. *p.* 58.

Orobus caule ſimplici, foliis ſenis, ellipticis, radice tuberoſa. *Hall. hiſt. n.* 417.

In Litoralis, Carnioliæ ſylvaticis. Fl. Majo, Iunio. Per.

albus. 4. O. foliis bijugis: foliolis enſiformibus petiolalis, ſtipulis ſimplicibus, caule ſimplici, *Syſt. veget. p.* 661.

Orobus pannonicus. *Iacqu. auſtr. t.* 39.

Orobus auſtriacus. *Crantz faſc. p.* 374. *t.* 1. *f.* 1.

In Auſtriæ, Pannoniæ collibus, pratis humidis. Fl. Majo, Iunio. Per.

niger. 5. O. caule ramoſo, foliis ſexjugis: foliolis ovato-oblongis. *Sp. pl. p.* 1028. *Scop. carn. tom.* 2. *p.* 59.

Orobus caule ramoſo, foliis ovatis duodenis. *Hall. hiſt. n.* 418.

Orobus pannonicus ſecundus. *Clus. hiſt. p.* CCXXX.

In ſylvis. Fl. Majo, Iunio. Per.

LATHYRUS. Calyx bilabiatus, ſuperne brevior. Stylus planus, ſupra villoſus, ſuperne latior.

Aphaca. 1. L. pedunculis unifloris, cirrhis aphyllis, ſtipulis ſagittato-cordatis. *Sp. pl. p.* 1029. *Scop. carn. tom.* 2. *p.* 61. *Curtis lond.*

Lathyrus aphyllos, ſtipulis ſagittatis latiſſimis. *Hall. hiſt. n.* 442.

In Litorali, Carniolia inter ſegetes, in vineis. Fl. Majo, Iunio.

Niſſolia. 2. L. pedunculis unifloris, foliis ſimplicibus, ſtipulis ſubulatis. *Sp. pl. p.* 1029. *Wulfen in Iacqu. collect. tom.* 4. *p.* 261.

Lathyrus pedunculis unifloris, foliis gramineis, ſtipulis minimis, *Hall. hiſt. n.* 441.

Niſſolia vulgaris. *Tourn. inſt.* 656. *Lindern. hort. p.* 73. *t.* 3.

Lathyrus anguſtifolius erectus, folio ſingulari, ſine capreolis, Niſſolii. *Magn. hort.* 112.

In agro goritienſi. *Wulfen.* Fl. Iunio.

ſetifolius. 3. L. pedunculis unifloris, cirrhis diphyllis, foliolis ſetaceo-linearibus. *Sp. pl. p.* 1031.

Lathyrus anguſtiſſimo folio, ſive capilaceo. *Bauh. prodr.* 148.

In vineis, ſaxoſis agri fluminenſis. Fl. a Majo ad Iulium.

angulatus. 4. L. pedunculis unifloris ariſtatis, cirrhis diphyllis ſimpliciſſimis, foliolis linearibus. *Sp. pl. p.* 1031. *Scop. carn. tom.* 2. *p.* 62.

Lathyrus foliis gramineis, petiolis unifloris ſetigeris, ſiliquis longis erectis. *Hall. hiſt. n.* 440.

Lathyrus folio tenuiore, floribus rubris. *Buxb. cent.* 3. *p.* 23. *t.* 42. *f.* 2.

In aſperis Litoralis, Forijulii. Fl. Iunio, Iulio.

tuberoſus. 5. L. pedunculis multifloris, cirrhis diphyllis: foliolis ovalibus, internodiis nudis. *Sp. pl. p.* 1033. *Curtis lond.*

Lathyrus ſcapis multifloris, foliis ovatis, capreolis trifidis. *Hall. hiſt. n.* 435.

In pratis, cultis, ad vias, ſepes Auſtriæ, Pannoniæ. Fl. æſtate. Per.

pratenſis. 6. L. pedunculis multifloris, cirrhis diphyllis ſimpliciſſimis: foliolis lanceolatis. *Sp. pl. p.* 1033. *Fl. dan. t.* 527. *Curtis lond. Scop. carn. tom.* 2. *p.* 64.

Lathyrus ſcapis multifloris, foliis lanceolatis, capreolis ſimplicibus. *Hall. hiſt. n.* 436.

Lathyrus luteus ſylveſtris dumetorum. *Bauh. hiſt.* 2. *p.* 304.

In pratis, ad ſepes, & alibi. Fl. Majo, Iunio, Iulio. Per.

ſylveſtris. 7. L. pedunculis multifloris, cirrhis diphyllis: foliolis enſiformibus, internodiis membranaceis. *Sp. pl. p.* 1032. *Fl. dan. t.* 325.

Lathyrus caule alato, foliis geminis enſiformibus. *Hall. hiſt. n.* 434.

Lathyrus ſylveſtris Dodonæi. *Clus. hiſt. p.* CCXXIX.

In Auſtria, Pannonia, Litorali ad ſepes. Fl. æſtate. Per.

latifolius. 8. L. pedunculis multifloris, cirrhis diphyllis: foliolis lanceolatis, internodiis membranaceis. *Sp. pl. p.* 1033. *Miller illuſtr. ic.*

Lathyrus caule alato, foliis ovato-lanceolatis, rigidis, ſcapis multifloris. *Hall. hiſt. n.* 433.

In ſylvaticis, ad ſepes. Fl. æſtate. Per.

paluſtris. 9. L. pedunculis multifloris, cirrhis polyphyllis, ſtipulis lanceolatis. *Sp. pl. p.* 1034. *Fl. dan. t.* 399.

Lathyrus foliis ſenis, floribus racemoſis. *Hall. hiſt. n.* 431.

In Auſtria in paluſtribus prope *Himberg.* Fl. Majo, Iunio. Per.

Cirrhi hexaphylli, internodia alata, foliola lanceolata, ſtipulæ ſemiſagittatæ, pedunculi ſubtriſtori, corollæ cæruleæ. Retzii obs. faſc. I. p. 24.

VICIA. Stigma latere inferiore transverſe barbatum.

* *Pedunculis elongatis.*

piſiformis. I. V. pedunculis multifloris, petiolis polyphyllis, foliolis ovatis: infimis ſeſſilibus. *Sp. pl. p.* 1034. *Iacqu. auſtr. t.* 364.

Vicia foliis ovatis, maximis, ſiliquis racemoſis pendulis. *Hall. hiſt. n.* 428.

Piſum ſylveſtre. *Clus. hiſt. p.* CCXXIX.

In Auſtriæ, Pannoniæ ſylvaticis. Fl. Iunio, Iulio. Per.

dumetorum. 2. V. pedunculis multifloris, foliolis reflexis ovatis mucronatis, ſtipulis ſubdentatis. *Sp. pl. p.* 1035.

Vicia foliis ovatis, oblongis, ſtipulis ſerratis, ſiliquis racemoſis, pendulis. *Hall. hiſt. n.* 427.

Cracca ſylvatica. *Riv. tetr. t.* 50.

Vicia maxima dumetorum. *Bauh. pin.* 345.

In Auſtriæ Pannoniæ ſylvis. Fl. æſtate. Per.

ſylvatica. 3. V. pedunculis multifloris, foliolis ovalibus, ſtipulis denticulatis. *Sp. pl. p.* 1035. *Fl. dan. t.* 277. *Scop. carn. tom.* 2. *p.* 66.

Vicia foliis ovatis, ſtipulis argute dentatis, ſiliquis racemoſis pendulis. *Hall. hiſt. n.* 426. *t.* 12.

Viciæ pulchrum genus multifolium, Galega quibusdam. *Bauh. hiſt.* 3. *p.* 309.

Vicia multiflora maxima perennis, tetro odore, floribus albentibus: lineis cœruleis notatis. *Pluk. alm.* 387. *t.* 71. *f.* 1.

In montanis Auſtriæ prope *Annaberg. Iacqu.* In Carnioliæ ſimilibus locis, & in aſcenſu alpium. *Scop.* Fl. a Iunio ad Auguſtum. Per.

caſſubica. 4. V. pedunculis ſubſexfloris, foliolis denis ovatis acutis, ſtipulis integris. *Sp. pl. p.* 1035. *Retzii ſcand. p.* 171. *obs. faſc.* 1. *p.* 82. *Fl. dan. t.* 98.

Vicia Gerardi. *Iacqu. auſtr. t.* 229. *Syſt. veget. p.* 664.

Vicia militans. *Crantz faſc. p.* 388. *t.* 1. *f.* 2.

In Auſtriæ, Pannoniæ pratenſibus, ſylvaticis. Fl. Iunio, Iulio. Per.

Cracca. 5. V. pedunculis multifloris, floribus imbricatis, foliolis lanceolatis pubescentibus, ſtipulis integris. *Sp. pl. p.* 1035. *Curtis lond. Fl. dan. t.* 804. *Scop. carn. tom.* 2. *p.* 66.

Vicia foliis linearibus ſericeis, racemis multifloris reflexis, ſtipulis integerrimis. *Hall. hiſt. n.* 424.

Vicia ſylveſtris ſecunda. *Cluſ. hiſt. p.* CCXXXV.

In ſylvaticis, pratis, ad ſepes, inter ſegetes. Fl. Iunio, Iulio. Per.

Vel ſola radice perenni diverſa a ſequente.

villoſa. 6. V. pedunculis multifloris, floribus imbricatis, foliolis ovalibus villoſis, ſtipulis nervoſis baſi dentatis. *Roth. germ. tom.* 2. *p.* 182.

In Auſtria, Pannonia inter ſegetes, in pratis, vineis. Fl. Iunio, Iulio.

oroboides. 7. V. pedunculis multifloris, caule erecto, petiolis abſque cirrhis. *Wulfen in Iacqu. collect. tom.* 4. *p.* 323.

Orobus pannonicus quartus. *Cluſ. hiſt. p.* CCXXXI.

In montanis, ſubalpinis Croatiæ, Carnioliæ. Fl. Iunio, Iulio. Per.

** *Floribus axillaribus, ſubſeſſilibus.*

ſativa. 8. V. leguminibus ſeſſilibus ſubbinatis erectis, foliolis retuſis, ſtipulis notatis. *Sp. pl. p.* 1037. *Fl. dan. t.* 522. *Scop. carn. tom.* 2. *p.* 64.

Vicia foliis imis ovatis, ſuperioribus linearibus, ſcapis breviſſimis, bifloris. *Hall. hiſt. n.* 430.

Vicia ſylveſtris. *Cluſ. hiſt. p.* CCXXXV.

Inter ſegetes. Fl. æſtate.

Lathyroides. 9. V. leguminibus ſeſſilibus ſolitariis erectis glabris, foliolis ſenis: inferioribus obcordatis. *Sp. pl. p.* 1037. *Fl. dan. t.* 58. *Iacqu. miſcell. tom.* 2. *p.* 299. *t.* 18.

In Auſtriæ, Pannoniæ herbidis, præcipue in inſulis danubialibus. Fl. Aprili, Majo.

Dantur plantæ, in quibus petioli, præſertim ſuperiores, in cirrhum elongantur; dantur quoque individua, in quibus omnes petioli cirrhis deſtituuntur.

hybrida. 10. V. leguminibus ſeſſilibus ſolitariis reflexis piloſis pentaſpermis, corollæ vexillis villoſis. *Sp. pl. p.* 1037. *Iacqu. hort. tom.* 2. *p.* 68. *t.* 146.

In cultis, & incultis Litoralis. Fl. Majo, Iunio.

Caules angulati, ſtriati, ramoſi; pedales. Foliola oblonga vel obverſe ovata, retuſa cum exigua ſetula terminante, piloſa. Stipulæ dorſo nulla macula tinctæ: inferiores ad baſin extrorſum producuntur, ibidem haſtatæ aut dentatæ; ſuperiores integerrimæ. Flores ſolitarii. Legumen pendulum vel patens. Iacqu. l. c.

pannonica. 11. V. pedunculis ſubtrifloris, corollæ vexillis villoſis, leguminibus piloſis, ſtipulis integerrimis notatis. *Iacqu. auſtr. t.* 34.

Vicia ſylveſtris albo flore. *Clus. hiſt. p.* CCXXXV.

In Auſtriæ, Pannoniæ arvis, pratis. Fl. Iunio, Iulio.

Caules bipedales, angulati, pubeſcentes aut glabriuſculi, ſi ſuſtentacula in vicinia, ut caules præcedentis, ſcandentes, ſi nulla, humi procumbentes. Foliola foliolis V. hybridæ ſimilia, niſi quod longiora & minus retuſa ſint. Stipulæ integerrimæ baſi macula fuſca notatæ. Pedunculi biaut triflori: rari quadrifloti; rariſſimi uniflori. Legumina oblonga, piloſa. Iacqu. l. c.

ſepium. 12. V. leguminibus pedicellatis ſubquaternis erectis, foliolis ovatis integerrimis: exterioribus decreſcentibus. *Sp. pl. p.* 1038. *Fl. dan. t.* 699. *Scop. carn. tom.* 2. *p.* 65.

Vicia foliis ovatis oblongis, ſcapis brevibus quadrifloris. *Hall. hiſt. n.* 429.

Vicia ſepium perennis. *Bauh. hiſt.* 2. *p.* 313.

In ſylvaticis, ad ſepes. Fl. Majo, Iunio. Per.

ſerratifolia. 13. V. caule erecto, petiolis abſque cirrho, foliolis ſerratis. *Syſt. veget. p.* 665. *Iacqu. auſtr. app. t.* 8.

In humidis Pannoniæ circa Sopronium, & ad lacum Peiſonis. Fl. Iunio.

ERVUM. Calyx quinquepartitus, longitudine corollæ.

Lens. 1. E. pedunculis ſubbifloris, ſeminibus compreſſis convexis. *Sp. pl. p.* 1039. *Hall. hiſt. n.* 421. *Scop. carn. tom.* 2. *p.* 67.

Inter ſegetes, ad pagos. Fl. æſtate.

tetraſpermum. 2. E. pedunculis ſubbifloris, ſeminibus globoſis quaternis. *Sp. pl. p.* 1039. *Curtis lond. Fl. dan. t.* 95. *Scop. carn. tom.* 2. *p.* 68.

Vicia foliis linearibus, ſiliquis gemellis glabris. *Hall. hiſt. n.* 423.

Viciæ ſeu Craccæ minimæ ſpecies cum ſiliquis glabris. *Bauh. hiſt.* 2. *p.* 315.

In herbidis, cultis. Fl. æſtate.

hirſutum. 3. E. pedunculis multifloris, ſeminibus globoſis binis. *Sp. pl. p.* 1039. *Curtis lond. Fl. dan. t.* 639. *Scop. carn. tom.* 2. *p.* 67.

Vicia foliis linearibus, ſiliquis racemoſis, diſpermis, hirſutis. *Hall. hiſt. n.* 422.

In agris. Fl. æſtate.

CICER. Calyx quinquepartitus, longitudine corollæ: laciniis quatuor ſuperioribus vexillo incumbentibus. Legumen rhombeum, turgidum, diſpermum.

arietinum. 1. C. foliolis ſerratis. *Sp. pl. p.* 1040. *Hall. hiſt. n.* 399.

Cicer ſativum. *Bauh. pin.* 347.

Le Pois chiche. Regn. bot.

In Iſtria, Tyroli inter ſegetes. Fl. æſtate.

CYTISUS. Calyx bilabiatus. Legumen baſi attenuatum.

Laburnum. 1. C. racemis ſimplicibus pendulis, foliolis ovato-oblongis. *Sp. pl. p.* 1041. *Iacqu. auſtr. t.* 306. *Scop. carn. tom.* 2. *p.* 68. *Hall. hiſt. n.* 360.

In Pannoniæ, Croatiæ, Carnioliæ montanis, ſubalpinis. Fl. Majo, Iunio. Fr.

nigricans. 2. C. racemis ſimplicibus erectis, foliolis ovato-oblongis. *Sp. pl. p.* 1041. *Iacqu. auſtr. t.* 387. *Scop. carn. tom.* 2. *p.* 69.

Cytiſus foliis ovatis nitidis, floribus ſpicatis, cernuis, calycibus & ſiliquis ſericeis. *Hall. hiſt. n.* 361.

Cytiſus quartus. *Clus. hiſt. p.* 95.

In ſylvaticis, collibus aſperis. Fl. Iunio, Iulio. Fr.

capitatus. 3. C. floribus umbellatis terminalibus, ramis erectis, foliolis ovatis. *Syſt. veget. p.* 666. *Iacqu. auſtr. t.* 33.

Cytiſus hirſutus. *Crantz faſc. p.* 398.

Cytiſi quinti ſpecies altera. *Clus. hiſt. p.* 96.

Cytiſi ſeptimi ſpecies altera. *Clus. hiſt. p.* 97.

In montoſis ſylvaticis. Fr.

Iam Aprili, Majo oriuntur ex axillis foliorum flores ſolitarii, modo bini, quandoque terni. Iunio ex ipſa ramorum ſummitate exoritur florum breviter pedunculatorum numero incerto capitulum, ſive umbella.

auſtriacus. 4. C. floribus umbellatis terminalibus, caulibus erectis, foliolis lanceolatis. *Sp. pl. p.* 1042. *Iacqu. auſtr. t.* 21.

Cytiſus quintus. *Clus. hiſt. p.* 95.

In aſperis, herbidis, ſylvaticis Auſtriæ, Pannoniæ. Fl. a Iunio ad Auguſtum. Fr.

ſupinus. 5. C. floribus pedunculatis, foliolis villoſis, caulibus decumbentibus fruteſcentibus. *Syſt. veget. p.* 667. *Iacqu. auſtr. t.* 20.

Cytiſus ſeptimus. *Cluſ. hiſt. p.* 96.

In collibus apricis, montoſis herbidis, aſperis. Fl. Aprili, Majo. Fr.

purpureus. 6. C. floribus pedunculatis ſolitariis, foliis glabris, caulibus decumbentibus fruteſcentibus, leguminibus ſuperne falcatis. *Syſt. veget. p.* 667. *Iacqu. auſtr. app. t.* 48. *Scop. carn. tom.* 2. *p.* 69. *t.* 43. *Zann. iſt. p.* 83. *t.* 2. 10.

In Carnioliæ, Croatiæ, Forijulii apricis aſperis, herbidis. Fl. Majo, Iunio. Fr.

COLUTEA. Calyx quinquefidus. Legumen inflatum, baſi ſuperiore dehiſcens.

arboreſcens. 1. C. arborea, foliolis obcordatis. *Sp. pl. p.* 1045. *Curtis magaz.* 81. *Scop. carn. tom.* 2. *p.* 57.

Colutea foliis ovatis, emarginatis, floribus racemoſis. *Hall. hiſt. n.* 400.

Le Baguenaudier. Regn. bot.

In collibus, ſylvaticis Litoralis, Iſtriæ, Forijulii, Carnioliæ calidioris, Croatiæ. Fl. æſtate.

GLYCYRRHIZA. Calyx bilabiatus. Legumen ovatum compreſſum.

echinata. 1. G. leguminibus echinatis, foliis ſtipulatis, foliolo impari ſeſſili. *Syſt. veget. p.* 669. *Iacqu. hort. tom.* 1. *p.* 41. *t.* 95.

Glycyrrhiza capite echinato. *Bauh. pin.* 352.

In Croatia inter *Novogrod* & *Gradiska. Hacquet.* Fl. Iunio, Iulio. Per.

CORONILLA. Calyx bilabiatus: dentibus duobus ſuperioribus connatis. Vexillum vix alis longius. Legumen iſthmis interceptum.

Emerus. 1. C. fruticoſa, pedunculis ſubtrifloris, corollarum unguibus calyce triplo longioribus, caule angulato. *Sp. pl. p.* 1046. *Scop. carn. tom.* 2. *p.* 71.

Coronilla frutescens, foliis retuſis, ſeptenis, ſiliquis longiſſimis, obiter articulatis. *Hall. hiſt. n.* 389.

Coronilla major. *Riv. tetr. t.* 95.

Colutea ſcorpioides prima elatior, & humilior. *Clus. hiſt. p.* 97.

In collibus, montoſis ſylvaticis. Fl. a Martio ad Iunium.

minima. 2. C. ſuffruticoſa procumbens, foliolis novenis ovatis, ſtipula oppoſitifolia emarginata, leguminibus angulatis nodoſis. *Syſt. veget. p.* 670. *Iacqu. auſtr. t.* 271.

Coronilla caulibus lignoſis, procumbentibus, foliis ovatis, floribus umbellatis, ſiliquis alatis. *Hall. hiſt. n.* 390.

In collibus herbidis, montanis, ſubalpinis. Fl. Majo, Iunio. Per.

coronata. 3. C. herbacea, foliolis novenis obovatis: internis cauli approximatis, ſtipula oppoſitifolia bipartita. *Sp. pl. p.* 1047. *Iacqu. auſtr. t.* 95.

Coronilla caule erecto, foliis undenis, ovatis, floribus umbellatis, ſiliquis articulatis pendulis. *Hall. hiſt. n.* 388.

Coronilla montana. *Riv. tetr. t.* 207. *Scop. carn. tom.* 2. *p.* 72.

Colutea ſcorpioides altera, *Clus. hiſt. p.* 98.

In collibus, montoſis. Fl. Iunio. Per.

varia. 4. C. herbacea, leguminibus erectis teretibus toroſis numeroſis, foliolis plurimis glabris. *Sp. pl. p.* 1048. *Scop. carn. tom. 2. p.* 73.

Coronilla caule brachiato, angulato, foliis vigenis, ariſtatis, floribus umbellatis. *Hall. hiſt. n.* 387.

Coronilla flore vario. *Riv. tetr. t.* 94.

Securidaca vera. *Clus. hiſt. p.* CCXXXVI.

In cultis, pratis, ad vias. Fl. æſtate. Per.

ORNITHOPUS. Legumen articulatum, teres, arcuatum.

ſcorpioides. 1. O. foliis ternatis ſubſeſſilibus: impari maximo. *Sp. pl. p.* 1049.

Ornithopodium ſcorpioides. *Scop. carn. tom.* 2. *p.* 73.

Ornithopodium portulacæ folio. *Tourn. inſt.* 400.

In olivetis, vineis, inter ſegetes Litoralis. Fl. Majo, Iunio.

HIPPOCREPIS. Legumen compreſſum, altera ſutura pluries emarginatum, curvum.

uniſiliquoſa. 1. H. leguminibus ſeſſilibus ſolitariis rectis. *Syſt. veget. p.* 670.

Ferrum equinum ſiliquis ſolitariis lunatim exciſſis. *Hall. hiſt. n.* 392.

Ferrum equinum ſiliqua ſingulari. *Gar. aix. t.* 114.

Tergeſti *alle Saule. I. Hoſt.* Fl. Majo, Iunio.

comoſa. 2. H. leguminibus pedunculatis confertis arcuatis: margine exteriore repandis. *Syſt. veget. p.* 671. *Iacqu. auſtr. t.* 431. *Scop. carn. tom. 2. p.* 74.

Ferrum equinum ſiliquis umbellatis, undulatis. *Hall. hiſt. n.* 391.

In collibus apricis, montoſis herbidis, alpinis Auſtriæ, Carnioliæ. In Litorali in ſimilibus locis, & in cultis. Fl. a Majo ad Iulium. Per.

SCORPIURUS. Legumen iſthmis interceptum, revolutum, teres.

ſubvilloſa. 1. S. pedunculis ſubquadrifloris, leguminibus extrorſum ſpinis conſertis acutis. *Sp. pl. p.* 1050.

In apricis herbidis, cultis agri fluminenſis, Iſtriæ. Fl. Majo.

HEDYSARUM. Corollæ carina transverſe obtuſa. Legumen articulis monoſpermis.

obſcurum. 1. H. foliis pinnatis, ſtipulis vaginalibus, caule erecto flexuoſo, floribus pendulis. *Syſt. veget. p.* 676. *Iacqu. auſtr. t.* 168.

Hedyſarum caule erecto, ramoſo, foliis ovatis, ſiliquis pendulis læviſſimis. *Hall. hiſt. n.* 395. *t.* 12.

In alpibus. Fl. Iulio, Auguſto. Per.

Onobrychis. 2. H. foliis pinnatis, leguminibus monoſpermis aculeatis, corollarum alis calycem æquantibus, caule elongato. *Syſt. veget. p.* 676. *Iacqu. auſtr. t.* 352.

Onobrychis caule erecto, ramoſo, floribus ſpicatis. *Hall. hiſt. n.* 396.

Onobrychis viciæfolia. *Scop. carn. tom.* 2. *p.* 76.

Onobrychis Dodonæi. *Clus. hiſt. p.* CCXXXII.

Le Sainfoin. Regn. bot.

In pratis. Fl. a Majo ad Iulium. Per.

Caput galli. 3. H. foliis pinnatis, leguminibus monoſpermis: criſtæ dentibus ſubulatis, alis breviſſimis, caule diffuſo. *Syſt. veget. p.* 676.

Onobrychis, fructu echinato, minor. *Bauh. pin.* 350. *prodr.* 149.

In Carniæ herbidis prope *Comeglians. I. Hoſt.* Fl. Iunio, Iulio.

GALEGA. Calyx dentibus ſubulatis, ſubæqualibus. Legumen ſtriis obliquis, ſeminibus interjectis.

officinalis. 1. G. leguminibus ſtrictis erectis, foliolis lanceolatis ſtriatis nudis. *Sp. pl. p.* 1062.

Galega foliis ellipticis, ſpinis & ſiliquis acutis. *Hall. hiſt. n.* 394.

Galega vulgaris. *Cluſ. hiſt. p.* CCXXXIII.

Le Galega. Regn. bot.

In udis. Fl. æſtate. Per.

PHACA. Legumen ſemibiloculare.

alpina. 1. P. caulescens erecta glabra, leguminibus oblongis inflatis ſubpiloſis. *Sp. pl. p.* 1064. *Wulfen in Iacqu. miscell. tom.* 2. *p.* 93. *ic. t.* 151.

Aſtragalus caule erecto, ramoſiſſimo, foliis ellipticis, hirſutis, ſiliquis veſicariis, pendulis. *Hall. hiſt. n.* 401.

In Tyroli in agro leontino in alpibus *Marienwald*, & in arenoſis Iſolæ. *Wulfen.* Fl. Iulio. Per.

frigida. 2. P. caule erecto, ſtipulis maximis, foliolis ovatis, novenis, leguminibus inflatis. *Hall. hiſt. n.* 402. ſub Aſtragalo. *Iacqu. auſtr. t.* 166.

In ſummis alpium auſtriacarum, ſtyriacarum jugis. Fl. Iulio, Auguſto. Per.

auſtralis. 3. P. caule ramoſo proſtrato, foliolis lanceolatis, florum alis ſemibifidis. *Syſt. veget. p.* 680. *Wulfen in Iacqu. miscell. tom.* 2. *p.* 43. *t.* 3.

Aſtragalus caule ramoſo, proſtrato, foliis lanceolatis, florum alis emarginatis. *Hall. hiſt. n.* 403.

In Carinthia in alpe *Trog* prope Pontebam, in alpibus reichenavienſibus ad S. Laurentium *hinter den zween Speichkogeln über dem Petauerſee* in ſic dicto *Garten* ſub altiſſima alpe *Prett*; in Tyroli cum P. alpina. *Wulfen.* Fl. Iunio, Iulio. Per.

ASTRAGALUS. Legumen biloculare, gibbum.

piloſus. 1. A. caulescens erectus piloſus, floribus ſpicatis, leguminibus ſubulatis piloſis. *Sp. pl. p.* 1065. *Iacqu. auſtr. t.* 51.

Aſtragalus caule erecto, ramoſo, ex alis ſpicifero, ſiliquis teretibus. *Hall. hiſt., n.* 411.

Onobrychis tertia. *Clus. hiſt. p.* CCXXXIX.

In collibus herbidis, pratis ſiccioribus, montoſis ſylvaticis. Fl. Iunio, Iulio. Per.

ſulcatus. 2. A. caulescens erectus, caule glabro ſulcato, foliolis lanceolatis, leguminibus ſubtriquetris. *Sp. pl. p.* 1065. *Iacqu. hort. tom.* 3. *p.* 23. *t.* 40.

In Pannonia in paluſtribus ad lacum Peiſonis. Fl. Iunio, Iulio. Per.

aſper. 3. A. caulescens ſtrictus ſubaſper, floribus longe ſpicatis erectis, leguminibus oblongis. *Syſt. veget. p.* 681. *Iacqu. miscell. tom.* 2. *p.* 335. *ic. t.* 33.

Cum præcedente. *Lübeck.* Fl. Majo, Iunio. Per.

Cicer. 4. A. caulescens proſtratus, leguminibus ſubgloboſis inflatis mucronatis piloſis. *Sp. pl. p.* 1067. *Iacqu. auſtr. t.* 251.

Aſtragalus caule ramoſo, diffuſo, ſpicis erectis, ſiliquis inflatis, rotundis, unguiculatis. *Hall. hiſt. n.* 409.

In pratis, & aliis herbidis locis. Fl. Iulio, Auguſto. Per.

glycyphyllos. 5. A. caulescens proſtratus, leguminibus ſubtriquetris arcuatis, foliolis ovalibus pedunculo longioribus. *Sp. pl. p.* 1067. *Scop. carn. tom.* 2. *p.* 76.

Aſtragalus caule proſtrato, foliis ovatis, ſcapis brevibus racemoſis, ſiliquis ſubtriquetris incurvis. *Hall. hiſt. n.* 413.

Glaux vulgaris. Glycyrrhiza ſylveſtris Gesneri. *Clus. hiſt. p.* CCXXXIII.

In pratenſibus, ſylvaticis. Fl. æſtate. Per.

Onobrychis. 6. A. caulescens diffuſus, pedunculis ſpicatis, vexillis flore duplo longioribus. *Sp. pl. p.* 1070. *Iacqu. auſtr. t.* 38. *Scop. carn. tom.* 2. *p.* 77.

Aſtragalus caule ſubereſto, ramoſo, foliis linearibus, ſericeis, vexillis longiſſimis. *Hall. hiſt. n.* 412.

Ad agrorum margines, aggeres, in pratis ſiccioribus, collibus. Fl. æſtate. Per.

leontinus. 7. A. caulescens proſtratus, leguminibus ovatis villoſis, floribus ſpicatis ereſtis. *Syſt. veget. p.* 683. *Wulfen in Iacqu. miscell. tom.* 2. *p.* 59. *ic. t.* 154.

In Tyroli in agro leontino prope Iſolam. Fl. Iulio. Per.

auſtriacus. 8. A. caulescens procumbens, floribus racemoſis, leguminibus utrinque acutis piloſis, foliolis linearibus emarginatis. *Sp. pl. p.* 1070. *Iacqu. auſtr. t.* 195.

Onobrychis ſecunda. *Clus. hiſt. p.* CCXXXIX.

In Auſtriæ, Pannoniæ collibus, campis, pratis. Fl. æſtate. Per.

alpinus. 9. A. caulescens procumbens, floribus pendulis racemoſis, leguminibus utrinque acutis piloſis. *Sp. pl. p.* 1070.

Aſtragalus caule procumbente ramoſo, foliis ovatis, ſiliquis inflatis, pendulis, hirſutis. *Hall. hiſt. n.* 404.

Aſtragalus alpinus minimus. *Fl. lapp. n.* 267. *t.* 9. *f.* 1.

Aſtragalus alpinus, foliis viciæ, ramoſus & procumbens. *Fl. dan. t.* 51.

In alpibus tyrolenſibus. Fl. Iulio. Per.

monta-nus. 10. A. ſubacaulis, ſcapis folio longioribus, floribus laxe ſpicatis erectis, leguminibus ovatis acumine inflexo. *Sp. pl. p.* 1070. *Iacqu. auſtr. t.* 167. *Scop. carn. tom.* 2. *p.* 78. *t.* 45.

Aſtragalus foliis ovato-lanceolatis, ſericeis, ſiliquis erectis, inflatis, ſubhirſutis. *Hall. hiſt. n.* 408.

Onobrychis quarta. *Clus. hiſt. p.* CCXL.

In alpinis. Fl. Iunio, Iulio. Per.

uralen-ſis. 11. A. acaulis, ſcapo erecto foliis longiore, leguminibus ſubulatis inflatis villoſis erectis. *Sp. pl. p.* 1071. *Wulfen in Iacqu. miscell. tom.* 2. *p.* 150. *ic. t.* 155.

Aſtragalus ſcapis aphyllis, foliis ovato-lanceolatis, ſericeis, ſiliquis turgidis, hirſutis, erectis, ſtyliferis. *Hall. hiſt. n.* 410. *t.* 14.

In Tyroli in Iſola prope Leontium, viciniſque alpibus. Fl. a Majo ad Iulium. Per.

campe-ſtris. 12. A. acaulis, calycibus leguminibusque villoſis, foliolis lanceolatis acutis, ſcapo decumbente. *Sp. pl. p.* 1072.

Aſtragalus ſpicis aphyllis, foliis lanceolatis hirſutis, ſiliquis erectis, inflatis, ſemibilocularibus. *Hall. hiſt. n.* 406. *t.* 13.

Aſtragalus pyrenaicus, barbæ Iovis folio, non ramoſus, flore ochroleuco, glomerato. *Scheuch. it.* 4. *p.* 330.

In Carinthia in declivibus herbidis Taureri malnizenſis, alpis melnikenſis diſtrictus gmündtenſis *in der Malta*; item in Tyroli in alpibus leontinis marienwaldenſibus. *Wulfen.* Ego in Styria in ſummis jugis alpium judenburgenſium. Fl. a Iulio ad Septembrim. Per.

incanus. 13. A. acaulis, ſcapis declinatis, foliis tomentoſis, leguminibus ſubulatis ſubarcuatis ſubtomentoſis apice reflexo. *Sp. pl. p.* 1072.

Aſtragalus monſpeſſulanus. *Scop. carn. tom.* 2. *p.* 77. *t.* 45.

In aſperis agri tergeſtini. Fl. Iunio. Per.

exſcapus. 14. A. acaulis exſcapus, leguminibus lanatis, foliolis villoſis. *Syſt. veget. p.* 685. *Iacqu. collect. tom.* 2. *p.* 269. *ic. t.* 561.

Glaux montana acaulon. *Bauh. pin.* 347. *prodr.* 147.

In Pannonia. Fl. Majo, Iunio. Per.

TRIFOLIUM. Flores ſubcapitati. Legumen vix calyce longius, non dehiſcens, deciduum.

M. cærulea. 1. T. racemis ovatis, leguminibus ſeminudis mucronatis, caule erecto. *Sp. pl. p.* 1077.

Lotus hortenſis odorata. *Bauh. pin.* 331.

Le Lotier odorant. Regn. bot.

In Carniolia, Tyroli ad pagos. Fl. Iunio, Iulio.

Foliorum pulverem cum farina miſcent incolæ ad gratum pani odorem conciliandum.

M. officinalis. 2. T. leguminibus racemoſis nudis diſpermis rugoſis acutis, caule erecto. *Sp. pl. p.* 1078. *Fl. dan. t.* 934. *Scop. carn. tom.* 2. *p.* 85.

Melilotus ſiliquis racemoſis, pendulis, acutis, rugoſis, diſpermis. *Hall. hiſt. n.* 362.

Le Melilot. Regn. bot.

In campis, pratis, inter ſegetes, ad vias. Fl. æſtate.

hybridum. 3. T. capitulis umbellaribus, leguminibus tetraſpermis, caule adſcendente. *Sp. pl. p.* 1079.

Trifolium caule ſubereĉto, ſpicis depreſſis, ſiliquis tetraſpermis. *Hall. hiſt. n.* 368.

Trifoliaſtrum pratente corymbiferum. *Mich. gen. p.* 28. *t.* 25. *f.* 2. 6.

In cultis, herbidis, ſucculentis. Fl. æſtate. Per.

repens. 4. T. capitulis umbellaribus, leguminibus tetraſpermis, caule repente. *Sp. pl. p.* 1080. *Curtis lond. Fl. dan t.* 990. *Scop. carn. tom.* 2. *p.* 84.

Trifolium caule repente, ſpicis depreſſis, ſiliquis tetraſpermis. *Hall. hiſt. n.* 367.

In pratis, pascuis, cultis. Fl. æſtate. Per.

alpinum. 5. T. capitulis umbellaribus, ſcapo nudo, leguminibus diſpermis pendulis, foliolis lineari-lanceolatis. *Sp. pl. p.* 1080.

Trifolium ſcapis radicatis, floribus racemoſis, foliis lanceolatis nervoſis. *Hall. hiſt. n.* 369.

Trifolium anguſtifolium alpinum. *Pon. in Clus. hiſt. p.* CCCXL.

Trifolium alpinum, flore magno, radice dulci. *Bauh. pin.* 328. *prodr.* 143.

In alpibus tyrolenſibus *in Neuſtift im Gerichte Stubay.* Fl. Iulio, Auguſto. Per.

rubens. 6. T. ſpicis villoſis longis, caule ereĉto, foliolis ſerrulatis. *Sp. pl. p.* 1081. *Iacqu. auſtr. t.* 385. *Scop. carn. tom.* 2. *p.* 80.

Trifolium foliis nervoſis, ciliatis, ſpicis ovatis, obeſis, calycis denticulis plumoſis, imo longiſſimo. *Hall. hiſt. n.* 375.

Trifolii majoris tertii altera ſpecies, *Clus. hiſt. p.* CCXLVI.

Trifolium montanum, ſpica longiſſima rubente. *Bauh. pin.* 328.

In montanis ſylvaticis. Fl. Iunio, Iulio. Per.

pratenſe. 7. T. ſpicis denſis, corollis inæqualibus, dentibus calycinis quatuor æqualibus, ſtipulis ariſtatis, caulibus adſcendentibus. *Retzii ſcand. p.* 173. *Sibthorp oxon. p.* 228. *Sp. pl. p.* 1082. *Scop. carn. tom.* 2. *p.* 70. *Fl. dan. t.* 989.

Trifolium caule obliquo, foliis ovatis hirſutis, ſupremis conjugatis, vaginis ariſtatis. *Hall. hiſt. n.* 377.

Trifolium pratenſe purpureum. *Bauh. pin.* 327.

Le Treſle, ou le Triolet des Prés. Regn. bot.

In pratis, paſcuis, & alibi. Fl. æſtate. Per.

flexuoſum. 8. T. ſpicis laxis, corollis ſubæqualibus, ſtipulis ſubulatis conniventibus, caulibus flexuoſis ramoſis. *Retzii ſcand. p.* 174. *Iacqu. auſtr. t.* 386.

Trifolium medium. *Sibthorp oxon. p.* 229. *Huds. angl. ed.* 1. 284.

Trifolium alpeſtre. *Huds. angl. ed.* 2. 326.

Trifolium majus tertium. *Clus. hiſt. p.* CCXLV.

Ad margines ſylvarum, in pratis montanis. Fl. Iunio, Iulio. Per.

alpeſtre. 9. T. ſpicis denſis, corollis ſubæqualibus, ſtipulis ſetaceis divergentibus, foliolis lanceolatis, caulibus ſtrictis ſimpliciſſimis. *Retzii ſcand. p.* 174. *Iacqu. auſtr. t.* 433. *Syſt. veget. p.* 688. *Scop. carn. tom.* 2. *p.* 79.

Trifolium majus ſecundum. *Clus. hiſt. p.* CCXLV.

In montanis, collibus herbidis. Fl. Iunio. Per.

pannonicum. 10. T. ſpicis villoſis longis, foliolis integerrimis cauleque erecto villoſiſſimis. *Syſt. veget. p.* 688. *Iacqu. obs.* 2. *p.* 21. *t.* 42.

In Pannoniæ pratis. In Carniolia in herbidis alpium wochinenſium. Fl. æſtate. Per.

ochroleucum. 11. T. ſpicis villoſis, caule erecto pubeſcente, foliolis infimis obcordatis. *Syſt. veget. p.* 689. *Iacqu. auſtr. t.* 40.

In ſylvaticis. Fl. Iunio, Iulio.

anguſti-folium. 12. T. ſpicis villoſis conico-oblongis, dentibus calycinis ſetaceis ſubæqualibus, foliolis linearibus. *Sp. pl. p.* 1083. *Scop. carn. tom.* 2. *p.* 82.

Lagopus anguſtifolius hiſpanicus. *Clus. hiſt. p.* CCXLII.

In herbidis Litoralis, Forijulii. Fl. Iunio, Iulio.

arvenſe. 13. T. ſpicis villoſis ovalibus, dentibus calycinis ſetaceis æqualibus. *Sp. pl. p.* 1083. *Fl. dan. t.* 724. *Scop. carn. tom.* 2. *p.* 83.

Trifolium caule ramoſo, foliis lanceolatis ſerratis, ſpicis villoſis ovatis. *Hall. hiſt. n.* 373.

In cultis, ad pagos, vias. Fl. Iulio, Auguſto.

ſtellatum. 14. T. ſpicis piloſis ovatis, calycibus patentibus, caule diffuſo, foliolis obcordatis. *Sp. pl. p.* 1083. *Scop. carn. tom.* 2. *p.* 80.

Lagopus maximus Lobelii. *Clus. hiſt. p.* CCXLVI.

In cultis, & incultis Litoralis, Forijulii. Fl. Iunio, Iulio.

fragiferum. 15. T. capitulis ſubrotundis, calycibus inflatis bidentatis reflexis, caulibus repentibus. *Sp. pl. p.* 1086. *Curtis lond. Scop. carn. tom.* 2. *p.* 84.

Trifolium caule repente, ſpicis glabris, calycibus ſericeis, ampullaſcentibus. *Hall. hiſt. n.* 370.

In pratis humidis, alibique. Fl. æſtate. Per.

montanum. 16. T. ſpicis ſubimbricatis ſubtribus, vexillis ſubulatis emarceſcentibus, calycibus nudis, caule erecto. *Sp. pl. p.* 1087. *Scop. carn. tom.* 2. *p.* 83.

Trifolium cauliculo paucifloro, foliis ellipticolanceolatis, nervoſis, ſubtus ſubhirſutis. *Hall. hiſt. n.* 372.

In pratis montanis, collibus herbidis. Fl. æſtate. Per.

agrarium. 17. T. ſpicis ovalibus imbricatis, vexillis deflexis perſiſtentibus, calycibus nudis, caule erecto. *Sp. pl. p.* 1087. *Fl. dan. t.* 558.

Trifolium caule erecto, ſpicis ſtrepentibus pauciſloris. *Hall. hiſt. n.* 364.

In pratis. Fl. Iunio, Iulio.

ſpadiceum. 18. T. ſpicis ovalibus imbricatis, vexillis deflexis perſiſtentibus, calycibus piloſis, caule erecto. *Sp. pl. p.* 1087.

Trifolium caule erecto, ſpicis ſtrepentibus ovatis. *Hall. hiſt. n.* 365.

Trifolium montanum lupulinum. *Bauh. pin.* 328. *prodr.* 140.

In herbidis alpium ſtyriacarum, tyrolenſium. Fl. Iulio, Auguſto.

procumbens. 19. T. ſpicis ſubovalibus imbricatis, vexillis deflexis perſiſtentibus, caulibus diffuſis. *Sibthorp oxon. p.* 230. *Sp. pl. p.* 1088.

Trifolium caulibus diffuſis, ſpicis ovatis ſtrepentibus. *Hall. hiſt. n.* 363.

Trifolium agrarium. *Curtis lond.*

In pratis, paſcuis, inter ſegetes. Fl. a Iunio ad Auguſtum.

LOTUS. Legumen cylindricum, ſtrictum. Alæ ſurſum longitudinaliter conniventes. Calyx tubuloſus.

ſiliquoſus. 1. L. leguminibus ſolitariis membranaceo-quadrangulis, caulibus procumbentibus, foliis ſubtus pubeſcentibus. *Sp. pl. p.* 1089. *Iacqu. auſtr. t.* 361.

Lotus caule proſtrato, ſcapis unifloris, ſiliquis alatis. *Hall. hiſt. n.* 386.

Tetragonolobus Scandalida. *Scop. carn. tom.* 2. *p.* 87.

Lotus pratorum ſiliquoſus. *Clus. hiſt. p.* CCXLIV.

In pratis humidis, alibique. Fl. a Majo ad Iulium, Per.

cornicu-latus. 2. L. capitulis depreſſis, caulibus decumbentibus, leguminibus cylindricis patentibus. *Sp. pl. p.* 1092. *Fl. dan. t.* 991.

Lotus ſiliquis umbellatis rectiſſimis, cylindricis. *Hall. hiſt. n.* 385.

In pratis, collibus, montoſis apricis. Fl. æſtate. Per.

Variat foliis glabris, villoſis.

Dorycnium. 3. L. capitulis aphyllis, foliis ſeſſilibus quinatis. *Sp. pl. p.* 1093.

Dorycnium pentaphyllum. *Scop. carn. tom.* 2. *p.* 87.

Dorycnium hiſpanicum. *Clus. hiſt. p.* 100. *Hall. hiſt. n.* 284.

In collibus, montanis ſylvaticis. Fl. æſtate. Per.

MEDICAGO. Legumen compreſſum, cochleatum. Carina corollæ a vexillo deflectens.

ſativa. 1. M. pedunculis racemoſis, leguminibus contortis, caule erecto glabro. *Sp. pl. p.* 1096.

Medica caule recto, foliis oblongis, ſerratis, racemis erectis, ſiliquis planis, iterato-contortis. *Hall. hiſt. n.* 382.

Medica legitima. *Clus. hiſt. p.* CCXLII.

La Luzerne. Regn. bot.

In pratenſibus. Fl. æſtate. Per.

Occurrit in agro vindobonenſi varietas caulibus diffuſis, floribus ex flavo, viridi & purpureo variegatis.

falcata. 2. M. pedunculis racemoſis, leguminibus lunatis, caule proſtrato. *Sp. pl. p.* 1096. *Fl. dan. t.* 233.

Medica foliis oblongis ſerratis, ſiliquis racemoſis falcatis. *Hall. hiſt. n.* 381.

Medica falcata. *Scop. carn. tom.* 2. *p.* 88.

Medica flavo flore. *Cluſ. hiſt. p.* CCXLIII.

In pratis, collibus, ad vias. Fl. æſtate. Per.

lupulina. 3. M. ſpicis ovalibus, leguminibus reniformibus monoſpermis, caulibus procumbentibus. *Sp. pl. p.* 1097. *Curtis lond.*

Medica caule diffuſo, capitulis hemisphæricis, ſiliquis reniformibus. *Hall. hiſt. n.* 380.

Medica lupulina. *Scop. carn. tom.* 2. *p.* 88.

In cultis, pratis, paſcuis. Fl. æſtate.

marina. 4. M. pedunculis racemoſis, leguminibus cochleatis ſpinoſis, caule procumbente tomentoſo. *Sp. pl. p.* 1097.

Trifolium cochleatum tomentoſum. *Bauh. pin.* 329. *Zann. iſt. p.* 181. *t.* 62.

Medica marina. *Cluſ. hiſt. p.* CCXLIII.

In arenoſis Litoralis. Fl. æſtate. Per.

carſtienſis. 5. M. caule erecto, floribus umbellatis, leguminibus cochleatis, ſetoſis. *Iacqu. collect. tom.* 1. *p.* 86. *ic. t.* 156.

In cultis, & incultis Litoralis. Fl. Majo, Iunio. Per.

proſtrata. 6. M. leguminibus cochleato-inermibus, foliis ternatis: foliolis cuneiformibus, apice dentatis, ſtipulis ſetaceis integerrimis, caule diffuſo. *Syſt. veget. p.* 694. *Iacqu. hort. tom.* 1. *p.* 39.

In herbidis, cultis agri fluminenſis. Fl. æſtate. Per.

POLYADELPHIA.

POLYANDRIA.

HYPERICUM. Calyx quinquepartitus. Petala quinque. Filamenta multa, in quinque phalanges baſi connata.

quadrangulare. 1. H. floribus trigynis, caule quadrato herbaceo. *Sp. pl. p.* 1104. *Curtis lond. Fl. dan. t.* 640. *Scop. carn. tom.* 2. *p.* 91.

Hypericum caule quadrangulo, foliis ovatis, perforatis, punctatis. *Hall. hiſt. n.* 1038.

In udis. Fl. æſtate. Per.

perforatum. 2. H. floribus trigynis, caule ancipiti, foliis obtuſis pellucido - punctatis. *Sp. pl. p.* 1105. *Curtis lond. Scop. carn. tom.* 2. *p.* 92.

Hypericum caule terete, alato, ramoſiſſimo, foliis ovatis perforatis. *Hall. hiſt. n.* 1037.

Le Mille - Pertuis. Regn. bot.

In pratis, montanis ſylvaticis, & alibi. Fl. æſtate. Per.

humifuſum. 3. H. floribus trigynis axillaribus ſolitariis, caulibus ancipitibus proſtratis filiformibus, foliis glabris. *Sp. pl. p.* 1105. *Curtis lond.*

Hypericum caule proſtrato, foliis ovatis, calycibus ſerratis punctatis. *Hall. hiſt. n.* 1039.

Hypericum ſupinum tertium minimum. *Clus. hiſt. p.* CLXXXI.

In herbidis: in Carniolia circa *Eck*, in Auſtria ſuperiore circa Lentiam. Fl. Iunio, Iulio. Per.

montanum. 4. H. floribus trigynis, calycibus ſerraturis glandulofis, caule tereti erecto glabro, foliis ovatis. *Sp. pl. p.* 1105. *Fl. dan. t.* 173.

Hypericum foliis ovatis, per oram punctatis, calycibus lanceolatis, ſerratis, globuligeris. *Hall. hiſt. n.* 1042.

In ſylvaticis. Fl. Iunio, Iulio. Per.

barbatum. 5. H. floribus trigynis, calycibus & petalis ciliatis iisdemque & foliis punctatis. *Syſt. veget. p.* 701. *Iacqu. auſtr. t.* 259.

In agri vindobonenſis pratis inter *Hädersdorf* & *Mauerbach. Iacqu.* Fl. Majo, Iunio. Per.

hirſutum. 6. H. floribus trigynis, calycibus ſerrato-glanduloſis, caule tereti erecto, foliis ovatis pubescentibus. *Sp. pl. p.* 1105. *Curtis lond. Fl. dan. t.* 802.

Hypericum foliis ovatis, hirſutis, calycibus lanceolatis, ſerratis, glanduligeris. *Hall. hiſt. n.* 1043.

In montanis ſylvaticis. Fl. Iunio, Iulio. Per.

dubium. 7. H. floribus trigynis, caule ſubquadrato herbaceo, foliis ovatis, calycibus obtuſiſſimis. *Leers herb. p.* 169.

In Styria in alpium judenburgenſium herbidis, in præcipitiis. Fl. Iulio, Auguſto. Per.

SYNGENESIA.

POLYGAMIA ÆQUALIS.

* *Semifloſculoſi: corollulis omnibus ligulatis.*

TRAGOPOGON. Receptaculum nudum. Calyx ſimplex. Pappus plumoſus, ſtipitatus.

pratenſe. 1. T. calycibus corollæ radium æquantibus, foliis integris ſtrictis. *Sp. pl. p.* 1109. *Fl. dan. t.* 906. *Scop. carn. tom.* 2. *p.* 94.

Tragopogon foliis gramineis, carinatis, amplexicaulibus, caudatis, ſubcriſpis, calyce flori æquali. *Hall. hiſt. n.* 8.

Le Salſifix des Prés. Bulliard. herb. t. 209.

In pratis. Fl. æſtate.

majus. 2. T. calycibus corollæ radio longioribus, foliis integris ſtrictis, pedunculis ſuperne incraſſatis, corollulis apice rotundatis. *Syſt. veget. p.* 710. *Iacqu. auſtr. t.* 29.

In agro vindobonenſi ad margines ſylvarum, ad vias, in pratis, cultis. Fl. Iunio, Iulio.

SCORZONERA. Receptaculum nudum. Calyx imbricatus, ſquamis margine ſcarioſis. Pappus plumoſus, ſtipitatus.

humilis. 1. S. caule ſubnudo unifloro, foliis lato-lanceolatis nervoſis planis. *Sp. pl. p.* 1112. *Iacqu. auſtr. t.* 36. *Scop. carn. tom.* 2. *p.* 95.

Scorzonera caule nudo unifloro, foliis lanceolatis. *Hall. hiſt. n.* 10.

Scorzonera humilis latifolia pannonica ſecunda. *Clus. hiſt. p. CXXXVIII.*

In collibus, pratis planitiei, montanis apricis. Fl. Aprili, Majo. Per.

parviflora. 2. S. caulibus ramoſis, foliis lineari-enſiformibus integris, radio corollæ breviſſimo. *Syſt. veget.* p. 710. *Iacqu. auſtr. t.* 305.

In Auſtria in pratis paluſtribus prope *Draſskirchen. Iacqu.* Ego in Pannonia ad litus lacus Peiſonis. Fl. æſtate. Per.

hiſpanica. 3. S. caule ramoſo, foliis amplexicaulibus integris ſerrulatis. *Sp. pl. p.* 1112.

Scorzonera major hiſpanica prima. *Clus. hiſt. p. CXXXVII.*

Scorzonera major pannonica prima. *Clus. hiſt. p. CXXXVIII.*

In Auſtriæ, Pannoniæ ſylvaticis. Fl. æſtate. Per.

graminifolia. 4. S. foliis lineari-enſiformibus integris carinatis. *Sp. pl. p.* 1112. *Iacqu. obs.* 4. *p.* 13. *t.* 100.

Scorzonera caule ramoſo, foliis linearibus, acuminatis, carinatis, calycibus acutis. *Gmel. ſib.* 2. *p.* 4.

Scorzonera purpurea. *Scop. carn. tom.* 2. *p.* 96. *t.* 46.

In herbidis alpium wochinenſium, tolmienſium. Fl. Iulio, Auguſto. Per.

Flos in noſtra planta non eſt luteus, ſed purpureus.

purpurea. 5. S. foliis lineari-ſubulatis integris canaliculato-triquetris, pedunculis cylindricis. *Syſt. veget. p.* 711. *Iacqu. auſtr. t.* 35.

Scorzonera major anguſtifolia cœruleo flore, pannonica quarta. *Clus. hiſt. p. CXXXIX.*

In Auſtriæ, Pannoniæ montoſis herbidis, collibus apricis. Fl. Majo, Iunio. Per.

anguſti-folia. 6. S. foliis ſubulatis integris, pedunculo incraſſato, caule ſimpliciſſimo baſi villoſo. *Iacqu. en. p.* 140. *Sp. pl. p.* 1113.

Scorzonera caule nudo unifloro, foliis linearibus. *Hall. hiſt n.* 11.

Scorzonera humilis anguſtifolia, pannonica tertia. *Clus. hiſt. p. CXXXVIII.*

In collibus, pratis Auſtriæ, Pannoniæ. Fl. cum præc. Per.

hirſuta. 7. S. foliis linearibus cauleque unifloro hirſutis. *Syſt. veget. p.* 711.

Tragopogon calycibus corolla brevioribus, foliis integris ſtrictis ciliatis, caule hirto, pedunculis ſuperne incraſſatis. *Ger. prov. p.* 234.

Tragopogon alter luteus. *Columm. ecphr.* 1. *p.* 234. *t.* 233.

In pratis, aliisque herbidis locis agri fluminenſis. Fl. cum præc. Per.

laciniata. 8. S. foliis linearibus dentatis acutis, caule erecto, ſquamis calycinis patulo-mucronatis. *Sp. pl. p.* 1114. *Iacqu. auſtr. t.* 356.

Scorzonera caule folioſo, ramoſo, foliis anguſtiſſimis, imis rariter ſemipinnatis, ſuperioribus linearibus, lanceolatis. *Hall. hiſt. n.* 12.

Tragopogon laciniatum luteum. *Bauh. pin.* 274.

In pratis, campis, ad vias Auſtriæ, Pannoniæ. Fl. æſtate. Per.

Folia radicalia primordialia integra, ſpathulata, longe petiolata: petiolis membranaceis, cætera ſinuata aut pinnatifida, laciniis utrinque tribus, vel quatuor lanceolato-linearibus; impari longiore. Folia caulina ſinuata aut integra. Caulis baſi ramoſus: ramis nudis. Calycis ſquamæ exteriores apice denticulo notatæ. Iacqu. l. c.

APARGIA. Receptaculum ſubnudum, alveolis marginatis. Pappus ſeſſilis plumoſus. Calyx imbricatus, erectus.

aurea. 1. A. foliis runcinatis, caule ſubunifolio, calyce hiſpido.

Taraxacum calycibus hirſutis, ſquamis rectis. *Hall. hiſt. n.* 56. *t.* 1.

Leontodon aureum. *Syſt. veget. p.* 715. *Iacqu. auſtr. t.* 297.

In alpinis, ſubalpinis. Fl. æſtate. Per.

haſtilis. 2. A. ſcapo calyceque lævi, foliis lenceolatis dentatis integerrimis glabris.

Picris caule unifloro, foliis glabris, ſemipinnatis, calyce lævi. *Hall. hiſt. n.* 26.

Leontodon haſtile. *Sp. pl. p.* 1123. *Iacqu. auſtr. t.* 164.

Apargia danubialis. *Scop. carn. tom.* 2. *p.* 114.

In pratis humidis. Fl. Iunio, Iulio. Per.

autumnalis. 3. A. caule ramoſo, pedunculis ſquamoſis, foliis lanceolatis dentatis integerrimis glabris.

Picris caule nudo, brachiato, foliis glabris ſemipinnatis. *Hall. hiſt. n.* 28.

Leontodon autumnale. *Sp. pl. p.* 1123. *Scop. carn. tom.* 2. *p.* 113.

In pratis, paſcuis. Fl. a Iulio ad Octobrim. Per.

Radix præmorſa. Folia radicalia lanceolata, dentata, raris pilis adſperſa. Pedunculi ſuperne incraſſati. Semina transverſim ſtriata.

alpina. 4. A. calyce toto erecto & ſcapo infra florem hiſpidis, foliis dentatis ſubpiloſis, ſetis ſimpliciſſimis.

Leontodon alpinum. *Syſt. veget. p.* 715. *Iacqu. auſtr. t.* 93.

In alpium herbidis, Fl, a Iulio ad Septembrim. Per.

hiſpida. 5. A. calyce toto erecto, foliis dentatis integerrimis hispidis: ſetis ſimplicibus furcatisque.

Picris caule nudo, unifloro, foliis aſperis, dentatis. *Hall. hiſt. n.* 25.

Leontodon hispidum. *Sp. pl. p.* 1124. *Scop. carn. tom.* 2. *p.* 111.

In pratis, paſcuis, & alibi. Fl. æſtate. Per.

Tota planta ſetis bifidis, trifidis, indiviſisque horret. Folia lanceolata, dentata. Semina ſtriata: ſuperne denticulata; inferne tuberculata.

hirta. 6. A. calyce toto erecto, foliis ſinuato-pinnatifidis aſperis: pilis ſimplicibus furcatisque.

Rhagadiolus foliis ſemipinnatis aſperrimis. *Hall. hiſt. n.* 7.

Leontodon hirtum. *Sp. pl. p.* 1123. *Scop. carn. tom.* 2. *p.* 112.

Hieracium parvum, hirtum, caule aphyllo, criſpum quando ſiccatum. *Bauh. hiſt.* 2. *p.* 1038.

In Carnioliæ alpinis, ſubalpinis. Fl. Iulio, Auguſto. Per.

Folia ſemipinnata: pinnis triangularibus, undique aſpera. Semina exteriora foliaceo brevi ornamento, interiora rigidis ſetis coronata. Haller l. c.

SONCHUS. Receptaculum nudum. Calyx imbricatus, ventricoſus. Pappus piloſus, ſeſſilis.

paluſtris. 1. S. pedunculis calycibusque hispidis ſubumbellatis, foliis runcinatis baſi ſagittatis. *Sp. pl. p.* 1116. *Curtis lond.*

Sonchus lævior auſtriacus quintus altiſſimus. *Clus. hiſt. p.* CXLVII.

In udis, paluſtribus Auſtriæ, Pannoniæ. Fl. Iulio, Auguſto. Per.

arvenſis. 2. S. pedunculis calycibusque hiſpidis ſubumbellatis, foliis runcinatis baſi cordatis. *Sp. pl. p.* 1116. *Curtis lond.*

Sonchus foliis amplexicaulibus, ſemipinnatis, ſerratis, calycibus hiſpidis. *Hall. hiſt. n.* 23.

Hieracium arvenſe. *Scop. carn. tom.* 2. *p.* 110.

In arvis, vineis, pratis, alibique. Fl. æſtate. Per.

oleraceus. 3. S. pedunculis tomentoſis, calycibus glabris. *Sp. pl. p.* 1116. *Curtis lond. Fl. dan. t.* 682. 843.

Hieracium oleraceum. *Scop. carn. tom.* 2. *p.* 110.

Le Laitron doux. Regn. bot.

In cultis, & alibi. Fl. æſtate.

Variat foliis integris, plus vel minus laciniatis; lævibus, ac aſperis.

alpinus. 4. S. pedunculis bracteisque longioribus hirſutis, floribus racemoſis, foliis runcinatis baſi ſagittalis. *Frœlich in Uſter. ann.* 1. *p.* 20. *Fl. dan. t.* 182. *Retzii ſcand. p.* 186.

Sonchus foliis amplexicaulibus, petiolatis, triangularibus, petiolo pinnato, floribus ſpicatis. *Hall. hiſt. n.* 20.

Hieracium cœruleum. *Scop. carn. tom.* 2. *p.* 111.

In alpinis, & ſubalpinis. Fl. æſtate. Per.

LACTUCA. Receptaculum nudum. Calyx imbricatus, cylindricus, margine membranaceo. Pappus ſimplex, ſtipitatus.

Scariola. 1. L. foliis verticalibus carina aculeatis. *Sp. pl. p.* 1119.

Lactuca foliis ſemipinnatis, nervo ſpinoſo. *Hall. hiſt. n.* 14.

Lactuca ſylveſtris, coſta ſpinoſa. *Bauh. pin.* 123. *Zann. iſt. p.* 157. *t.* 294.

La Laitue ſauvage. Regn. bot.

In cultis, ruderatis, ad vias. Fl. Iulio, Auguſto.

viroſa. 2. L. foliis horizontalibus carina aculeatis dentatis. *Sp. pl. p.* 1119.

Lactuca foliis ovatis, patulis, nervo ſpinoſo. *Hall. hiſt. n.* 15.

Lactuca ſylveſtris, odore viroſo. *Bauh. pin.* 123.

Lactuca agreſtis, Scariolæ hortenſis folio, odore opii vehementi ſoporifero, & viroſo. *Lob. adv.* 89.

In Litorali, Forojulio ad ſepes, in cultis. Fl. Iunio.

ſaligna. 3. L. foliis haſtato-linearibus ſeſſilibus carina aculeatis *Sp. pl. p.* 1119. *Iacqu. auſtr. t.* 250.

Lactuca foliis imis ſemipinnatis, ſuperioribus amplexicaulibus, hamatis, nervo ſpinoſo. *Hall. hiſt. n.*. 13.

In aſperis, ad vineas, agrorum margines Auſtriæ, Pannoniæ. Fl. Auguſto.

perennis. 4. L. foliis linearibus dentato-pinnatis: laciniis ſurſum dentatis. *Sp. pl. p.* 1120. *Scop. carn. tom.* 2. *p.* 102.

Lactuca foliis glabris, pinnatis, caule brachiato. *Hall. hiſt. n.* 16.

In montanis Carnioliæ, Croatiæ, Forijulii, Litoralis. Fl. æſtate.

CHONDRILLA. Receptaculum nudum. Pappus piloſus, ſtipitatus. Calyx calyculatus, multiflorus.

juncea. 1. C. foliis radicalibus runcinatis, caulinis linearibus integris, *Syſt. veget. p.* 713. *Iacqu. auſtr. t.* 427. *Hall. hiſt. n.* 17.

Chondrilla viminalibus virgis. *Cluſ. hiſt. p. CXLIII.*

In collibus, aggeribus, ad agrorum margines. Fl. æſtate.

PRENANTHES. Receptaculum nudum. Calyx calyculatus. Pappus ſimplex. Floſculi ſimplici ſerie.

viminea. 1. P. foliorum ramentis cauli adnatis. *Sp. pl. p.* 1120. *Iacqu. auſtr. t.* 9.

Chondrilla viminea viſcoſa monſpeliaca. *Bauh. pin.* 131. *prodr.* 68.

In collibus Auſtriæ aſperis. Fl. Iulio.

purpurea. 2. P. floſculis quinis, foliis lanceolatis denticulatis. *Sp. pl. p.* 1121. *Iacqu. auſtr. t.* 317. *Scop. carn. tom.* 2. *p.* 103.

Prenanthes foliis amplexicaulibus, petiolatis, lanceolatis, ſerratis. *Hall. hiſt. n.* 19.

Sonchus lævior pannonicus quartus, purpureo flore. *Cluſ. hiſt. p. CXLVII.*

In ſylvis montanis, & ſubalpinis. Fl. Iulio, Auguſto. Per.

muralis. 3. P. floſculis quinis, foliis runcinatis. *Sp. pl. p.* 1121. *Curtis lond. Fl. dan. t.* 509. *Scop. carn. tom.* 2. *p.* 103.

Prenanthes foliis ſerratis, pinnatis, pinna ſuprema triangulari, trilobata. *Hall. hiſt. n.* 18.

Sonchus lævior vulgaris ſecundus. *Cluſ. hiſt. p. CXLVI.*

In ſylvaticis, nemorenſibus. Fl. Iulio, Auguſto.

LEONTODON. Receptaculum convexum, nudum. Calyx oblongus, calyculatus duplici ſerie ſquamarum. Pappus ſtipitatus, ſimplex.

Taraxacum. 1. L. ſquamis calycinis exterioribus reflexis, foliis runcinatis acutis dentatis glabris. *Sp. pl. p.* 1122. *Curtis lond Fl. dan. t.* 574. *Scop. carn. tom.* 2. *p.* 99.

Taraxacum calycibus glabris, ſquamis imis reflexis. *Hall. hiſt. n.* 56.

Taraxacum officinale. *Wigg. prim.* 56. *Sibthorp oxon. p.* 239.

Le Piſſenlit, ou Dent de Lion. Regn. bot.

In pratis, paſcuis, cultis, alibique. Fl. a primo vere ad ſerum autumnum. Per.

HIERACIUM. Receptaculum nudum. Calyx imbricatus, ovatus. Pappus ſimplex, ſeſſilis.

incanum. 1. H. foliis integerrimis ſubdenticulatis lanceolatis ſcabris, ſcapo unifloro. *Syſt. veget. p.* 716. *Iacqu. auſtr. t.* 287.

Apargia incana. *Scop. carn. tom.* 2. *p.* 113.

Hieracium ſextum montanum. *Cluſ. hiſt. p. CXLI.*

In collibus, montibus Auſtriæ, Carnioliæ. Fl. Majo, Iunio. Per.

Folia ſunt rariter dentata, utrinque pilis albis apice ſtellatis incana. Pappus ſeſſilis, plumoſus. Iacqu. l. c.

alpinum. 2. H. foliis oblongis integris dentatis, ſcapo ſubnudo unifloro, calyce piloſo. *Sp. pl. p.* 1124. *Scop. carn. tom.* 2. *p.* 106. *Fl. dan. t.* 27. *Allion. pedem. tom.* 1. *p.* 212. *t.* 14. *f.* 2.

Hieracium caule unifloro, foliis lingulatis, dentatis, hirſutis, calyce hirſuto. *Hall. hiſt. n.* 49.

In alpibus ſtyriacicis, carniolicis, tyrolenſibus. Fl. Iulio, Auguſt. Per.

Tota planta pilis, baſi nigris, valde hirſuta. Scapus modo uniflorus, modo biflorus.

Taraxaci. 3. H. foliis lanceolatis dentatis glabris, ſcapo ſubnudo unifloro, calyce hirſuto. *Sp. pl. p.* 1125. *Allion. pedem. tom.* 1. *p.* 211. *t.* 31. *f.* 1.

Picris caule hirſuto, unifloro, foliis glabris, ſemipinnatis, calycibus hirſutiſſimis. *Hall. hiſt. n.* 27.

In alpibus. Fl. Iulio, Auguſto. Per.

alpeſtre. 4. H. foliis lanceolatis dentatis glabris, ſcapo ſubunifolio & unifloro, calyce hirſuto cylindrico. *Syſt. veget. p.* 716. *Iacqu. auſtr. t.* 191.

In pratis Auſtriæ ſubalpinis. Fl. Iunio, Iulio. Per.

Folia radicalia plura, oblonga, magis minusve profunde dentata, ſæpe inferne pinnatifida. Folia duo, vel unicum in ſuperiori caulis parte. Iacqu. l. c.

Piloſella. 5. H. foliis integerrimis ovatis ſubtus tomentoſis, ſtolonibus repentibus, ſcapo unifloro. *Sp. pl. p.* 1125. *Curtis lond.*

Hieracium caule unifloro, ſtolonibus reptantibus, foliis petiolatis, ovatis, longe piloſis, ſubtus tomentoſis. *Hall. hiſt. n.* 55.

La Piloſelle. Regn. bot.

In campis, pratis, collibus. Fl. æſtate. Per.

dubium. 6. H. foliis integris ovato-oblongis, ſtolonibus repentibus, ſcapo nudo multifloro. *Sp. pl. p.* 1125.

Hieracium caule nudo, paucifloro, ſtolonibus reptantibus, foliis longe piloſis, lanceolatis. *Hall. hiſt. n.* 53.

In pratis, paſcuis, in uliginoſis. Fl. æſtate. Per.

Auricula. 7. H. foliis integerrimis lanceolatis, ſtolonibus repentibus, ſcapo nudo multifloro. *Sp. pl. p.* 1126.

Hieracium caule nudo, ſtolonibus reptantibus, foliis longe piloſis, lanceolatis, floribus umbellatis. *Hall. hiſt. n.* 52.

In pratis, collibus, ad agrorum margines. Fl. æſtate. Per.

cymoſum. 8. H. foliis lanceolatis integris piloſis, ſcapo ſubnudo baſi piloſo, floribus ſubumbellatis. *Sp. pl. p.* 1126. *Fl. dan. t.* 810.

Hieracium caule ſubnudo, foliis hirſutis, lanceolatis, floribus umbellatis, petiolis ramoſis. *Hall. hiſt. n.* 51.

Piloſella minori flore, hirſutior & altior non repens. *Bauh. hiſt.* 2. *p.* 1040.

In collibus agri vindobonenſis. Fl. Iunio, Iulio. Per.

præmorſum. 9. H. foliis ovatis ſubdentatis, ſcapo nudo ramoſo, floribus ſuperioribus primoribus. *Sp. pl. p.* 1126. *Fl. dan. t.* 942.

Hieracium caule nudo, foliis petiolatis, ovatis, integerrimis, hirſutis, floribus ſpicatis. *Hall. hiſt. n.* 51.

In collibus, paſcuis, arvis reſtibilibus Auſtriæ, Pannoniæ. Fl. Majo, Iunio. Per.

aurantiacum. 10. H. foliis integris, caule ſubnudo ſimpliciſſimo piloſo corymbifero. *Sp. pl. p.* 1126. *Iacqu. auſtr. t.* 410.

Hieracium caule ſubnudo, foliis ovatis, integris, floribus umbellatis, aurantiis. *Hall. hiſt. n.* 50.

In alpinis, & ſubalpinis. Fl. Iunio, Iulio, Auguſto. Per.

incarnatum. 11. H. foliis ovatis denticulatis hirſutis, caule ſubnudo multifloro. *Wulfen in Iacqu. collect. tom.* 2. *p.* 127. *ic. t.* 578.

Hieracium montanum. *Scop. carn. tom. 2. p. 106. t. 50.*

In montanis Carnioliæ, Carinthiæ. Fl. Majo, Iunio. Per.

montanum. 12. H. caule folioſo ſubunifloro. *Syſt. veget. p. 717. Iacqu. auſtr. t. 190.*

Hieracium caule unifloro, foliis caulinis ovato-lanceolatis, dentatis, amplexicaulibus. *Hall. hiſt. n. 38.*

Hypochæris montana. *Sp. pl. p. 1140.*

In alpium herbidis. Fl. æſtate. Per.

porrifolium. 13. H. caule ramoſo folioſo, foliis lanceolato-linearibus ſubintegerrimis. *Syſt. veget. p. 717. Iacqu. auſtr. t. 286.*

In rupeſtribus ſubalpinis, & alpinis. Fl. Iulio, Auguſto. Per.

ſaxatile. 14. H. caule ramoſo folioſo, foliis anguſte lanceolatis utrinque acuminatis, baſi piloſis. *Iacqu. collect. tom. 1. p. 159. ic. t. 163. obs. 2. p. 30. t. 50.*

In rupeſtribus, aſperis Auſtriæ prope *Medling*, *Briel*, *Baaden*, *Bertholdsdorf*, alibique. Fl. æſtate. Per.

ſtaticifolium. 15. H. ſcapo nudo ramoſo, foliis lanceolato-linearibus obſolete denticulatis obtuſis: pedunculis longiſſimis unifloris, ſuperne ſquamoſis. *Hænke in Iacqu. collect. tom. 2. p. 34.*

Hieracium caule ramoſo, foliis radicalibus linearibus ſubdenticulatis, caulinis minimis. *Ger. prov. p. 167.*

In Carinthia in monte *Loibel*. In Auſtria in arenoſis prope *Baaden*. Fl. æſtate. Per.

Variat caule unifloro.

chondrilloides. 16. H. caule ramoſo, foliis caulinis elongato-dentatis glabris; radicalibus lanceolatis integris, caule ramoſo. *Sp. pl. p.* 1128. *Iacqu. auſtr. t.* 429.

In ſaxoſis alpinis. Fl. Iulio, Auguſto, Septembri. Per.

murorum. 17. H. caule ramoſo, foliis radicalibus ovatis dentatis; caulino minori. *Sp. pl. p.* 1128. *Scop. carn. tom.* 2. *p.* 108. *Allion. pedem. tom.* 1. *p.* 215. *t.* 28. *f.* 1.

Hieracium caule ſubnudo, paucifloro, foliis ovato-lanceolatis, circa petiolum dentatis. *Hall. hiſt. n.* 46.

In collibus, montanis ſylvaticis. Fl. æſtate. Per.

humile. 18. H. caule paucifloro foliis inferioribus hirſutis & inciſis vix altiore. *Syſt. veget. p.* 717.

Hieracium humilum. *Iacqu. auſtr. t.* 189.

In Auſtria in rupeſtribus alpinis, & ſubalpinis. Fl. Iunio, Iulio. Per.

paludoſum. 19. H. caule paniculato, foliis amplexicaulibus dentatis glabris, calycibus hiſpidis. *Sp. pl. p.* 1129. *Scop. carn. tom.* 2. *p.* 109. *Allion. pedem. tom.* 1. *p.* 216. *t.* 28. *f.* 2. *t.* 31. *f.* 2.

Hieracium foliis glabris, dentatis, imis petiolatis, ſuperioribus amplexicaulibus, calycibus hirſutis. *Hall. hiſt. n.* 45.

In udis ſylvaticis, montanis, & alpinis Styriæ, Carnioliæ, Carinthiæ, Tyrolis. Fl. æſtate. Per.

pyrenaicum. 20. H. foliis amplexicaulibus obovato-lanceolatis retrorſum dentatis, caule ſimplici, calycibus laxis. *Syſt. veget. p.* 718.

Crepis auſtriaca. *Iacqu. auſtr. t.* 441.

In alpinis, & ſubalpinis. Fl. Iulio, Auguſto. Per.

intyba-ceum. 21. H. foliis oblongis aſperis lanceolatis dentatis, calyce hirſutiſſimo. *Hall. hiſt. n.* 41. *Iacqu. auſtr. app. t.* 43. *Allion. pedem. tom.* I. *p.* 217. *t.* 29. *f.* 3.

In alpibus ſtyriacis, carinthiacis, tyrolenſibus. Fl. æſtate. Per.

molle. 22. H. foliis lanceolatis ſubintegerrimis mollibus: inferioribus petiolatis, floribus ſubcorymboſis. *Syſt. veget. p.* 718. *Iacqu. auſtr. t.* 119.

Hieracium foliis lævibus, integerrimis, radicalibus longe petiolatis, ellipticis, caulinis amplexicaulibus, hamis obtuſis. *Hall. hiſt. n.* 47.

In Auſtriæ pratis ſubalpinis. Fl. Iulio, Auguſto. Per.

ſtipita-tum. 23. H. caule ſubunifolio ramoſo, foliis dentatis, pappo ſtipitato. *Syſt. veget. p.* 718. *Iacqu. auſtr. t.* 293.

Habitat & floret cum præc. Per.

villoſum. 24. H. caule ramoſo folioſo, foliis hirtis: radicalibus lanceolato-ovatis dentatis; caulinis amplexicaulibus cordatis. *Sp. pl. p.* 1130. *Iacqu. auſtr. t.* 87.

Hieracium foliis villoſis integerrimis, radicalibus ligulatis, caulinis ovatis, amplexicaulibus. *Hall. hiſt. n.* 44.

Hieracium quintum villoſum. *Cluſ. hiſt. p. CXLI.*

In alpinis, & ſubalpinis. Fl. a Iunio ad Septembrim.

ſpicatum. 25. H. caule multifloro, foliis amplexicaulibus piloſis, rariter dentatis. *Hall. hiſt. n.* 43. *Allion. pedem. tom.* I. *p.* 218. *t.* 27. *f.* 3.

Hieracium *prenanthoides*; caule recto ſummo conice ramoſiſſimo, foliis ellipticis hirſutis, baſi amplexicaulibus. *Vill. proſp. p.* 35. *t.* 9. *f.* 2. *t.* 10. *f.* 1.

In ſubalpinis, & alpinis. Fl. a Iunio ad Septembrim. Per.

ſabaudum. 26. H. caule erecto multifloro, foliis ovato-lanceolatis dentatis ſemiamplexicaulibus, *Sp. pl. p.* 1131. *Scop. carn. tom.* 2. *p.* 109. *Allion. pedem. tom.* 1. *p.* 218. *t.* 27. *f.* 2.

Hieracium foliis hirſutis, dentatis, inferioribus ellipticis, ſuperioribus lanceolatis. *Hall. hiſt. n.* 35.

Hieracium ſecundum majus latifolium. *Cluſ. hiſt. p.* CXL.

In montanis ſylvaticis. Fl. Iulio, Auguſto. Per.

umbellatum. 27. H. foliis linearibus ſubdentatis ſparſis, floribus ſubumbellatis. *Sp. pl. p.* 1131. *Fl. dan. t.* 68c.

Hieracium foliis linearibus, ſubdentatis, ſparſis, floribus umbellatis. *Hall. hiſt. n.* 34.

Hieracii tertii genus alterum anguſtifolium. *Cluſ. hiſt. p.* CXL.

In ſylvaticis, ad vineas, vias. Fl. Iulio, Auguſto. Per.

CREPIS. Receptaculum nudum. Calyx calyculatus ſquamis difformibus, deciduis. Pappus piloſus.

fœtida. 1. C. foliis runcinato-pinnatis hirtis, petiolis dentatis. *Sp. pl. p.* 1133.

Crepis foliis ſcabris pinnatis, pinnis anguloſis, dentatis, retroverſis, extrema triangulari maxima. *Hall. hiſt. n.* 29.

Hieracium luteum, cichorii ſylveſtris folio amygdalas amaras olens. *Moris. hiſt.* 3. *p.* 63. *ſect.* 7. *t.* 4. *f.* 4.

In arvis, vineis, collibus, ad vias. Fl. æſtate.

tectorum. 2. C. foliis lanceolato-runcinatis ſeſſilibus lævibus: inferioribus dentatis. *Syſt. veget. p.* 719. *Curtis lond. Scop. carn. tom.* 2. *p.* 98.

Crepis foliis radicalibus pinnatis ſuperne amplexicaulibus, haſtatis. *Hall. hiſt. n.* 31.

Leontodon autumnale. *Fl. dan. t.* 501.

In pratis, collibus, ad vias. Fl. æſtate.

biennis. 3. C. foliis runcinato-pinnatifidis ſcabris: baſi ſuperne dentatis, calycibus muricatis. *Sp. pl. p.* 1136.

Crepis foliis ſubſcabris, ſemipinnatis, pinna extrema maxima. *Hall. hiſt. n.* 30.

In pratis, ad agrorum margines, ad vias. Fl. æſtate.

virens. 4. C. foliis runcinatis glabris amplexicaulibus, calycibus ſubtomentoſis. *Sp. pl. p.* 1134.

Crepis caule ramoſiſſimo, foliis glaberrimis, integris & dentatis. *Hall. hiſt. n.* 33.

In arenoſis, in arvis, ad vias, margines ſylvarum Styriæ, Auſtriæ. Fl. æſtate.

pulchra. 5. C. foliis ſagittatis dentatis, caule paniculato, calycibus pyramidatis glabris. *Sp. pl. p.* 1134.

Hieracium annuum montanum, caule canaliculato. *Moris. hiſt.* 3. *p.* 68. *ſect.* 7. *t.* 5. *f.* 37.

In Litoralis herbidis, ad vias. Fl. æſtate.

HYOSERIS. Receptaculum nudum. Calyx ſubæqualis. Pappus ſeſſilis, paleaceo-ariſtatus, capillari-cinctus; in ſeminibus marginalibus breviſſimus.

fœtida. 1. H. ſcapis unifloris, foliis pinnatifidis, ſeminibus nudis. *Sp. pl. p.* 1137.

Lapſana caule nudo indiviſo; foliis ſemipinnatis, pinnis retroverſis, dentatis. *Hall. hiſt. n.* 5.

Lampſana fœtida, *Scop. carn. tom.* 2. *p.* 118.

Leontodoides alpinus glaber, eryſimi folio, radice craſſa fœtida. *Mich. gen. p.* 31. *t.* 28.

In Carniolie, Carinthiæ, Litoralis alpinis, ſubalpinis, montaniſque. Fl. Majo, Iunio. Per.

LAPSANA. Receptaculum nudum. Calyx calyculatus, ſquamis ſingulis interioribus canaliculatis.

[illegible] 1. L. calycibus fructus angulatis, pedunculis tenuibus ramoſiſſimis. *Sp. pl. p.* 1141. *Fl. dan. t.* 500. *Curtis lond. Scop. carn. tom.* 2. *p.* 117.

Lapſana caule brachiato, foliis ovatis, longe petiolatis, petiolis pinnatis. *Hall. hiſt. n.* 6.

La Lampſane. Regn. bot.

In ſylvis, cultis, ad ſepes. Fl. Iunio, Iulio.

Zacintha. 2. L. calycibus fructus toruloſis depreſſis obtuſis ſeſſilibus. *Sp. pl. p.* 1141.

Cichorium verrucatum, Zacintha. *Clus. hiſt. p.* CXLIV.

In Litoralis, Iſtriæ cultis, ad ſepes. Fl. Majo, Iunio.

ſtellata. 3. L. calycibus fructus undique patentibus: radiis ſubulatis, foliis caulinis lanceolatis indiviſis. *Sp. pl. p.* 1141.

In Litorali, Iſtria. Fl. primo vere.

CICHORIUM. Receptaculum ſubpaleaceum. Calyx calyculatus. Pappus ſubquinquedentatus, obſolete piloſus.

Intybus. 1. C. floribus geminis ſeſſilibus, foliis runcinatis. *Sp. pl. p.* 1142. *Fl. dan. t.* 907. *Scop. carn. tom.* 2. *p.* 119.

Cichoreum foliis pinnatis, pinnis triangularibus dentatis, floribus ſeſſilibus. *Hall. hiſt. n.* 1.

Cichorium ſylveſtre ſeu officinarum. *Bauh. pin.* 125.

In pratis, paſcuis, collibus, ad vias. Fl. æſtate. Per.

Variat foliis integris; floribus cæruleis, rubris, albis.

HYPOCHÆRIS. Receptaculum paleaceum: Calyx ſubimbricatus. Pappus plumoſus.

maculata. 1. H. caule ſubnudo, ramo ſolitario, foliis ovato-oblongis integris dentatis. *Sp. pl. p.* 1140. *Fl. dan. t.* 149.

Hypochæris caule unifloro, foliis dentatis hirſutis. *Hall. hiſt. n.* 2. *t.* 1.

Hypochæris helvetica. *Wulfen in Iacqu. miſcell. tom.* 2. *p.* 25. *ic. t.* 165. *Syſt. veget. p.* 721.

Achyrophorus maculatus. *Scop. carn. tom.* 2. *p.* 116.

Hieracium latifolium primum. *Clus. hiſt. p.* CXXXIX.

In pratis, paſcuis, & in alpibus. Fl. æſtate. Per.

Varietas caule unifloro in alpibus ſtyriacis, carinthiacis, & tyrolenſibus creſcens in hortum tranſlata jam ſecundo anno ita mutatur, ut eam a planta in demiſſis creſcente diſtinguere non valeas.

radicata. 2. H. foliis runcinatis obtuſis ſcabris, caule ramoſo nudo lævi, pedunculis ſquamoſis. *Sp. pl. p.* 1140. *Fl. dan. t.* 150. *Curtis lond.*

Hypochæris foliis ſemipinnatis hirſutis, caule nudo brachiato. *Hall. hiſt. n.* 3.

Achyrophorus radicatus. *Scop. carn. tom.* 2. *p.* 117.

In pratis, paſcuis. Fl. æſtate. Per.

SCOLYMUS. Receptaculum paleaceum. Calyx imbricatus ſpinoſus. Pappus nullus. Semen biariſtatum.

maculatus. 1. S. floribus ſolitariis. *Syſt. veget. p.* 722. *Scop. carn. tom.* 2. *p.* 120.

Scolymus Theophraſti narbonenſis. *Clus. hiſt. p.* CLIII.

Scolymus chryſanthemos. *Bauh. pin.* 384. *Zann. iſt. p.* 240. *t.* 175.

In Litorali ad vias, hortos, alibique. In agro vindobonenſi prope *Schœnbrunn.* Fl. Iulio, Auguſto. Per.

** Capitati.*

ARCTIUM. Calyx globoſus: ſquamis apice hamis inflexis.

Lappa. 1. A. foliis cordatis inermibus petiolatis. *Sp. pl. p.* 1143. *Fl. dan. t.* 642. *Curtis lond. Scop. carn. tom.* 2. *p.* 122. *Hall. hiſt. n.* 161.

Lappa major ſeu Arctium Dioscoridis. *Bauh. pin.* 198.

La Bardane, ou le Glouteron. Regn. bot.

In ruderatis, ad ſepes, vias. Fl. æſtate. Per.

Perſonata. 2. A. foliis decurrentibus ciliato-ſpinoſis: radicalibus pinnatis; caulinis oblongo-ovatis. *Sp. pl. p.* 1144.

Carduus inermis, foliis imis pinnatis, ſuperioribus ovato-lanceolatis, ſerratis, ſquamis calycinis reflexis. *Hall. hiſt. n.* 162. *t.* 3.

Carduus Perſonata. *Iacqu. auſtr. t.* 348.

In montanis, ad torrentes, fluvios Styriæ, Tyrolis. In Auſtriæ ſubalpinis. Fl. Iulio, Auguſto. Per.

Carduelis. 3. A. foliis pinnatifidis aculeatis. *Syſt. veget. p.* 723.

Cirſium Arctioides. *Scop. carn. tom.* 2. *p.* 124. *t.* 53.

In Carniolia in alpibus wochinenſibus. Fl. Iulio, Auguſto. Per.

SERRATULA. Calyx ſubcylindricus, imbricatus, muticus.

tinctoria. 1. S. foliis lyrato-pinnatifidis: pinna terminali maxima, floſculis conformibus. *Sp. pl. p.* 1144. *Fl. dan. t.* 281.

Carduus inermis, foliis glabris, ſerratis, imis ovatis, ſuperioribus ſemipinnatis. *Hall. hiſt. n.* 163.

Carduus tinctorius. *Scop. carn. tom.* 2. *p.* 132.

Serratula Matthioli. *Clus. hiſt. p. VIII.*

In pratis, ad ſylvarum margines, & alibi. Fl. æſtate. Per.

alpina. 2. S. calycibus ſubhirſutis ovatis, foliis indiviſis. *Sp. pl. p.* 1145. *Fl. dan. t.* 37.

Cirſium foliis ovato-lanceolatis, dentatis, ſubtus tomentoſis, calycibus villoſis. *Hall. hiſt. n.* 179. *t.* 6.

Carduus mollior ſecundus. *Clus. hiſt. p. CLI.*

In alpibus. Fl. Iulio, Auguſto. Per.

arvenſis. 3. S. foliis dentatis ſpinoſis. *Sp. pl. p.* 1149. *Fl. dan. t.* 644. *Curtis lond.*

Cirſium caule anguloſo, ramoſo, foliis ſemipinnatis, pinnis anguloſis, ſpinoſis, calycibus inermibus, cylindricis. *Hall. hiſt. n.* 171.

Le Chardon hemorroidal. Regn. bot.

In cultis, aſperis. Fl. æſtate. Per.

Variat foliis utrinque unicoloribus, & facie viridibus, dorſo albidis.

CARDUUS. Calyx ovatus, imbricatus ſquamis ſpinoſis. Receptaculum piloſum.

lanceolatus. 1. C. foliis decurrentibus pinnatifidis hiſpidis: laciniis divaricatis, calycibus ovatis ſpinoſis villoſis, caule piloſo. *Sp. pl. p.* 1149.

Cirſium foliis ſemipinnatis, pinnis bilobatis, ſpinoſis, capitulis congeſtis, tomentoſis, conicis. *Hall. hiſt. n.* 169.

Cirſium lanceolatum. *Scop. carn. tom.* 2. *p.* 130.

Ad ſepes, vias, in ſylvaticis, & alibi. Fl. æſtate.

nutans. 2. C. foliis ſemidecurrentibus ſpinoſis, floribus cernuis: ſquamis calycinis ſuperne patentibus. *Sp. pl. p.* 1150. *Fl. dan. t.* 675. *Scop. carn. tom.* 2. *p.* 131.

Carduus foliis ſemipinnatis, pinnis angulofis, ſpinoſis, floribus nutantibus, calycinis ſquamis pungentibus, patulis. *Hall. hiſt. n.* 167.

Cirſium tertium Dodonæi. *Clus. hiſt. p.* CL.

Ad vias, pagos, in campeſtribus. Fl. æſtate.

acanthoides. 3. C. foliis decurrentibus ſinuatis margine ſpinoſis, calycibus pedunculatis ſolitariis, erectis villoſis. *Sp. pl. p.* 1150. *Iacqu. auſtr. t.* 249.

Cum præcedente. Fl. a Iulio ad hyemem.

criſpus. 4. C. foliis decurrentibus ſinuatis: margine ſpinoſis, floribus aggregatis terminalibus, ſquamis inermibus ſubariſtatis patulis. *Sp. pl. p.* 1150. *Fl. dan. t.* 621.

In Auſtriæ, Pannoniæ nemoribus, ſylvaticis humidis. Fl. æſtate.

paluſtris. 5. C. foliis decurrentibus dentatis: margine ſpinoſis, floribus racemoſis erectis, pedunculis inermibus. *Sp. pl. p.* 1151. *Curtis lond.*

Cirſium caule recto, ſubnudo, alato, foliis ſemipinnatis, ſpinoſis, calycibus ariſtatis. *Hall. hiſt. n.* 170.

Cirſium paluſtre. *Scop. carn. tom. 2. p.* 128.

In udis, paluſtribus. Fl. æſtate.

canus. 6. C. foliis decurrentibus lanceolatis eroſo - dentatis ciliato - aculeatis : utrinque arachnoideo - ſubvilloſis. *Syſt. veget. p.* 725. *Iacqu. auſtr. t.* 42. 43.

Cirſium maximum montanum incano folio, bulboſa radice. *Clus. hiſt. p.* CXLIX.

In pratis humidis Auſtriæ, Pannoniæ. Fl. Auguſto, Septembri. Per.

defloratus. 7. C. foliis decurrentibus lanceolatis ſerratis ſubſpinoſo - ciliatis nudis, pedunculis longiſſimis lanuginoſis unifloris. *Sp. pl. p.* 1152. *Iacqu. auſtr. t.* 89.

Carduus inermis, ſcapo nudo paucifloro. *Hall. hiſt. n.* 164.

Cirſium defloratum. *Scop. carn. tom.* 2. *p.* 127.

Cirſium tertium montanum. *Clus. hiſt. p.* CXLIX.

In montanis, alpinis, ſubalpinis. Fl. Iulio, Auguſto. Per.

Occurrit in alpibus varietas, ſpinis valde horrens.

marianus. 8. C. foliis amplexicaulibus haſtato - pinnatifidis ſpinoſis, calycibus aphyllis: ſpinis canaliculatis duplicato - ſpinoſis. *Sp. pl. p.* 1153. *Curtis lond.*

Silybum nervis foliorum albis. *Hall. hiſt. n.* 181.

Cirſium maculatum. *Scop. carn. tom.* 2. *p.* 130.

Carduus albis maculis notatus, vulgaris. *Bauh. pin.* 381. *Zann. iſt. p.* 51. *t.* 111.

Le Chardon Marie. Regn. bot.

Ad vias Litoralis. Fl. Iunio.

eriophorus. 9. C. foliis ſeſſilibus bifariam pinnatifidis: laciniis alternis erectis, calycibus globoſis villoſis. *Sp. pl. p.* 1153. *Iacqu. auſtr. t.* 171.

Cirſium foliis pinnatis, pinnis bilobatis, lanceolatis, ſpinoſis, calycibus ſphæricis, ſpinoſis, tomentoſis. *Hall. hiſt. n.* 168.

Cirſium eriophorum. *Scop. carn. tom.* 2. *p.* 130.

Carduus eriocephalus. *Clus. hiſt. p.* CLIV.

In ſubalpinis, montanis. Fl. Iulio, Auguſto. Per.

hetero-phyllus. 10. C. foliis amplexicaulibus lanceolatis ciliatis integris laciniatisque, caule ſubunifloro, calyce inermi. *Sp. pl. p.* 1154.

Cirſium foliis ciliatis, ſubtus tomentoſis, radicalibus petiolatis, ovato-lanceolatis & ſemipinnatis. *Hall. hiſt. n.* 180. *t.* 7.

In alpinis, & ſubalpinis Styriæ, Carinthiæ, Tyrolis. Fl. Iulio, Auguſto. Per.

Ex unica radice exeunt caules plures, quidam foliis omnibus integris, alii laciniatis, nonnulli integris & laciniatis veſtiti.

panno-nicus. 11. C. foliis ſemidecurrentibus nudis indiviſis ciliatis, flore ſubſolitario. *Syſt. veget. p.* 725.

Carduus ſerratuloides. *Iacqu. auſtr. t.* 127.

Cirſium pannonicum primum pratenſe. *Clus. hiſt. p.* CXLVIII.

In montoſis aſperis Auſtriæ, Pannoniæ. Fl. Iunio, Iulio. Per.

Folia caulina ſunt longe lanceolata, ſemiamplexicaulia, acuminata: facie virentia, piloſa & ſcabra; dorſo arachnoideo-lanuginoſa & pallentiora, craſſiuſcula, obſolete ſerrulata, & ciliata ſpinulis innocuis, infima obtuſiora & obſolete decurrentia, ſuperiora gradatim minora. Iacqu. l. c.

tatari-cus. 12. C. foliis amplexicaulibus lanceolatis: ſerraturis ſpinoſo-ſetaceis, floribus triphyllis. *Sp. pl. p.* 1155. *Iacqu. auſtr. t.* 90.

In Auſtriæ udis. Fl. Iulio. Per.

rivularis. 13. C. foliis ſeſſilibus ſerratis : inferne pinnatifidis, ſuperne integris, caule inermi, floribus capitatis. *Syſt. veget. p.* 727. *Iacqu. auſtr. t.* 91.

In Auſtria, Pannonia ad rivulos, in pratis udis. Fl. Iunio, Iulio. Per.

mollis. 14. C. foliis pinnatifidis linearibus ſubtus tomentoſis, caule inermi unifloro. *Sp. pl. p.* 1156. *Iacqu. auſtr. t.* 18.

Carduus mollior primus. *Clus. hiſt. p.* CL.

Cirſium molle. *Scop. carn. tom.* 2. *p.* 126.

In collibus asperis. Fl. Majo, Iunio.

acaulis. 15. C. acaulis, calyce glabro. *Sp. pl. p.* 1156. *Wulfen in Iacqu. colle͡ct. tom.* 1. *p.* 239. *ic. t.* 579.

Cirſium floribus acaulis. *Hall. hiſt. n.* 178.

Carlina minor, purpureo flore. *Clus. hiſt. p.* CLVI.

In herbidis apricis, ad vias Carnioliæ calidioris, Forijulii. Fl. Iulio, Auguſto. Per.

CNICUS. Calyx ovatus, imbricatus ſquamis ramoſo-ſpinoſis, obvalatus bra͡cteis. Corollulæ æquales.

oleraceus. 1. C. foliis pinnatifidis carinatis nudis, bra͡cteis concavis integris ſubcoloratis. *Sp. pl. p.* 1156. *Fl. dan t.* 860.

Cirſium foliis ciliatis, amplexicaulibus, inermibus, flores comprehendentibus. *Hall. hiſt. n.* 173.

Cirſium oleraceum. *Scop. carn. tom.* 2. *p.* 124.

Ad fluvios, in pratis humidis. Fl. a Iulio ad ſerum Autumnum. Per.

Eriſithales. 2. C. foliis amplexicaulibus pinnatifidis ariſtato-ſerratis: floribus cernuis, calycibus glutinoſis. *Sp. pl. p.* 1157.

Cirſium foliis ciliatis, imis ovatis, ſuperioribus ſemipinnatis, floribus ſeſſilibus obtuſis. *Hall. hiſt. n.* 175.

Cirſium Eriſithales. *Scop. carn. tom.* 2. *p.* 125.

Carduus Eriſithales. *Iacqu. auſtr. t.* 310.

In ſubalpinis, & alpinis. Fl. Iulio, Auguſto. Per.

pygmæus. 3. C. foliis ſeſſilibus ſublinearibus confertiſſimis, calice inermi. *Sp. pl. p.* 1156.

Serratula pygmæa. *Iacqu. auſtr. t.* 440.

Cirſium pygmæum. *Scop. carn. tom.* 2. *p.* 123.

Carduus mollior humilis anguſtifolius. *Clus. hiſt. p.* CLI.

In alpibus auſtriacis, ſtyriacis, carniolicis. Fl. Iulio, Auguſto. Per.

ſpinoſiſſimus. 4. C. foliis amplexicaulibus ſinuato-pinnatis ſpinoſis, caule ſimplici, floribus ſeſſilibus. *Sp. pl. p.* 1157.

Cirſium foliis caulinis ſemipinnatis, pinnis anguloſis, ſpinoſiſſimis terminantibus mollibus, ciliatis, umbellam floralem comprehendentibus. *Hall. hiſt. n.* 172. *t.* 5.

Cirſium ſpinoſiſſimum. *Scop. carn. tom.* 2. *p.* 129.

In Carniolia in alpibus wochinenſibus. In Tyroli ad torrentes alpinos, alibique. Fl. Iulio, Auguſto. Per.

ONOPORDUM. Receptaculum favoſum. Calycis ſquamæ mucronatæ.

Acanthium. 1. O. calycibus ſquamoſis: ſquamis patentibus, foliis ovato-oblongis ſinuatis. *Sp. pl. p.* 1158. *Fl. dan. t.* 909. *Curtis lond. Scop. carn. tom.* 2. *p.* 132.

Onopordum caule alato, foliis ovatis, dentatis, dentibus anguloſis, ariſtatis. *Hall. hiſt. n.* 159.

In ruderatis, ad vias. Fl. æſtate.

CARLINA. Calyx radiatus ſquamis marginalibus longis, coloratis.

acaulis. 1. C. caule unifloro flore breviore. *Sp. pl. p.* 1160. *Scop. carn. tom.* 2. *p.* 133.

Carlina caule unifloro. *Hall. hiſt. n.* 183.

Chamæleon album Dioscoridis, Ixia Theophraſti. *Clus. hiſt. p. CLV.*

La Carline, ou le Chameleon blanc. Regn. bot.

In collibus, campis, pratis ſiccioribus. Fl. Iunio, Iulio. Per.

acanthifolia. 2. C. acaulis, foliis tomentoſis pinnatifidis plicatis, pinnis ſemibilobis pungentibus. *Allion. pedem. tom.* 1. *p.* 156. *t.* 51.

Carlina *chardouſſe*, acaulis, foliis ſinuato-pinnatifidis palearum ſegmentis ſubulato-incraſſatis. *Vill. proſp. p.* 27.

Carlina Utzka. *Hacqu. alp. p.* 9. *t.* 1.

In Litorali in monte majore. *Hacquet.* Fl. Iulio. Per.

lanata. 3. C. caule bifido, calycibus ſanguineis terminalibus: primo axillari ſeſſili. *Syſt. veget. p.* 728.

Carlina ſquamis calycinis terminalibus purpuraſcentibus, floribus inferioribus proliferis. *Gerard prov.* 183.

Carlina Pola. *Hacqu. alp. p.* 29. *t.* 4. *f.* 3.

Carlina flore purpuro-rubente. *Tourn. inſt.* 500. *Garid. t.* 21.

In Iſtria veneta circa Polam. *Hacquet. I. Hoſt.* Fl. Iulio, Auguſto.

corymboſa. 4. C. caule multifloro ſubdiviſo, floribus ſeſſilibus, calycibus radio flavo. *Sp. pl. p.* 1160. *Wulfen in Iacqu. collect. tom.* 4. *p.* 306.

In Litoralis aſperis. Fl. Iunio, Iulio.

vulgaris. 5. C. caule multifloro corymboſo, floribus terminalibus, calycibus radio albo. *Sp. pl. p.* 1161.

Carlina foliis radicalibus ſemipinnatis, pinnis angulofis, ſpinoſis, caulinis ovato-lanceolatis, amplexicaulibus. *Hall. hiſt. n.* 182.

Carlina ſylveſtris vulgaris. *Clus. hiſt. p. CLV.*

In collibus, montoſis, aridis. Fl. æſtate.

** *Discoidei.*

BIDENS. Receptaculum paleaceum. Pappus ariſtis erectis, ſcabris. Calyx imbricatus. Corolla rarius flosculo uno alterne radiante inſtruitur.

triparti-ta. 1. B. foliis trifidis, calycibus ſubfolioſis, ſeminibus erectis. *Sp. pl. p.* 1165. *Scop. carn. tom.* 2. *p.* 178.

Bidens foliis petiolatis trilobatis, & quinquelobatis, ſerratis, floribus circumvallatis. *Hall. hiſt. n.* 121.

In aquoſis. Fl. æſtate.

cernua. 2. B. foliis lanceolatis amplexicaulibus, floribus cernuis. *Sp. pl. p.* 1165. *Curtis lond. Fl. dan. t.* 841. *Scop. carn. tom.* 2. *p.* 177.

Bidens foliis ſeſſilibus ſerratis, floribus nutantibus, circumvallatis. *Hall. hiſt. n.* 120.

In aquoſis, pratis udis. Fl. Iulio, Auguſto.

Variat floribus radiatis. Hæc varietas eſt Coreopſis Bidens. Linn.

CACALIA. Receptaculum nudum. Pappus piloſus. Calyx cylindricus, oblongus, baſi tantum ſubcalyculatus.

alpina. 1. C. herbacea, foliis reniformi - cordatis acutis denticulatis. *Syſt. veget. p.* 734. *Iacqu. auſtr. t.* 234.

In ſubalpinis. Fl. Iulio, Auguſto. Per.

albifrons. 2. C. herbacea, foliis cordatis biſerratis acutis ſubtus tomentoſis, ſtipulis oblongis rotundatis. *Syſt. veget. p.* 734.

Cacalia tomentoſa. *Iacqu. auſtr. t.* 235.

In alpibus. Fl. cum præc. Per.

EUPATORIUM. Receptaculum nudum. Pappus plumoſus. Calyx imbricatus, oblongus. Stylus longus, ſemibifidus.

cannabinum. 1. E. foliis digitatis. *Sp. pl. p.* 1173. *Fl. dan. t.* 745. *Scop. carn. tom.* 2. *p.* 155.

Eupatorium foliis ovato - lanceolatis, tripteris. *Hall. hiſt. n.* 136.

L' Eupatoire d' Avicenne. Regn. bot.

In udis. Fl. Iulio, Auguſto. Per.

STÆHELINA. Receptaculum breviſſime paleaceum. Pappus ramoſus. Antheræ caudatæ.

dubia. 1. S. foliis linearibus denticulatis, ſquamis calycinis lanceolatis, pappo calycibus duplo longiore. *Sp. pl. p.* 1176. *Ger. prov. p.* 190. *t.* 6.

In inſulis venetis, copioſe Auxerii verſus l' *Iſola longa. I. Hoſt.* Fl. Iunio. Fr.

CHRYSOCOMA. Receptaculum nudum. Pappus ſimplex. Calyx hemisphæricus, imbricatus. Stylus vix floſculis longior.

Linoſyris. 1. C. herbacea foliis linearibus glabris, calycibus laxis. *Sp. pl. p.* 1178. *Hall. hiſt. n.* 144. *Allion. pedem. tom.* 1. *p.* 174. *t.* 11. *f.* 2.

Oſyris auſtriaca. *Clus. hiſt. p.* 325.

In Auſtriæ, Styriæ collibus, montoſis ſterilibus. Fl. a Iulio ad Octobrim. Per.

SANTOLINA. Receptaculum paleaceum. Pappus nullus. Calyx imbricatus, hemisphæricus.

roſmarinifolia. 1. S. pedunculis unifloris, foliis linearibus margine tuberculatis. *Sp. pl. p.* 1180.

Abrotonum femina quartum. *Clus. hiſt. p.* 342.

Le petit Cyprés; ou la Garde-Robe. Regn. bot.

In rupeſtribus Iſtriæ. *I. Hoſt.* Ego in ſimilibus locis agri fluminenſis in *Raſtozzine.* Fl. Iunio, Iulio. Fr.

POLYGAMIA SUPERFLUA.

* *Discoidei.*

TANACETUM. Receptaculum nudum. Pappus ſubmarginatus. Calyx hemisphæricus, imbricatus. Corollæ radii obſoletæ, trifidæ.

vulgare. 1. T. foliis bipinnatis inciſis ſerratis. *Sp. pl. p.* 1184. *Fl. dan. t.* 871. *Scop. carn. tom.* 2. *p.* 143.

Tanacetum foliis pinnatis, pinnis ſemipinnatis, acute dentatis. *Hall. hiſt. n.* 132.

Tanacetum vulgare luteum. *Bauh. pin.* 132.

La Tanaiſie. Regn. bot.

Ad fluvios, ſepes, & alibi. Fl. Iulio, Auguſto.

Iuveni varietatem floribus albis in Croatia inter Fucine & Merslagora.

ARTEMISIA. Receptaculum ſubvilloſum vel nudiuſculum. Pappus nullus. Calyx imbricatus ſquamis rotundatis, conniventibus. Corollæ radii nullæ.

Abrotanum. 1. A. fruticoſa, foliis ſetaceis ramoſiſſimis. *Sp. pl. p.* 1185. *Scop. carn. tom.* 2. *p.* 144.

Abrotanum mas anguſtifolium majus. *Bauh. pin.* 136. *Zann. iſt. p.* 1. *t.* 30.

In aſperis Litoralis. Fl. Auguſto.

campeſtris. 2. A. foliis multifidis linearibus, caulibus procumbentibus virgatis. *Sp. pl. p.* 1185. *Scop. carn. tom.* 2. *p.* 145.

Artemiſia caulibus declinatis, foliis ſericeis, calveſcentibus, pinnatis; pinnis multifidis, acutis. *Hall. hiſt. n.* 131.

Artemiſia tenuifolia ſecunda. *Clus. hiſt. p.* 340.

Abrotanum campeſtre. *Bauh. pin.* 136.

In campis, collibus, montanis aſperis. Fl. æſtate. Per.

erithmifolia. 3. A. foliis compoſitis divaricatis linearibus carnoſis glabris, caule adſcendente panniculato. *Sp. pl. p.* 1186.

Abrotanum hiſpanicum maritimum, folio craſſo ſplendente & rigido. *Tourn. inſt.* 459.

In aſperis, arenoſis Litoralis. Fl. Auguſto. Per.

glacialis. 4. A. foliis palmatis multifidis ſericeis, caulibus adſcendentibus, floribus glomeratis faſtigiatis. *Sp. pl. p.* 1187. *Iacqu. auſtr. app. t.* 35. *Wulſen in Iacqu. collect. tom.* 2. *p.* 118. *t.* 7. *f.* 1. 2. 3.

Abſinthium foliis ſericeis petiolatis pinnatis, pinnulis trifidis & quinquefidis. *Hall. hiſt. n.* 125.

Abſinthium alpinum candidum humile. *Bauh. pin.* 139. *prodr.* 71.

In alpibus carniolicis, carinthiacis, tyrolenſibus. Fl. Iulio, Auguſto. Per.

ſpicata. 5. A. foliis radicalibus biternatis, caule adſcendente ſpicato, floribus erectis. *Iacqu. auſtr. app. t.* 34.

Artemiſia foliis ſericeis, radicalibus petiolatis, pinnatis, inciſis, caulinis ſemipinnatis. *Hall. hiſt. n.* 127.

In Tyroli in alpibus leontinis. Fl. Iulio, Auguſto. Per.

pontica. 6. A. foliis multipartitis ſubtus tomentoſis, floribus ſubrotundis nutantibus: receptaculo nudo. *Sp. pl. p.* 1187. *Iacqu. auſtr. t.* 99.

Artemiſia foliis ſericeis, duplicato-pinnatis, pinnulis parallelis. *Hall. hiſt. n.* 129.

Abſinthium ponticum vulgare. *Cluſ. hiſt. p.* 339.

Abſinthium ponticum tenuifolium, incanum. *Bauh. pin.* 138.

In ſiccis, apricis Auſtriæ, Pannoniæ. Fl. Autumno. Per.

Folia caulina ſunt bipinnata, utrinque tomentoſa, facie glauca, dorſo magis incana, plana: laciniis linearibus, acutis. Ramea inferiora & caulina ſuperiora ſunt ſimpliciter pinnata. Summa autem omnia integerrima & ſimpliciſſima. Iacqu. l. c.

auſtriaca. 7. A. foliis multipartitis tomentoſis incanis, floribus oblongis nutantibus: receptaculis nudis. *Syſt. veget. p.* 744. *Iacqu. auſtr. t.* 100.

Abſinthium auſtriacum tenuifolium. *Cluſ. hiſt. p.* 339.

In apricis ſiccis Auſtriæ, Pannoniæ. Fl. Octobri. Per.

Folia ſunt tomentoſa, ex glauco incana; plana: in ramulis primo prodeuntibus integra; ſubſequa in elongatis trifida, atque ſic paulatim diviſione increſcentia, ut tandem evadant ſubtriplicato-pinnata: laciniis in prioribus magis obtuſis & patentibus; in poſterioribus magis anguſtis, linearibus, acutioribus & conniventibus. Summa iterum decreſcunt. Iacqu. l. c.

Abſinthium. 8. A. foliis compoſitis multifidis, floribus ſubgloboſis pendulis: receptaculo villoſo. *Sp. pl. p.* 1188. *Scop. carn. tom.* 2. *p.* 144.

Abſinthium foliis incanis, pinnatis, pinnis primis bilobatis, pinnulis inciſis, lanceolatis, ſpicis heteromallis, floribus pendulis. *Hall. hiſt. n.* 124.

L' Abſinthe. Regn. bot.

Ad vias, pagos, in ruderatis, in apricis ſterilibus. Fl. Iulio, Auguſto. Per.

vulgaris. 9. A. foliis pinnatifidis planis inciſis ſubtus tomentoſis, racemis ſimplicibus recurvatis, floribus radio quinquefloro. *Sp. pl. p.* 1188. *Scop. carn. tom.* 2. *p.* 144.

Artemiſia foliis inferne tomentoſis, pinnatis, pinnis acute dentatis, ſpica panniculata erecta. *Hall. hiſt. n.* 130.

L' Armoiſe. Regn. bot.

In ruderatis, ad ſepes, & alibi. Fl. Iulio, Auguſto. Per.

caeruleſcens. 10. A. foliis caulinis lanceolatis integris: radicalibus multifidis, floſculis femineis ternis. *Sp. pl. p.* 1189. *Scop. carn. tom.* 2. *p.* 145.

Abſinthium maritimum, lavendulae folio. *Bauh. pin.* 139. *Zann. iſt. p.* 245. *t.* 156. 139.

Abſinthium marinum patavinum. *Clus. hiſt. p.* 338.

In Litorali ad mare. Fl. Auguſto. Per.

Folia radicalia caulinaque inferiora citius marceſcunt, quam reliqua: hinc ſit, quod plerumque in planta florente folia omnia indiviſa inveniantur.

GNAPHALIUM. Receptaculum nudum. Pappus plumoſus. Calyx imbricatus ſquamis marginalibus rotundatis, ſcarioſis, coloratis.

Stœchas. 1. G. fruticoſum, foliis linearibus, corymbo compoſito, ramis virgatis. *Sp. pl. p.* 1193. *Scop. carn. tom.* 2. *p.* 154.

Filago foliis linearibus, oris contractis, ramis ſimpliciſſimis, umbella convexa terminatis. *Hall. hiſt. n.* 145.

Elichryſium, ſeu Stœchas citrina anguſtifolia. *Bauh. pin.* 264. *Zann. iſt. p.* 92. *t.* 279.

In aſperis, ſaxoſis agri fluminenſis. Fl. Iunio, Iulio.

arenarium. 2. G. herbaceum, foliis lanceolatis: inferioribus obtuſis, corymbo compoſito, caulibus ſimpliciſſimis. *Sp. pl. p.* 1195. *Fl. dan. t.* 641.

Stœchas citrina germanica, latiore folio. *Bauh. hiſt.* 3. *p.* 153.

In agri vindobonenſis collibus prope *Weinhaus.* Fl. æſtate. Per.

luteo-album. 3. G. herbaceum, foliis ſemiamplexicaulibus enſiformibus repandis obtuſis utrinque pubeſcentibus, floribus conglomeratis. *Sp. pl. p.* 1196.

Filago foliis molliſſimis, tomentoſis, obtuſis, umbellis convexis, electrinis. *Hall. hiſt. n.* 147.

Chryſocome citrina ſupina, latifolia, italica. *Barr. ic.* 367.

Elichryſium ſylveſtre latifolium capitulis conglobatis. *Bauh. pin.* 264. *Zann. iſt. p.* 92.

In arenoſis Litoralis. Fl. Iulio.

dioicum. 4. G. ſarmentis procumbentibus, caule ſimpliciſſimo, corymbo ſimplici, floribus dioicis. *Sp. pl.* *p.* 1199. *Scop. carn. tom.* 2. *p.* 149.

Filago flagellis reptans, ſexu diſtincta, flosculis omnibus androgynis. *Hall. hiſt. n.* 157.

Piloſella minor. *Clus. hiſt. p.* 330.

La Perliere dioïque. Bulliard. herb. t. 325.

In collibus, pratis, montanis ſylvaticis. Fl. Majo, Iunio. Per.

Flores feminini elongantur, dum mares eandem formam, & figuram globoſam conſervant.

ſylvaticum. 5. G. caule herbaceo ſimpliciſſimo erecto, floribus ſparſis. *Sp. pl. p.* 1200. *Scop. carn. tom.* 2. *p.* 151. *t.* 56.

Filago foliis linearibus, alis ſpiciferis. *Hall. hiſt. n.* 148.

In montanis ſylvaticis, & in alpibus. Fl. æſtate. Per.

In alpibus ſtyriacis, carinthiacis, & tyrolenſibus occurrit varietas floribus nigerrimis, in ſpicam brevem congeſtis.

uliginoſum. 6. G. caule herbaceo ramoſo diffuſo, floribus confertis terminalibus. *Sp. pl. p.* 1200. *Fl. dan. t.* 859.

Filago caule ramoſiſſimo, diffuſo, umbella ſeſſili terminato. *Hall. hiſt. n.* 151.

In uliginoſis. Fl. Iulio, Auguſto.

XERANTHEMUM. Receptaculum paleaceum. Pappus ſetaceus. Calyx imbricatus, radiatus: radio colorato.

annuum. 1. X. herbaceum, foliis lanceolatis patentibus. *Sp. pl. p.* 1201. *Iacqu. auſtr. t.* 388.

Xeranthemum herbaceum, foliis lanceolatis. *Hall. hiſt. n.* 122.

Ptarmica auſtriaca. *Cluſ. hiſt. p. XI.*

In Auſtriæ, Pannoniæ, Litoralis campeſtribus, collibus aridis, arvis. Fl. Iulio, Auguſto.

Flores feminei ſunt in ambitu, habentque limbum patulum, cujus tres laciniæ longæ ſunt & æquales, duæ vero ambitum reſpicientes duplo breviores; habent quoque ſtylum petalo duplo longiorem. In hermaphroditis germen quinque ſetis tanquam calyculo proprio inſtruitur, quæ in femineis deſunt. Iacqu. l. c.

CARPESIUM. Receptaculum nudum. Pappus nullus. Calyx imbricatus: ſquamis exterioribus reflexis.

cernuum. 1. C. floribus terminalibus. *Sp. pl. p.* 1203. *Iacqu. auſtr. t.* 204. *Scop. carn. tom.* 2. *p.* 142. *Hall. hiſt. n.* 134.

In Auſtria ad ſepes, in herbidis udis prope *Hütteldorf.* Vulgare in Carnioliæ convallibus, Forjulii humidis. Fl. Iulio, Auguſto. Per.

abrotanoides. 2. C. floribus lateralibus. *Sp. pl. p.* 1204. *Wulfen in Iacqu. collect. tom.* 1. *p.* 267. *Osb. it. t.* 10.

In umbroſis, ſucculentis agri aquilejenſis. Fl. Iulio.

CONYZA. Receptaculum nudum. Pappus ſimplex. Calyx imbricatus, ſubrotundus. Corollæ radii trifidæ.

ſquarroſa. 1. C. foliis lanceolatis acutis, caule herbaceo corymboſo, calycibus ſquarroſis. *Sp. pl. p.* 1205. *Fl. dan. t.* 622. *Scop. carn. tom.* 2. *p.* 154.

Conyza foliis ovato-lanceolatis, rariter ſerratis, floribus umbellatis. *Hall. hiſt. n.* 135.

Conyza Matthioli. Bacharis monſpelienſium. *Cluſ. hiſt. p. XXI.*

In collibus, ſylvaticis. Fl. Iulio, Auguſto. Per.

** *Radiati.*

ERIGERON. Receptaculum nudum. Pappus piloſus. Corollæ radii lineares, anguſtiſſimæ.

canadenſe. 1. E. caule floribus paniculatis, hirtis, foliis lanceolatis ciliatis. *Syſt. veget. p.* 754. *Scop. carn. tom.* 2. *p.* 159.

Erigeron floribus paniculatis ſpicatis. *Hall. hiſt. n.* 84.

In cultis, & incultis. Fl. æſtate.

acre. 2. E. pedunculis alternis unifloris. *Sp. pl. p.* 1211.

Erigeron caule alterne ramoſo, petiolis unifloris, ſemifloſculis pappum æquantibus. *Hall. hiſt. n.* 85.

In collibus apricis, montanis ſterilibus, muris, cultis. Fl. æſtate. Per.

alpinum. 3. E. caule ſubbifloro, calyce ſubhirſuto, foliis obtuſis ſubtus villoſis. *Syſt. veget. p.* 754. *Scop. carn. tom.* 2. *p.* 160.

Erigeron caule alterne ramoſo, petiolis unifloris, ſemifloſculis pappum ſuperantibus. *Hall. hiſt. n.* 86.

In alpinis, & ſubalpinis. Fl. Iulio, Auguſto. Per.

uniflorum. 4. E. caule unifloro, calyce piloſo. *Sp. pl. p.* 1211. *Fl. dan. t.* 292.

Erigeron caule unifloro, calyce tomentoſo, radiis ſemifloſculoſis. *Hall. hiſt. n.* 87.

Aſter caule unifloro, foliis integerrimis, calyce villoſo, ſubrotundo. *Fl. lapp. n.* 307. *t.* 9. *f.* 3.

In alpibus tyrolenſibus. Fl. Auguſto.

TUSSILAGO. Receptaculum nudum. Pappus ſimplex. Calycis ſquamæ æquales, discum æquantes, ſubmembranaceæ.

alpina. 1. T. ſcapo unifloro ſubnudo, foliis cordato-orbiculatis crenatis, utrinque virentibus. *Syſt. veget. p.* 755. *Iacqu. auſtr. t.* 246.

Tuſſilago alpina ſecunda glabro folio. *Cluſ. hiſt. p. CXIII.*

In alpinis, & ſubalpinis. Fl. Iulio, Auguſto. Per.

Folia ſunt ſupra glabra, nec lucida; ſubtus virentia & villoſa, nec tomentoſa aut incana. Iacqu. l. c.

discolor. 2. T. ſcapo unifloro ſubnudo, foliis cordato-orbiculatis crenatis ſupra lucidis, ſubtus tomentoſo-incanis. *Syſt. veget. p.* 755. *Iacqu. auſtr. t.* 247.

Tuſſilago alpina. *Scop. carn. tom.* 2. *p.* 156.

Tuſſilago alpina prima canescente folio. *Cluſ. hiſt. p. CXII.*

Habitat & floret cum præc. Per.

Folia radicalia, ſubrotunda, cordata, acute crenulata, craſſa, firma: prona parte læte virentia, lucida, glaberrima; dorſo vero denſiſſimo breviſſimoque tomento, quod adulta minime dejiciunt, incana. Iacqu. l. c.

ſylveſtris. 3. T. ſcapo ſubunifloro ſubnudo, foliis cordato-orbiculatis, inciſo-lobatis. *Syſt. veget. p.* 755. *Iacqu. auſtr. app. t.* 12. *Scop. carn. tom.* 2. *p.* 157.

In montanis ſylvaticis, ſubalpinis Styriæ, Carnioliæ, Croatiæ. Fl. Majo, Iunio. Per.

Radix non repens ut in præcedentibus. Folia radicalia reniformi-ſubrotunda, utrinque virentia. Folia caulina a duobus ad quinque: inferiora ex petiolo latiſſimo, concavo & amplexicauli oriuntur, lobate inciſa; ſumma ſeſſilia, lanceolata & integra. Flos plerumque unicus, huic quandoque ex foliorum ſummorum alis flos pedunculatus unus alterve accedit. Iacqu. l. c.

Farfara. 4. T. ſcapo unifloro imbricato, foliis ſubcordatis angulatis denticulatis. *Sp. pl. p.* 1214. *Curtis lond. Fl. dan. t.* 595. *Scop. carn. tom.* 2. *p.* 159.

Petaſites ſcapo unifloro, flore radiato. *Hall. hiſt. n.* 143.

Le Tuſſilage. Regn. bot.

In pratis humidis, in uliginoſis, ad torrentes, fluvios. Fl. primo vere. Per.

frigida. 5. T. thyrſo faſtigiato, floribus radiatis. *Sp. pl. p.* 1214. *Fl. dan. t.* 61.

Petaſites floribus ſpicatis, erectis, foliis calycinis lanceolatis, floſculis pauciſſimis androgynis. *Hall. hiſt. n.* 141.

In alpium udis, præcipitiis. Fl. Iunio. Per.

alba. 6. T. thyrſo faſtigiato, floſculis femineis nudis paucis. *Sp. pl. p.* 1214. *Fl. dan. t.* 524.

Petaſites ſpicis laxis, umbellatis, floſculis plerisque androgynis. *Hall. hiſt. n.* 139.

In ſylvis Croatiæ ſubalpinis. Fl. Iunio. Per.

hybrida. 7. T. thyrſo oblongo, floſculis femineis nudis plurimis. *Sp. pl. p.* 1214.

Petaſites floribus ſpicatis, pendulis, petiolis multifloris, calycibus æqualibus, floſculis pauciſſimis androgynis. *Hall. hiſt. n.* 140.

Ad fluvios, torrentes. Fl. primo vere. Per.

Petaſites. 8. T. thyrſo ovato, floſculis femineis nudis paucis. *Sp. pl. p.* 1215. *Fl. dan.* 842.

Petaſites floribus denſe ſpicatis, floſculis androgynis. *Hall. hiſt. n.* 138.

Le Petaſite. Regn. bot.

Habitat & floret cum præc. Per.

T. frigida donatur quandoque floribus radio deſtitutis. In Petaſite flores feminei deſunt. In frigida, hybrida, alba, & Petaſite ſtigma quandoque tam craſſum eſt, ut egrediendo tubum ab antheris, margine coalitis, formatum dilaceret, faciatque, ut eæ liberæ appareant.

SENECIO. Receptaculum nudum. Pappus ſimplex. Calyx cylindricus, calyculatus: ſquamis apice ſphacelatis.

vulgaris. 1. S. corollis nudis, foliis pinnato-ſinuatis amplexicaulibus, floribus ſparſis. *Sp. pl. p.* 1216. *Curtis lond. Fl. dan. t.* 513. *Scop. carn. tom.* 2. *p.* 162. *Hall. hiſt. n.* 58.

Le Seneçon. Regn. bot.

In cultis, ruderatis. Fl. a primo vere ad hyemem.

viscoſus. 2. S. corollis revolutis, foliis pinnatifidis viſcidis, ſquamis calycinis laxis longitudine perinanthii. *Sp. pl. p.* 1217.

Senecio foliis viſcidis, pinnatis, pinnis ſemipinnatis, anguloſis, calycibus hirſutis. *Hall. hiſt. n.* 60.

Senecio hirſutus, viſcidus, graveolens. *Dill. elth.* 347. *t.* 258. *f.* 336.

In montanis ſterilibus. Fl. æſtate.

ſylvaticus. 3. S. corollis revolutis, foliis pinnatifidis denticulatis, caule corymboſo erecto. *Sp. pl. p.* 1217. *Fl. dan. t.* 869.

Senecio foliis pinnatis, pinnis alterne majoribus, ſemipinnatis, anguloſis, calycibus lævibus, radiis breviſſimis revolutis. *Hall. hiſt. n.* 59.

In montanis ſylvaticis, copioſe ad pedes alpium. Fl. æſtate.

ſqualidus. 4. S. corollis radiantibus calyce longioribus integris, foliis pinnatifidis: laciniis linearibus diſtantibus. *Sp. pl. p.* 1218.

In Carnia. *I. Hoſt.* Fl. Iunio, Iulio.

incanus. 5. S. corollis radiantibus, foliis utrinque tomentoſis ſubpinnatis obtuſis, corymbo ſubrotundo. *Sp. pl. p.* 1219.

Senecio tomentoſus, foliis pinnatis, pinnis ſubrotundis, umbella globoſa. *Hall. hiſt. n.* 61.

Chryſanthemum alpinum primum. *Clus. hiſt. p.* 333.

In alpibus ſtyriacis, carinthiacis, tyrolenſibus. Fl. Iulio, Auguſto. Per.

abrotanifolius. 6. S. corollis radiantibus, foliis pinnato-multifidis linearibus nudis acutis, pedunculis ſubbifloris. *Syſt. veget. p.* 757. *Iacqu. auſtr. t.* 79. *Scop. carn. tom.* 2. *p.* 167.

Solidago foliis vaginoſis, pinnatis, pinnis perangustis, acutis, ſimplicibus & bifidis. *Hall. hiſt. n.* 71.

Chryſanthemum alpinum ſecundum. *Clus. hiſt. p.* 333.

Chryſanthemum alpinum, foliis abrotani multifidis. *Bauh. pin.* 134.

In alpinis, & ſubalpinis. Fl. Iulio, Auguſto. Per.

Iacobæa. 7. S. corollis radiantibus, foliis pinnato-lyratis: laciniis lacinulatis, caule erecto. *Sp. pl. p.* 1220. *Fl. dan. t.* 944. *Scop. carn. tom.* 2. *p.* 167.

Iacobæa vulgaris. *Clus. hiſt. p. XXII.*

In pratis, collibus, montanis. Fl. æſtate. Per.

aquaticus. 8. S. corollis radiantibus, foliis dentatis: radicalibus ovatis, caulinis pinnatifidis: lacinia extima majore. *Hud. angl.* 2. 366. *Retzii ſcand. p.* 195. *Fl. dan. t.* 784.

Iacobæa tertia latifolia. *Cluſ. hiſt. p. XXIII.*

In humidis. Fl. æſtate. Per.

tenuifolius. 9. S. corollis radiantibus, foliis omnibus pinnatis, pinnis linearibus, pedunculis corymboſis, caule erecto. *Syſt. veget. p.* 758. *Iacqu. auſtr. t.* 278.

In Auſtria prope *Bertholdsdorf*, *Medling*, *Giſshübel*, alibique ad maceries, ſepes, vineas. In agri fluminenſis montanis ſylvaticis. Fl. Auguſto, Septembri. Per.

Differt a S. Iacobæa floribus odoratis minoribus; tempore florendi magis autumnali; calycum apicibus pallentibus nec fuſcis; foliisque tenuioribus & omnibus ſubpinnatis; radice repente. Iacqu. l. c.

paludoſus. 10. S. corollis radiantibus, foliis enſiformibus acute ſerratis ſubtus ſubvilloſis, caule ſtricto. *Sp. pl. p.* 1220. *Fl. dan. t.* 385. *Scop. carn. tom.* 2. *p.* 163.

Senecio foliis linearibus ſerratis, floribus umbellatis. *Hall. hiſt. n.* 66.

Conyza paluſtris, ſerratifolia. *Bauh. pin.* 266. *Zann. iſt. p.* 146. *t.* 131.

In Littorali in paluſtribus aquilejenſibus, duinenſibus. In Auſtria in ſimilibus locis prope *Rohrau*, *Bruck an der Leitha*. Fl. Iulio, Auguſto. Per.

nemorenſis. 11. S. corollis radiantibus octonis, foliis lanceolatis biſerratis ſubtus villoſis, caule ramoſo. *Sp. pl. p.* 1221. *Iacqu. auſtr. t.* 184.

Senecio foliis ovato-lanceolatis, ſerratis, ſubtus hirſutis, floribus umbellatis, *Hall. hiſt. n.* 64.

In nemorenſibus. Fl. Iulio, Auguſto. Per.

ſarracenicus. 12. S. corollis radiantibus, floribus corymboſis, foliis lanceolatis ſerratis glabriusculis. *Sp. pl. p.* 1221. *Iacqu. auſtr. t.* 186.

Senecio foliis lanceolatis, ſerratis, glabris, floribus umbellatis. *Hall. hiſt. n.* 65.

In Auſtria, Pannonia in inſulis danubialibus. Fl. Iulio, Auguſto. Per.

Doria. 13. S. corollis radiantibus, floribus corymboſis, foliis ſubdecurrentibus nudis lanceolatis denticulatis, ſuperioribus ſenſim minoribus. *Sp. pl. p.* 1221. *Iacqu. auſtr. t.* 185.

Ad ſepes, in pratis humidis: in Pannonia prope Poſonium, *Laide*, alibique; in Croatia prope Varasdinum. In Auſtria prope *Bruck an der Leitha.* Fl. Iulio, Auguſto. Per.

Folia craſſa, ferme glauca, & ad tactum pinguia indicant vel ſola Senecionem Doriam. Radix valde repens, folia anguſta, conferta & flores radiati octoni S. ſarracenicum. Senecio nemorenſis totus odoratus habet corollas radiantes quinas. Iacqu. l. c.

Doronicum. 14. S. corollis radiantibus, caule indiviſo ſubunifloro, foliis indiviſis ſerratis: radicalibus ovatis ſubtus villoſis. *Sp. pl. p.* 1222. *Iacqu. auſtr. app. t.* 45. *Scop. carn. tom.* 2. *p.* 163.

Senecio foliis craſſis, ſerratis, imis petiolatis ſubrotundis, caulinis lanceolatis. *Hall. hiſt. n.* 67.

In alpibus carniolicis, carinthiacis, tyrolenſibus. Fl. Iulio, Auguſto. Per.

Variat caule multifloro, unifloro.

ASTER. Receptaculum nudum. Pappus ſimplex. Corollæ radii plures. Calyces imbricati: ſquamis inferioribus patulis.

alpinus. 1. A. foliis ſubſpathulatis hirtis integerrimis, caulibus ſimplicibus, unifloris. *Ait. Kew. tom.* 3. *p.* 198. *Sp. pl. p.* 1226. *Iacqu. auſtr. t.* 88.

Aſter foliis radicalibus petiolatis, ovatis, ſcabris, caule unifloro. *Hall. hiſt. n.* 82.

Aſter ſeptimus, auſtriacus quintus. *Cluſ. hiſt. p. XV.*

Aſter hirſutus auſtriacus cœruleus, magno flore, foliis ſubrotundis. *Bauh. pin.* 267. *prodr.* 124.

In ſummis alpium auſtriacarum, ſtyriacarum jugis. Fl. Iulio, Auguſto. Per.

Tripolium. 2. A. foliis lineari-lanceolatis integerrimis carnoſis glabris trinerviis, foliolis calycinis ſubmembranaceis obtuſis. *Ait. Kew. tom.* 3. *p.* 199. *Sp. pl. p.* 1226. *Scop. carn. tom.* 2. *p.* 170.

Aſter pannonicus. *Iacqu. hort. tom.* 1. *p.* 3. *t.* 8.

In Auſtria in paluſtribus prope *Bruck an der Leitha.* In Pannonia ad lacum Peiſonis. In Litorali in paludibus duinenſibus, aquilejenſibus. Fl. a Iunio ad ſerum Autumnum. Per.

Amellus. 3. A. foliis oblongo-lanceolatis integris ſcabris, ramis corymboſis, calycibus imbricatis ſubſquarroſis: foliolis obtuſis: interioribus membranaceis apice coloratis. *Ait. Kew. tom.* 3. *p.* 199. *Sp. pl. p.* 1226. *Iacqu. auſtr. t.* 425. *Scop. carn. tom.* 2. *p.* 170.

Aſter caule folioſo, foliis ſcabris, ciliatis, lanceolatis, obtuſis, calycinis ſquamis ovatis, ciliatis. *Hall. hiſt. n.* 83.

Aſter octavus italorum & Fuchſii. *Cluſ. hiſt. p. XVI.*

Aſter atticus cœruleus vulgaris. *Bauh. pin.* 267.

In collibus, montanis apricis, ſylvaticis. Fl. Auguſto, Septembri. Per.

SOLIDAGO. Receptaculum nudum. Pappus ſimplex. Corollæ radii circiter quinque. Squamæ calycinæ imbricatæ, clauſæ.

Virgaurea. 1. S. caule erecto tereti pubescente ſuperne ramoſo, foliis inferioribus ellipticis piloſiuſculis ſerratis, racemis erectis, ligulis elongatis. *Ait. Kew. tom.* 3. *p.* 218. *Sp. pl. p.* 1235. *Fl. dan. t.* 663.

Solidago caule anguloſo, hirſuto, foliis ovato-lanceolatis, dentatis, racemis multifloris ſpicatis. *Hall. hiſt. n.* 69.

Doria Virgaurea. *Scop. carn. tom.* 2. *p.* 176.

La Verge d'Or. Regn. bot.

In ſylvaticis, & in alpibus. Fl. Iulio, Auguſto. Per.

CINERARIA. Receptaculum nudum. Pappus ſimplex. Calyx ſimplex, polyphyllus, æqualis.

cordifolia. 1. C. panicula pauciflora, foliis caulinis petiolatis cordatis acute ſerratis glabris, caule angulato. *Syſt. veget. p.* 764. *Iacqu. auſtr. t.* 176. 177.

Senecio foliis cordato-lanceolatis ſerratis. *Hall. hiſt. n.* 63.

Senecio alpinus. *Linn. ſuppl.* 371. *Scop. carn. tom.* 2. *p.* 164.

Jacobæa pannonica tertia, latifolia prima. *Cluſ. hiſt. p. XXII.*

In alpinis, & ſubalpinis. Fl. Iulio, Auguſto. Per.

Huic ſunt petioli foliorum radicalium & caulinorum infimorum longi, nec vel minime alati.

crispa. 2. C. floribus paniculatis, foliis caulinis ſpathulato-oblongis ſerratis obtuſis amplexicaulibus, baſi dilatato-dentatis. *Syſt. veget. p.* 764. *Iacqu. auſtr. t.* 178.

In alpibus. Fl. Iulio, Auguſto. Per.

Folia radicalia ſunt cordata & per totam petioli longitudinem utrinque decurrentia, ibidemque laciniato - inciſa. Caulina ſuperiora ſunt oblonga, acuta, argute & inæqualiter inciſo - ſerrata, ſeſſilia. Omnia ad margines criſpata & undulata. Iacqu. l. c.

integrifolia. 3. C. floribus umbellatis, foliis denticulatis lanceolatis, radicalibus ſpathulatis, *Syſt. veget. p.* 764. *Iacqu. auſtr. t.* 179.

Ad margines ſylvarum ſubalpinarum. Fl. Iunio, Iulio. Per.

campeſtris. 4. C. floribus corymboſis, foliis radicalibus petiolatis ovatis tomentoſis: caulinis ſeſſilibus linearibus. *Retzii ſcand. p.* 196. *Iacqu. auſtr. t.* 180.

Iacobæa pannonica ſecunda. *Cluſ. hiſt. p. XXII.*

In pratenſibus, ſylvaticis Auſtriæ, Pannoniæ. Fl. Iunio. Per.

longifolia. 5. C. foliis obſolete denticulatis, omnibus oblongis. *Syſt. veget. p.* 765. *Iacqu. auſtr. t.* 181.

In alpinis. Fl. Iulio. Per.

Huic ſunt folia omnia oblonga, dum integrifoliæ ſunt ferme integerrima, radicalia ſpathulata potius, quam vere petiolata. Iacqu. l. c.

INULA. Receptaculum nudum. Pappus ſimplex. Antheræ baſi in ſetas duas deſinentes.

Helenium. 1. I. foliis amplexicaulibus ovatis rugoſis, ſubtus tomentoſis, calycum ſquamis ovatis. *Sp. pl. p.* 1236. *Fl. dan. t.* 728.

Aſter foliis ovato - lanceolatis, ſerratis, ſubtus tomentoſis, calycinis ovato - lanceolatis maximis. *Hall. hiſt. n.* 72.

Aſter Helenium. *Scop. carn. tom.* 2. *p.* 171.

L' Aunée. Regn. bot.

In Croatiæ, agri fluminenſis montanis ſylvaticis. Occurrit quoque in Styriæ ſuperioris pomariis. Fl. Iunio. Per.

Oculus chriſti. 2. I. foliis amplexicaulibus oblongis integerrimis hirſutis, caule piloſo corymboſo. *Sp. pl. p.* 1237. *Iacqu. auſtr. t.* 223.

Conyza tertia auſtriaca. *Cluſ. hiſt. p. XX.*

In Auſtriæ, Pannoniæ aridis. Fl. Iulio. Per.

britannica. 3. I. foliis amplexicaulibus lanceolatis diſtinctis ſerratis ſubtus villoſis, caule ramoſo villoſo erecto. *Sp. pl. p.* 1237.

Aſter foliis ſubhirſutis, radicalibus petiolatis, ellipticis, caulinis ſeſſilibus, lanceolatis, calycinis perangustis. *Hall. hiſt. n.* 78.

Conyza aquatica, aſteris flore aureo. *Bauh. pin.* 266. *prodr.* 124.

In uliginoſis, humidis. Fl. Iulio, Auguſto. Per.

dyſenterica. 4. I. foliis amplexicaulibus cordato-oblongis, caule villoſo paniculato, ſquamis calycinis ſetaceis. *Sp. pl. p.* 1237. *Cur. lond.*

Aſter foliis amplexicaulibus, undulatis, ſubtus tomentoſis. *Hall. hiſt. n.* 79.

Aſter dyſentericus. *Scop. carn. tom.* 2. *p.* 171.

Conyza media vulgaris. *Cluſ. hiſt. p. XXI.*

L' Inule conizière. Bulliard herb. t. 299.

In paluſtribus, ad foſſas. Fl. Iulio, Auguſto. Per.

Bubonium. 5. I. foliis lanceolatis rigidulis denticulatis ſubvilloſis ſeſſilibus, caule ramisque ſubbifloris, calyce ſquarroſo. *Syſt. veget. p.* 767. *Iacqu. auſtr. app. t.* 19.

Aſter Bubonium. *Scop. carn. tom.* 2. *p.* 173. *t.* 58.

Ad ſepes, in herbidis Litoralis, Carnioliæ calidioris, Forijulii. Fl. Iulio, Auguſto. Per.

ſalicina. 6. I. foliis lanceolatis recurvis ſerrato-ſcabris, ramis angulatis, floribus inferioribus altioribus. *Sp. pl. p.* 1238. *Fl. dan. t.* 786.

Aſter foliis glabris, ciliatis, venoſis. *Hall. hiſt. n.* 76.

Aſter ſalicinus. *Scop. carn. tom.* 2. *p.* 172.

Aſter quartus. Auſtriacus ſecundus. *Clus. hiſt. p. XIV.*

Ad margines ſylvarum. Fl. Iulio, Auguſto. Per.

hirta. 7. I. foliis ſeſſilibus lanceolatis recurvis ſubſerrato-ſcabris, caule teretiuſculo ſubpiloſo, floribus inferioribus altioribus. *Sp. pl. p.* 1239. *Iacqu. auſtr. t.* 358.

Aſter foliis ovatis, venoſis, ſcabris, hirſutis. *Hall. hiſt. n.* 75.

Aſter hirtus. *Scop. carn. tom.* 2. *p.* 173. *t.* 58.

Aſter quintus. *Clus. hiſt. p. XIV.*

In collibus herbidis. Fl. Iunio. Per.

germanica. 8. I. foliis ſeſſilibus lanceolatis recurvis ſcabris, floribus ſubfaſciculatis. *Sp. pl. p.* 1239. *Iacqu. auſtr. t.* 134.

In aſperis, ad agrorum margines Auſtriæ, Pannoniæ. Fl. Iulio. Per.

enſifolia. 9. I. foliis ſeſſilibus linearibus acuminatis nervoſis glabris ſparſis, caule ſubunifloro. *Sp. pl. p.* 1240. *Kram. auſtr.* 253. *Iacqu. auſtr. t.* 162.

Aſter enſifolius. *Scop. carn. tom.* 2. *p.* 174.

Aſter anguſtifolius, ſive ſextus, auſtriacus quartus. *Clus. hiſt. p. XV.*

In collibus, montoſis apricis. Fl. Auguſto. Per.

crithmifolia. 10. I. foliis linearibus carnoſis tricuſpidatis. *Sp. pl. p.* 1240.

Senecio crithmifolius. *Scop. carn. tom.* 2. *p.* 163.

Aſter maritimus, folio tereti, craſſo, tridentato. *Tourn. inſt.* 483.

Crithmum maritimum, flore aſteris attici. *Bauh. pin.* 288. *Zann. iſt. p.* 29. *t.* 23.

In Litorali in rupibus, quas alluit mare adriaticum. Fl. Auguſto, Septembri. Per.

ARNICA. Receptaculum nudum. Pappus ſimplex. Corollulæ radii filamentis quinque abſque antheris.

montana. 1. A. foliis ovatis integris: caulinis geminis oppoſitis. *Sp. pl. p.* 1245. *Fl. dan. t.* 63. *Scop. carn. tom.* 2. *p.* 175.

Arnica foliis conjugatis, ovatis, integerrimis. *Hall. hiſt. n.* 90.

Doronicum quintum germanicum & ſextum pannonicum. *Clus. hiſt. p. XVIII.*

L' Arnica. Regn. bot.

In pratis alpinis & ſubalpinis. Copioſſime in Croatia prope *Fucine*; in Litorali in monte majore; in Styria ſuperiore in alpibus judenburgenſibus. Fl. æſtate. Per.

ſcorpioides. 2. A. foliis alternis dentatis mollibus. *Syſt. veget. p.* 768. *Iacqu. auſtr. t.* 349.

Arnica foliis alternis, inæqualiter ſinuato-denticulatis, radicalibus petiolatis. *Hall. hiſt. n.* 89.

Aſter ſcorpioides. *Scop. carn. tom.* 2. *p.* 169.

Doronicum quartum. *Clus. hiſt. p. XVII.*

In alpibus. Fl. Iulio, Auguſto. Per.

Doronicum. 3. A. foliis alternis ſubſerratis, oblongis aſperis. *Syſt. veget. p.* 768. *Iacqu. auſtr. t.* 92.

Arnica uniflora, foliis alternis ovato-lanceolatis, hirſutis, integerrimis, calyce hirto. *Hall. hiſt. n.* 91.

Doronicum ſecundum, auſtriacum primum. *Clus. hiſt. p. XVI.*

In ſaxoſis alpium auſtriacarum, ſtyriacarum. Fl. Iulio, Auguſto. Per.

DORONICUM. Receptaculum nudum. Pappus ſimplex. Calycis ſquamæ duplicis ordinis æquales, disco longiores. Semina radii nuda pappoque deſtituta.

Pardelianches. 1. D. foliis cordatis obtuſis denticulatis: radicalibus petiolatis; caulinis amplexicaulibus. *Sp. pl. p.* 1247. *Iacqu. auſtr. t.* 350.

Doronicum foliis cordatis, obtuſis, imis longe petiolatis, ſuperioribus amplexicaulibus. *Hall. hiſt. n.* 88.

Doronicum latifolium. *Clus. hiſt. p. XVI.*

In alpibus. Fl. Iulio, Auguſto. Per.

In Carniolia in alpibus wochinenſibus inveni varietatem ramoſam, qualem figura cluſiana ſiſtit.

auſtriacum. 2. D. foliis imis cordato-ovatis auriculatis: ſummis lanceolatis. *Iacqu. auſtr. t.* 130.

Doronicum ſeptimum, auſtriacum tertium. *Clus. hiſt. p. XIX.*

In alpinis, & ſubalpinis. Fl. Iulio, Auguſto. Per.

Bellidiaſtrum. 3. D. ſcapo nudo ſimpliciſſimo unifloro. *Sp. pl. p.* 1247. *Iacqu. auſtr. t.* 400.

Arnica caule nudo, unifloro, foliis ovato-lanceolatis, ſerratis. *Hall. hiſt. n.* 92.

After Bellidiaſtrum. *Scop. carn. tom.* 2. *p.* 168.

Bellis media. *Cluſ. hiſt. p. XLIV.*

In alpinis, & ſubalpinis. Fl. æſtate.

BELLIS. Receptaculum nudum, conicum. Pappus nullus. Calyx hemisphæricus: ſquamis æqualibus. Semina obovata.

perennis. 1. B. ſcapo nudo. *Sp. pl. p.* 1248. *Fl. dan. t.* 503. *Curtis lond. Scop. carn. tom.* 2. *p.* 146. *Hall. hiſt. n.* 93.

La Paquerette vivace. Bulliard herb. t. 173.

In pratis. Fl. a primo vere ad hyemem.

CHRYSANTHEMUM. Receptaculum nudum. Pappus marginatus. Calyx hemisphæricus, imbricatus: ſquamis marginalibus membranaceis.

atratum. 1. C. foliis omnibus cuneiformibus oblongis ſerratis carnoſis. *Sp. pl. p.* 1252.

In alpinis, & ſubalpinis. Fl. Iulio, Auguſto. Per.

alpinum. 2. C. foliis cuneiformibus pinnatifidis, laciniis integris, caulibus unifloris. *Sp. pl. p.* 1253.

Pyrethrum caule unifloro, foliis petiolatis palmatis. *Hall. hiſt. n.* 96.

Leucanthemum alpinum. *Cluſ. hiſt. p.* 335.

In alpibus ſtyriacis, carinthiacis, tyrolenſibus, carnicis. Fl. Iulio, Auguſto, Septembri. Per.

Leucanthemum. 3. C. foliis amplexicaulibus oblongis: ſuperne ſerratis, inferne dentatis. *Sp. pl. p.* 1251. *Curtis lond.*

Matricaria foliis radicalibus petiolatis, ovatis, crenatis, caulinis amplexicaulibus dentatis. *Hall. hiſt. n.* 98.

Matricaria Leucanthemum. *Scop. carn. tom. 2, p.* 147.

Le Leucantheme vulgaire. Regn. bot.

In pratis, collibus, Fl, Iunio, Iulio. Per,

montanum, 4. C. foliis imis ſpathulato-lanceolatis ſerratis: ſummis linearibus. *Sp. pl. p.* 1252.

Leucanthemum montanum minus, *Tourn. inſt.* 492.

In Carnioliæ, Croatiæ alpinis, & ſubalpinis, Fl, Iunio, Iulio. Per.

inodorum. 5. C. foliis pinnatis multifidis, caule ramoſo diffuſo. *Sp. pl. p.* 1253. *Fl. dan. t.* 696.

In Auſtria, Pannonia ad pagos, vias, in arvis, alibique, Fl, Iunio, Iulio.

corymboſum, 6. C. foliis pinnatis inciſo-ſerratis, caule multifloro. *Sp. pl. p.* 1251. *Iacqu. auſtr. t.* 379.

Pyrethrum foliis pinnatis, pinnis ſemipinnatis, acute dentatis, floribus umbellatis. *Hall. hiſt. n.* 95.

Pyrethrum corymboſum. *Scop. carn. tom.* 2, *p.* 148.

Tanacetum inodorum primum & ſecundum, *Clus. hiſt. p.* 337.

In collibus, montanis ſylvaticis, Fl. Iunio, Iulio. Per,

MATRICARIA, Receptaculum nudum. Pappus nullus, Calyx hemisphæricus, imbricatus: ſquamis marginalibus acutiuſculis, ſolidis.

Parthenium, 1. M. foliis compoſitis planis: foliolis ovatis inciſis, pedunculis ramoſis. *Sp. pl. p.* 1255. *Fl. dan. t.* 674.

Matricaria foliis pinnatis, pinnis ſemipinnatis, laciniis obtuſis, floribus umbellatis, *Hall. hiſt. n.* 100.

La Matricaire. Regn. bot.

In ruderatis, cultis, collibus aridis, ad pagos. Fl. æſtate. Per.

Chamomilla. 2. M. receptaculis conicis, radiis patentibus, ſquamis calycinis margine æqualibus. *Sp. pl. p.* 1256. *Curtis lond. Scop. carn. tom.* 2. *p.* 148.

Matricaria foliis capillaribus, duplicato-pinnatis, pinnis lanceolatis, bifidis, trifidisque. *Hall. hiſt. n.* 101.

In cultis, ruderatis, ad vias, pagos. Fl. Majo, Iunio.

ANTHEMIS. Receptaculum paleaceum. Pappus nullus. Calyx hemiſphæricus, ſubæqualis. Flosculi radii plures quam quinque.

alpina. 1. A. foliis dentato-pinnatis integerrimis linearibus, caule villoſo ſubunifloro, petalis ovatis, paleis ſphacelatis. *Sp. pl. p.* 1261. *Iacqu. auſtr. app. t.* 30.

In alpibus carniolicis, auſtriacis, tyrolenſibus. Fl. Iulio, Auguſto. Per.

Variat caule unifloro, bifloro, & multifloro.

arvenſis. 2. A. receptaculis conicis: paleis ſetaceis, ſeminibus coronato-marginatis. *Sp. pl. p.* 1261. *Scop. carn. tom.* 2. *p.* 178.

Chamæmelum foliis glabris, duplicato-pinnatis, nervo foliaceo, pinnulis lanceolatis, ſemine lævi. *Hall. hiſt. n.* 103.

In arvis. Fl. Iunio, Iulio.

auſtriaca. 3. A. receptaculis conicis: paleis oblongis mucronatis, ſeminibus nudis, foliis bipinnatis lanuginoſo-villoſis. *Syſt. veget. p.* 776. *Iacqu. auſtr. t.* 444.

In arvis Auſtriæ, Styriæ, Pannoniæ. Fl. æſtate.

tinctoria. 4. A. foliis bipinnatis ſerratis ſubtus tomentoſis, caule corymboſo. *Sp. pl. p.* 1263. *Fl. dan. t.* 741.

Chamæmelum foliis hirſutis, pinnatis, pinnulis ſimplicibus lanceolatis. *Hall. hiſt. n.* 105.

Buphthalmum vulgare chryſanthemo congener. *Clus. hiſt. p.* 332.

L' Oeil de Beuf. Regn. bot.

In collibus Auſtriæ, Pannoniæ. Fl. Iunio, Iulio. Per.

ACHILLEA. Receptaculum paleaceum. Pappus nullus. Calyx ovatus, imbricatus. Flosculi radii circiter quatuor.

Ageratum. 1. A. foliis lanceolatis obtuſis acute ſerratis. *Sp. pl. p.* 1264.

Ageratum foliis ſerratis. *Bauh. pin.* 221.

L' Eupatoire de Méſué. Regn. bot.

In agro tergeſtino. *I. Hoſt.* Fl. Auguſto, Septembri. Per.

macrophylla. 2. A. foliis pinnatis planis, inciſo-ſerratis: extimis majoribus coadunatis. *Sp. pl. p.* 1265.

Achillea foliis pinnatis, pinnis acute ſerratis, extremis confluentibus. *Hall. hiſt. n.* 115.

Dracunculus alpinus, foliis ſcabioſæ. *Bauh. pin.* 98. *prodr.* 39.

In Carnia in monte *Croſtis. I. Hoſt.* Fl. Iulio. Per.

Clavennæ. 3. A. foliis laciniatis planis obtuſis tomentoſis. *Sp. pl. p.* 1266. *Iacqu. auſtr. t.* 76. *Scop. carn. tom.* 2. *p.* 179.

Achillea foliis tomentoſis, longe petiolatis, palmatis. *Hall. hiſt. n.* 114.

Abſinthium alpinum umbelliferum. *Clus. hiſt. p.* 339.

In ſummis alpium jugis. Fl. Iulio, Auguſto. Per.

Ptarmica. 4. A. foliis lanceolatis acuminatis argute ſerratis. *Sp. pl. p.* 1266. *Curtis lond.*

Achillea foliis linearibus, lanceolatis, acutiſſime ſerratis. *Hall. hiſt. n.* 117.

Dracunculus pratenſis ſerrato folio. *Bauh. pin.* 98.

Ptarmica vulgaris. *Clus. hiſt. p. XII.*

L' Herbe á éternuer. Regn. bot.

In Carniæ pratis. *I. Hoſt.* Fl. Iulio. Per.

atrata. 5. A. foliis pinnulis pectinatis integriusculis, pedunculis villoſis. *Sp. pl. p.* 1267. *Iacqu. auſtr. t.* 77. *Scop. carn. tom.* 2. *p.* 179.

Achillea foliis glabris, pinnatis, pinnis ſimplicibus & multifidis. *Hall. hiſt. n.* 111.

Parthenium alpinum. *Clus. hiſt. p.* 336.

In ſummis alpium jugis. Fl. Auguſto, Septembri. Per.

moſchata. 6. A. foliis pinnatis, pinnulis ſimplicibus, floribus umbellatis. *Syſt. veget. p.* 778. *Iacqu. auſtr. app. t.* 33.

In alpibus tyrolenſibus. Fl. Iulio, Auguſto. Per.

magna. 7. A. foliis bipinnatis ſubpiloſis: laciniis linearibus dentatis: auriculis decuſſatis. *Sp. pl. p.* 1267. *Allion. pedem. tom.* 1. *p.* 184. *t.* 53. *f.* 1.

In Carnia in monte *Croſtis. I. Hoſt.* Fl. Iulio. Per.

Millefolium. 8. A. foliis bipinnatis nudis: laciniis linearibus dentatis, caulibus ſuperne ſulcatis. *Sp. pl. p.* 1267. *Curtis lond. Fl. dan. t.* 737. *Scop. carn. tom.* 2. *p.* 180.

Achillea foliis pinnatis, punctatis, pinnis æqualibus, pinnulis trifidis & quinquefidis. *Hall. hiſt. n.* 107.

La Mille-feuille. Regn. bot.

In pratis, ad agrorum margines, vias, & alibi. Fl. æſtate. Per.

nobilis. 9. A. foliis bipinnatis: inferioribus nudis planis, ſuperioribus obtuſis tomentoſis, corymbis convexis conſertiſſimis. *Sp. pl. p.* 1268.

Achillea foliis tomentoſis, duplicato-pinnatis, pinnis crenatis, pinnulis ſemipinnatis. *Hall. hiſt. n.* 109.

Tanacetum minus album, odore camphoræ. *Bauh. pin.* 132.

In apricis aridis Auſtriæ, Pannoniæ. Fl. Iulio, Auguſto. Per.

Calyces magis oblongi, quam in præcedente, cylindrici, radius quidem albus, ſed discus ochroleucus, & calycinarum ſquamarum margo minime nigricans. Retzii obs. faſc. 2. p. 26.

BUPHTHALMUM. Receptaculum paleaceum. Pappus: margo obſoletus. Seminum latera, præſertim radii, marginata. Stigma flosculorum hermaphroditicorum indiviſum.

aquaticum. 1. B. calycibus obtuſe folioſis ſeſſilibus axillaribus, foliis alternis oblongis obtuſis, caule herbaceo. *Sp. pl. p.* 1274.

Chryſanthemum conyzoides luſitanicum. *Breyn. cent.* 157. *t.* 77.

In paludibus aquilejenſibus. Fl. æſtate.

ſalicifolium. 2. B. folis alternis lanceolatis ſubſerratis villoſis, calycibus nudis, caule herbaceo. *Sp. pl. p.* 1275. *Iacqu. auſtr. t.* 370. *Scop. carn. tom.* 2. *p.* 177.

Buphthalmum foliis hirſutis, imis petiolatis, ovatis, ſuperioribus amplexicaulibus, lanceolatis, rariter dentatis. *Hall. hiſt. n.* 118.

Aſter tertius, auſtriacus primus. *Cluſ. hiſt. p. XIII.*

In collibus, montanis ſylvaticis. Fl. æſtate. Per.

POLYGAMIA FRUSTRANEA.

CENTAUREA. Receptaculum ſetoſum. Pappus ſimplex. Corollæ radii infundibuliformes, longiores, irregulares.

phrygia. 1. C. calycibus recurvato - plumoſis, foliis indiviſis oblongis ſcabris. *Syſt. veget. p.* 785. *Fl. dan. t.* 520. *Scop. carn. tom.* 2. *p.* 137.

Cyanus foliis ſubhirſutis, ovato - lanceolatis amplexicaulibus, dentatis, plumulis calycinis reflexis. *Hall. hiſt. n.* 188.

Iacea quarta auſtriaca villoſo capite. *Cluſ. hiſt. p. VI.*

In montanis ſylvaticis, & in alpibus. Fl. æſtate. Per.

In Carnioliæ alpibus wochinenſibus occurrit varietas caule unifloro.

nigra. 2. C. calycibus ciliatis: ſquamula ovata: ciliis capillaribus erectis, foliis lyrato - angulatis, floribus floſculoſis. *Sp. pl. p.* 1288. *Fl. dan. t.* 996.

Iacea foliis radicalibus ſemipinnatis, caulinis ovato - lanceolatis. *Hall. hiſt. n.* 184.

Iacea auſtriaca ſexta. *Cluſ. hiſt. p. VII.*

In Auſtriæ, Pannoniæ pratis, ad ſepes. Fl. Iulio, Auguſto. Per.

montana. 3. C. calycibus ſerratis, foliis lanceolatis decurrentibus, caule ſimpliciſſimo. *Sp. pl. p.* 1289. *Iacqu. auſtr. t.* 371. *Scop. carn. tom.* 2. *p.* 137.

Cyanus caule unifloro, alato, foliis ellipticis lanceolatis. *Hall. hiſt. n.* 190.

In collibus, montanis, & alpinis. Fl. a Iunio ad Auguſtum. Per.

Folia ſunt lanceolata, acuta, ſeſſilia, decurrentia, utrinque hirſuta, denticulata vel integerrima, indiviſa vel lyrato - angulata.

Cyanus. 4. C. calycibus ſerratis, foliis linearibus integerrimis: infimis dentatis. *Sp. pl. p.* 1289. *Fl. dan. t.* 993. *Scop. carn. tom.* 2. *p.* 138.

Cyanus foliis imis ellipticis, dentatis, ſuperioribus linearibus, ſemifloſculis latiſſimis. *Hall. hiſt. n.* 191.

Le Bluet. Regn. bot.

Inter ſegetes. Fl. æſtate.

paniculata. 5. C. calycibus cyliatis: ſquamis planis, foliis bipinnatis: rameis pinnatifidis linearibus, caule paniculato. *Sp. pl. p.* 1289. *Iacqu. auſtr. t.* 320. *Scop. carn. tom.* 2. *p.* 139.

Cyanus pinnulis foliorum pinnatis, linearibus. *Hall. hiſt. n.* 187.

Stœbe gallica, & auſtriaca elatior. *Clus. hiſt. p. X.*

In muris, arvis, ad vias. Fl. a Iulio ad Octobrim.

Scabioſa. 6. C. calycibus ciliatis, foliis pinnatifidis: pinnis lanceolatis. *Sp. pl. p.* 1291. *Scop. carn. tom.* 2. *t.* 139.

Cyanus foliis pinnatis, pinnis dentatis, obtuſis, ſquamis calycinis triangularibus. *Hall. hiſt. n.* 186.

In pratis, ruderatis, & alibi. Fl. æſtate. Per.

Iacea. 7. C. calycibus ſcarioſis laceris, foliis lanceolatis: radicalibus ſinuato - dentatis, ramis angulatis. *Sp. pl. p.* 1293. *Fl. dan. t.* 519.

Rhaponticum foliis tomentoſis, radicalibus ſemipinnatis, caulinis lanceolatis. *Hall. hiſt. n.* 195.

Rhaponticum Iacea. *Scop. carn. tom.* 2. *p.* 135.

La Centaurée Iacée. Bulliard herb. t. 227.

In pratis. Fl. æſtate. Per.

ſplendens. 8. C. calycibus ſcarioſis obtuſis, foliis radicalibus bipinnatifidis; caulinis pinnatis: dentibus lanceolatis. *Sp. pl. p.* 1293.

Rhaponticum foliis pinnatis, pinnis ſemipinnatis, pinnulis lanceolatis, calycibus nitidis. *Hall. hiſt. n.* 197.

Stœbe ſalmantica tertia. *Clus. hiſt. p. X.*

In aſperis agri fluminenſis. Fl. Iunio, Iulio.

Rhapontica. 9. C. calycibus ſcarioſis, foliis ovato-oblongis denticulatis integris petiolatis: ſubtus tomentoſis. *Sp. pl. p.* 1294. *Wulfen in Iacqu. collect. tom.* 1. *p.* 223.

Centaurium foliis petiolatis, lanceolatis ad pediculum emarginatis. *Hall. hiſt. n.* 160.

Iacea alpina, capite maximo. *Segu. ver.* 2. *p.* 146.

In Carniolia in monte *Grindovitz* verſus torrentem, *Kokra* dictum; & in alpibus wochinenſibus in monte, *Per Ieſerh* dicto. *Zois.* Fl. Iulio, Auguſto. Per.

Calcitrapa. 10. C. calycibus ſubduplicato-ſpinoſis ſeſſilibus, foliis pinnatifidis linearibus dentatis, caule piloſo. *Sp. pl. p.* 1297.

Calcitrapa foliis pinnatis, pinnis dentatis, floribus in alis ſeſſilibus. *Hall. hiſt. n.* 194.

Rhaponticum Calcitrapa. *Scop. carn. tom.* 2. *p.* 136.

Carduus ſtellatus foliis Papaveris erratici. *Bauh. pin.* 387. *Zann. iſt. p.* 76. *t.* 152.

Le Chardon etoilé. Regn. bot.

In Litoralis, Forijulii, Carniolia calidioris, agri vindobonenſis aſperis, ad vias. Fl. Iunio, Iulio.

ſolſtitia-lis. 11. C. calycibus duplicato-ſpinoſis ſolitariis, foliis rameis decurrentibus inermibus lanceolatis, radicalibus lyrato-pinnatifidis. *Sp. pl. p.* 1297. *Scop. carn. tom.* 2. *p.* 140.

Calcitrapa caule alato, foliis tomentoſis, radicalibus ſemipinnatis, caulinis lanceolatis, dentatis. *Hall. hiſt. n.* 193.

In apricis aſperis, ad vias. Fl. Iulio, Auguſto.

collina. 12. C. calycibus ciliatis inermi-ſpinoſis, foliis bipinnatifidis, caule angulato. *Sp. pl. p.* 1298. *Scop. carn. tom.* 2. *p.* 141.

In aſperis Litoralis. Fl. æſtate. Per.

POLYGAMIA NECESSARIA.

CALENDULA. Receptaculum nudum. Pappus nullus. Calyx polyphyllus, æqualis. Semina hermaphroditis centralia disci nulla: peripheriæ propria rarius ſolitaria, membranacea. Femineis ſolitaria, incurva.

arvenſis. 1. C. ſeminibus cymbiformibus muricatis incurvatis: extimis erectis protenſis. *Sp. pl. p.* 1303.

Calendula foliis lanceolatis amplexicaulibus, dentatis, floribus concoloribus. *Hall. hiſt. n.* 94.

Le Souci. Regn. bot.

In arvis, vineis Forijulii, Litoralis. Fl. æſtate.

FILAGO. Receptaculum nudum. Pappus brevis, ſimplex. Calyx imbricatus. Floſculi inter ſquamas calycis locati.

germanica. 1. F. panicula dichotoma, floribus rotundatis axillaribus hirſutis, foliis acutis. *Sp. pl. p.* 1311. *Fl. dan. t.* 997.

Filago caule dichotomo, corymbo ſphærico, in angulo ſeſſili. *Hall. hiſt. n.* 153.

In Styria ſuperiore prope *Leoben* in aſcenſu montis *Windisberg.* Fl. æſtate.

arvenſis. 2. F. caule paniculato, floribus conicis lateralibus. *Sp. pl. p.* 1312.

Filago caule erecto, ramoſiſſimo, foliis molliſſimis, tomentoſis, florum acervis ſeſſilibus, in ſpicas continuatis. *Hall. hiſt. n.* 154.

Gnaphalium arvenſe. *Scop. carn. tom.* 2. *p.* 153.

In arvis. Fl. Iulio, Auguſto.

Leontopodium. 3. F. caule ſimpliciſſimo, capitulo terminali bracteis hirſutiſſimis radiato. *Sp. pl. p.* 1312.

Filago floribus ſeſſilibus, umbellatis, foliis umbellam excedentibus, tomentoſis. *Hall. hiſt. n.* 152.

Gnaphalium Leontopodium. *Iacqu. auſtr. t.* 86. *Scop. carn. tom.* 2. *p.* 150.

Gnaphalium alpinum. *Clus. hiſt. p.* 328.

In ſummis alpium jugis. Fl. Iulio, Auguſto. Per.

POLYGAMIA SEGREGATA.

ECHINOPS. Calyx uniflorus. Corollæ tubuloſæ hermaphroditæ. Receptaculum ſetoſum. Pappus obſoletus.

ſphærocephalus. 1. E. capitulis globoſis, foliis ſinuatis pubeſcentibus. *Sp. pl. p.* 1314. *Miller illuſtr. ic. Scop. carn. tom.* 2. *p.* 121.

Echinops caule ramoſo, foliis ſemipinnatis, pinnis dentatis, dentibus ariſtatis. *Hall. hiſt. n.* 158.

In uliginoſis, ad fluvios, foſſas. Fl. æſtate. Per.

Ritro. 2. E. capitulo globoſo, foliis pinnatifidis ſupra glabris. *Sp. pl. p.* 1314. *Scop. carn. tom.* 2. *p.* 121.

In aſperis Litoralis. Fl. Iunio, Iulio. Per.

MONOGAMIA.

IASIONE. Calyx communis polyphyllus. Corolla pentapetala, regularis. Capſula infera, bilocularis.

montana. 1. Iaſione. *Sp. pl. p.* 1317. *Curtis lond. Fl. dan. t.* 319. *Iacqu. en. p.* 289.

Rapunculus foliis linearibus ſubaſperis, ſpica planiuscula, petalis liberis. *Hall. hiſt. n.* 678.

In Auſtriæ collibus herbidis, montanis ſylvaticis. Fl. æſtate.

VIOLA. Calyx pentaphyllus. Corolla pentapetala, irregularis poſtice cornuta. Capſula ſupera, trivalvis, unilocularis.

pinnata. 1. V. acaulis, foliis pinnatifidis. *Sp. pl. p.* 1323. *Wulfen in Iacqu. collect. tom.* 1. *p.* 314.

Viola acaulis, foliis multifidis obtuſis. *Hall. hiſt. n.* 561.

Viola acaulis foliis pinnato-multifidis, & laciniatis. *Allion. miscell. taur. tom.* 3. *p.* 181. *t.* 5. *f.* 2. *tom.* 5. *p.* 84.

In Tyroli in rupeſtribus alpium leontinarum. *Wulfen.*

hirta. 2. V. acaulis, foliis cordatis piloſo-hiſpidis. *Sp. pl. p.* 1324. *Curtis lond. Fl. dan. t.* 618. *Hall. hiſt. n.* 559.

In pratis, ſylvaticis, ad ſepes. Fl. primo vere. Per.

odorata. 3. V. foliis cordatis, ſtolonibus reptantibus. *Sp. pl. p.* 1324. *Curtis lond. Fl. dan. t.* 309. *Scop. carn. tom.* 2. *p.* 181. *Hall. hiſt. n.* 558.

La Violette odorante. Bulliard herb. t. 169.

In pratis, ad ſepes. Fl. primo vere. Per.

canina. 4. V. caule adultiore adſcendente, foliis oblongo-cordatis. *Sp. pl. p.* 1324. *Curtis lond. Scop. carn. tom.* 2. *p.* 182. *Hall. hiſt. n.* 562.

In ſylvaticis, ad vias. Fl. Majo, Iunio. Per.

montana. 5. V. caulibus erectis, foliis cordatis oblongis. *Sp. pl. p.* 1325.

In pratis humidis, montanis Auſtriæ, Pannoniæ, copioſe in inſulis danubialibus. Fl. Iunio. Per.

mirabilis. 6. V. caule triquetro, foliis reniformi-cordatis; floribus caulinis apetalis. *Sp. pl. p.* 1326. *Iacqu. auſtr. t.* 19.

In montanis ſylvaticis. Per.

Folia ſunt ob baſeos lobos plus vel minus conniventes plerumque cucullata. Flores radicales habent petala, & ſunt ſteriles. Caulini ſunt plerumque apetaloidei, & ſemper fertiles. Fertilium ſtylus brevior verſus germen circulari incurvatione redit. Sterilium ſtylus ad egreſſum de germine anguſtatus inflectitur, ut ſæpe emarcidus in inflexionis loco videatur. Flores ſteriles initio Aprilis, fertiles usque ad finem Maji videndi ſunt. Iacqu. l. c.

biflora. 7. V. caule bifloro, foliis reniformibus ſerratis. *Sp. pl. p.* 1326. *Fl. dan. t.* 46. *Scop. carn. tom.* 2. *p.* 182.

Viola caule debili, foliis petiolatis reniformibus, obtuſe dentatis. *Hall. hiſt. n.* 563.

Viola montana prima. *Clus. hiſt. p.* 309.

In alpinis, & ſubalpinis. Fl. æſtate. Per.

tricolor. 8. V. caule triquetro diffuſo, foliis oblongis inciſis, ſtipulis pinnatifidis. *Sp. pl. p.* 1326. *Curtis lond. Fl. dan. t.* 623. *Scop. carn. tom.* 2. *p.* 183.

Viola caule anguloſo, diffuſo, foliis ovatis, dentatis, flore calyce duplo longiore. *Hall. hiſt. n.* 568.

Iacea ſeu flos Trinitatis. *Cam. epit. p.* 912.

In arivis, collibus. Fl. aeſtate.

grandiflora. 9. V. caule triquetro ſimplici, foliis oblongiuſculis, ſtipulis pinnatifidis. *Syſt. veget. p.* 803.

Viola caule erecto, paucifloro, foliis imis ſubrotundis, caulinis ellipticis, ſtipulis ſemipinnatis maximis. *Hall. hiſt. n.* 566. *t.* 17.

Viola alpina. *Iacqu. auſtr. t.* 242.

Viola Zoiſii. *Wulfen in Iacqu. collect. tom.* 4. *p.* 297. *t.* 11. *f.* 1.

Viola montana ſecunda. *Clus. hiſt. p.* 309.

In alpibus. Fl. Iulio, Auguſto. Per.

Variat floris colore flavo, cœruleo, albo.

IMPATIENS. Calyx diphyllus. Corolla pentapetala, irregularis, nectario cucullato. Capſula ſupera, quinquevalvis.

noli tangere. 1. I. pedunculis multifloris ſolitariis, foliis ovatis, geniculis caulinis tumentibus. *Sp. pl. p.* 1329. *Fl. dan. t.* 588.

Impatiens caule geniculato, pedunculis ramoſis, foliis petiolatis, ovatis, crenatis *Hall. hiſt. n.* 557.

Balſamina noli tangere. *Scop. carn. tom.* 2. *p.* 184.

In nemorenſibus, udis. Fl. aeſtate.

GYNANDRIA.

DIANDRIA.

ORCHIS. Nectarium corniforme pone florem.

biſolia. 1. O. bulbis indiviſis, nectarii labio lanceolato integerrimo: cornu longiſſimo, petalis patentibus. *Sp. pl. p.* 1331. *Fl. dan. t.* 235. *Scop. carn. tom.* 2. *p.* 186.

Orchis radicibus oblongis, labello lineari. *Hall. hiſt. n.* 1285. *t.* 35.

In ſylvis, pratis ſicc[i]oribus. Fl. Iulio, Auguſto. Per.

Columna ſtaminea nulla; ſed ſtigma maximum fornicatum, cujus latera fiſſa includunt partes maſculas. Stamina omnino libera, ſed in rima piſtilli laterali abſcondita, filamento anthera longiore, pelta laterali prope baſin viscida inſtructo. Retzii obs. faſc. 1. p. 30.

ornithis. 2. O. bulbis indiviſis, nectarii labio ſubrotundo, cornu germine duplo longiori, petalis tribus exterioribus conniventibus, reliquis patentiſſimis. *Syſt. veget. p.* 808. *Iacqu. auſtr. t.* 138.

In alpibus auſtriacis. *Iacqu.* Fl. Auguſto. Per.

globoſa. 3. O. bulbis indiviſis, nectarii labio reſupinato trifido: media emarginata: cornu brevi, petalis apice ſubulatis. *Syſt. veget. p.* 808. *Iacqu. auſtr. t.* 265. *Scop. carn. tom.* 2. *p.* 188.

Orchis radicibus ſubrotundis, ſpica denſiſſima, petalis exterioribus ariſtatis. *Hall. hiſt. n.* 1272. *t.* 27.

In pratis alpinis, & ſubalpinis. Fl. Iulio, Auguſto. Per.

pyramidalis. 4. O. bulbis indiviſis, nectarii labio bicorni trifido æquali integerrimo: cornu longo, petalis ſublanceolatis. *Syſt. veget. p.* 808. *Iacqu. auſtr. t.* 266. *Scop. carn. tom.* 2. *p.* 187.

Orchis radicibus ſubrotundis, ſpica denſa, calcare longiſſimo, labello bicorni, æqualiter tripartito. *Hall. hiſt. n.* 1286. *t.* 35.

In pratis, paſcuis montanis, & ſubalpinis. Fl. Iulio, Auguſto. Per.

coriophora. 5. O. bulbis indiviſis, nectarii labio trifido reflexo crenato: cornu brevi, petalis conniventibus. *Sp. pl. p.* 1332. *Iacqu. auſtr. t.* 122. *Scop. carn. tom.* 2. *p.* 189.

Orchis radicibus ſubrotundis, galea connivente, labello trifido reflexo. *Hall. hiſt. n.* 1284. *t.* 34.

Orchis odore hirci minor. *Bauh. pin.* 82.

In pratis. Fl. Iunio. Per.

paluſtris. 6. O. bulbis indiviſis, nectarii labio trifido: media emarginata, petalis erectis, duobus interioribus conniventibus. *Iacqu. collect. tom.* 1. *p.* 75. *ic. t.* 181.

Orchis maſcula. *Crantz ſtirp. auſtr. p.* 500.

In Auſtria in pratis paluſtribus prope *Erla, Himberg, Lachſenburg.* In Pannonia ad lacum Peiſonis. Fl. Iunio. Per.

Bulbi indiviſi. Caulis folioſus, teres. Folia acuta, inferne vaginantia, dein conniventia. Spathæ germini æquales. Flos inodorus, ſaturate purpureus, vel carneus. Petala erecta, oblonga, ovata, obtuſa: duobus interioribus conniventibus. Nectarii cornu obtuſum, germine paullo brevius. Labii ampli & trifidi laciniæ laterales rotundatæ: intermedia minima & emarginata. Iacqu. l. c.

Morio. 7. O. bulbis indiviſis, nectarii labio quadrifido crenulato: cornu obtuſo, petalis omnibus conniventibus. *Sp. pl. p.* 1333. *Fl. dan. t.* 253. *Curtis lond. Scop. carn. tom.* 2. *p.* 190.

Orchis radicibus ſubrotundis, galeæ petalis lineatis, labello trifido, crenato, medio ſegmento emarginato. *Hall. hiſt. n.* 1282. *t.* 33.

In pratis, paſcuis. Fl. Majo, Iunio. Per.

Flores variant labio nectarii inflexo & plano, quadrifido, indiviſo, & leviter tantum emarginato; cornu præterea obtuſo & gibbo, cylindrico, compreſſo, vel ſubcompreſſo. Retzii obs. faſc. 1. p. 30.

aſcula. 8. O. bulbis indiviſis, nectarii labio quadrilobo crenulato: cornu obtuſo, petalis dorſalibus reflexis. *Sp. pl. p.* 1333. *Iacqu. miſcell. tom.* 2. *p.* 375. *ic. t.* 180. *Curtis lond. Fl. dan. t.* 457. *Scop. carn. tom.* 2. *p.* 189.

Orchis radicibus ſubrotundis, petalis lateralibus reflexis, labello trifido, ſegmento medio longiori, bifido. *Hall. hiſt. n.* 1283. *t.* 33.

In pratis, paſcuis ſubalpinis, & alpinis. Fl. Iulio. Per.

ſtulata. 9. O. bulbis indiviſis, nectarii labio quadrifido punctis ſcabro: cornu obtuſo, petalis diſtinctis. *Sp. pl. p.* 1333. *Scop. carn. tom.* 2. *p.* 189.

Orchis radicibus ſubrotundis, labello quadrifido, calcare breviſſimo. *Hall. hiſt. n.* 1273. *t.* 28.

Orchis anthropophora. *Fl. dan. t.* 103.

Orchis pannonica quarta. *Clus. hiſt. p.* 268.

In pratis. Fl. Majo, Iunio. Per.

ariegata. 10. O. bulbis indiviſis, nectarii labio quadrilobo, cum denticulo medio: lobis ſerratis, diſco nudo glabro. *Iacqu. collect. tom.* 2. *p.* 267. *ic. t.* 599.

Ochis radicibus ſubrotundis, ſpica breviſſima, labello breviter quadrifido, circumſerrato, punctato. *Hall. hiſt. n.* 1275.

Orchis militaris pratenſis elatior, floribus variegatis. *Segu. ver.* 2. *p.* 123. *t.* 15. *f.* 3.

In Auſtriæ, Pannoniæ pratis Fl. Majo, Iunio. Per.

Petala in cucullum erectum conniventia, pallide purpurea: tribus exterioribus duplo longioribus & lanceolato-acuminatis; duobus interioribus lanceolato-linearibus & acutiuſculis. Hæc

& petalum exterius dorſale linea unica, intus *atropurpurea*, *foris minus ſaturata*, percurrit, *dum tales duæ percurrunt petala exteriora* duo *lateralia. Nectarii cornu ſubulatum*, *ſtriatum*, *germine brevius. Labium inferius petalis* concolor, *& totum adſperſum punctis purpureis*, quadrifidum: *lobis ad apicem ſerratis*; *lateralibus oblongis*, *intermediis rotundatis*, *cum denticulo inter utrumque. Discus glaber.* Iacqu. l. c.

militaris. 11. O. bulbis indiviſis, nectarii labio quinquefido punctis ſcabro: cornu obtuſo, petalis confluentibus. *Sp. pl. p.* 1333. *Iacqu. collect. tom.* 2. *p.* 268. *ic. t.* 598.

Orchis radicibus ſubrotundis, ſpica conica, labello quadrifido; brachiolis & pectore perangustis. *Hall. hiſt. n.* 1277. *t.* 28.

Habitat & floret cum præc. Per.

Petala foris purpureo - cinerea, *intus purpura lineisque ſaturatioribus variegata*, *acuta. Tria exteriora ſunt lanceolata*; *duo interiora lanceolato - linearia*, *prioribus paullo breviora. Nectarii cornu cinereum*, *breve*, *obtuſum compreſſulum. Labium quadrilobum cum denticulo medio*: *lobis integerrimis*, *obtuſis*, *& ſaturate purpureis. Discus pallidus*, *muricatus piliſque faſciculatis purpureis tectus. Uti variat colore floris magis minusve purpureo*, *ſic variat quoque albido.* Iacqu. l. c.

fusca. 12. O. bulbis indiviſis, nectarii labio quadrifido: laciniis oblongis, cornu brevi, petalis confluentibus. *Syſt. veget. p.* 809. *Iacqu. auſtr. t.* 307.

Orchis radicibus ſubrotundis, ſpica longa, labello quadrifido, brachiolis anguſtis, crusculis latis, ſerratis. *Hall. hiſt. n.* 1276. *t.* 31.

Orchis militaris. *Scop. carn. tom.* 2. *p.* 192.

In montium declivibus, in pratis, ſylvis. Fl. Majo, Iunio. Per.

pallens. 13. O. bulbis indiviſis, nectarii labio trifido integerrimo, cornu obtuſo mediocri, petalis patentibus. *Syſt. veget. p.* 809. *Iacqu. auſtr. t.* 45.

Orchis radicibus ſubrotundis, petalis galeæ lineatis, labello trifido, integerrimo, *Hall. hiſt. n.* 1281.

Orchis pannonica octava. *Cluſ. hiſt. p.* 269.

In collibus Auſtriæ herbidis, montoſis ſylvaticis. Fl. Aprili, Majo. Per.

ſambucina. 14. O. bulbis ſubpalmatis erectis, nectarii cornu conico: labio ovato ſubtrilobo, bracteis longitudine florum. *Syſt. veget. p.* 810. *Iacqu. auſtr. t.* 108.

Orchis radicibus palmatis, bracteis coloratis, labello circumſerrato, trilobato, lobo medio emarginato. *Hall. hiſt. n.* 1280.

Orchis *mixta*, bulbis palmatis: nectarii cornu conico: labio trilobo ſubſerrato, petalis dorſalibus reflexis. *Retzii ſcand. p.* 205.

Orchis incarnata. *Sp. pl. p.* 1335.

Orchis latifolia. *Scop. carn. tom.* 2. *p.* 196.

Orchis palmata lutea labio floris maculato. *Segu. ver. ſuppl. p.* 248. *t.* 8. *f.* 5.

Ad margines ſylvarum aſperos, & herbidos, in pratis Auſtriæ, Croatiæ, Carnioliæ. Fl. Majo, Iunio. Per.

Bulbus palmatus plerumque bifidus, raro trifidus, idque obſolete. Invenitur tamen in adulta ſtirpe uterque integerrimus & ovatus. In ſambucina folia dilutius virent, ſunt vere caulina, perque intervalla diſpoſita. In pallente ſunt ſubradicalia, horum vaginæ confluunt. Flores O. pallentis florum ſambuci nigræ odorem redolent; bracteæ ſunt multo anguſtiores & breviores; nectarii labellum perpetuo immaculatum eſt; & cornu manifeſte emarginatum retrorſum adſcendit. In ſambucina flores inodori, nectarii labellum purpureis punctis fere ſemper variegatum, & cornu vix emarginatum tendit deorſum. Iacqu. l. c.

Color florum albidus. In Carniolia, Croatia invenitur varietas floribus rubris, O. incarnatam Linn. ſiſtens.

latifolia. 15. O. bulbis ſubpalmatis rectis, nectarii cornu conico: labio trilobo lateribus reflexo, bracteis flore longioribus. *Sp. pl. p.* 1334. *Fl. dan. t.* 266. *Curtis lond.*

Orchis radicibus palmatis, caule fiſtuloſo, bracteis maximis, labello trifido, ſerrato, medio ſegmento obtuſo. *Hall. hiſt. n.* 1279. *t.* 32.

Orchis comoſa. *Scop. carn. tom.* 2. *p.* 198.

In pratis udis. Fl. Iunio. Per.

maculata. 16. O. bulbis palmatis patentibus, nectarii cornu germinibus breviore: labio plano, petalis dorſalibus patulis. *Sp. pl. p.* 1335. *Fl. dan. t.* 933. *Scop. carn. tom.* 2. *p.* 198.

Orchis radicibus palmatis, caule ſolido, labello trifido, ſerrato, medio ſegmento minore, ovato. *Hall. hiſt. n.* 1278. *t.* 32.

In ſylvaticis, pratis montanis, & ſubalpinis. Fl. Iunio, Iulio. Per.

odoratiſſima. 17. O. bulbis palmatis, nectarii cornu recurvo, labio ovato acuto, foliis linearibus. *Sp. pl. p.* 1335. *Iacqu. auſtr. t.* 264.

Orchis radicibus palmatis, labello obtuſe trifido, concolore, calcare germine breviori. *Hall. hiſt. n.* 1274. *t.* 29.

Orchis palmata anguſtifolia minor odoratiſſima. *Bauh. pin.* 86. *prodr.* 30.

In Auſtriæ, Styriæ pratis ſubalpinis. Fl. Iulio. Per.

conopſea. 18. O. bulbis palmatis, nectarii cornu ſetaceo germinibus longiore: labio trifido, petalis duobus patentiſſimis. *Sp. pl. p.* 1335. *Fl. dan. t.* 224. *Scop. carn. tom.* 2. *p.* 199.

Orchis radicibus palmatis, calcare longiſſimo, labello unicolore, obtuſe trifido. *Hall. hiſt. n.* 1287. *t.* 29.

In paſcuis, pratis montanis. Fl. Iulio. Per.

abortiva. 19. O. bulbis fasciculatis filiformibus, nectarii labio ovato integerrimo. *Sp. pl. p.* 1336. *Iacqu. auſtr. t.* 193.

Epipactis aphylla, calcare longo, labello ovato lanceolato. *Hall. hiſt. n.* 1288. *t.* 36.

Serapias abortiva. *Scop. carn. tom.* 2. *p.* 205.

Pſeudolimodorum. *Clus. hiſt. p.* 270.

In montoſis ſylvaticis Auſtriæ, Pannoniæ, Litoralis, Fl. Iunio. Per.

SATYRIUM. Nectarium inflato-didymum pone florem.

hircinum. 1. S. bulbis indiviſis, foliis lanceolatis, nectarii labio trifido: intermedia lineari elongata obliqua præmorſa. *Sp. pl. p.* 1337. *Iacqu. auſtr. t.* 367.

Orchis radicibus ſubrotundis, labello longiſſimo, tripartito, plicato. *Hall. hiſt. n.* 1268. *t.* 25.

Orchis hircina. *Scop. carn. tom.* 2. *p.* 193.

Orchis barbata, odore hirci, breviore latioreque folio. *Bauh. pin.* 82.

In Auſtriæ, Pannoniæ, Litoralis collibus herbidis, paſcuis, montanis ſylvaticis. Fl. Iunio, Iulio. Per.

viride. 2. S. bulbis palmatis, foliis oblongis obtuſis: nectarii labio lineari trifido: intermedia obſoleta. *Sp. pl. p.* 1337.

Orchis radicibus palmatis, galea connivente, labello triſulco, calcare breviſſimo. *Hall. hiſt. n.* 1269. *t.* 26.

Orchis virens. *Scop. carn. tom.* 2. *p.* 199.

Orchis palmata, flore viridi. *Bauh. pin.* 86. *prodr.* 30.

In pratis, paſcuis montanis, ſubalpinis, & alpinis. Fl. Iunio, Iulio. Per.

nigrum. 3. S. bulbis palmatis, foliis linearibus, floribus reſupinatis, nectarii labio indiviſo ovato acuminato. *Sp. pl. p.* 1338. *Iacqu. auſtr. t.* 368. *Fl. dan. t.* 998.

Orchis radicibus palmatis, ſpica denſiſſima, flore reſupinato, calcare breviſſimo. *Hall. hiſt. n.* 1271. *t.* 27.

Orchis nigra. *Scop. carn. tom.* 2. *p.* 200.

Orchis palmata anguſtifolia alpina, nigro flore. *Bauh. pin.* 86.

In alpinis, & ſubalpinis. Fl. Iulio, Auguſto. Per.

albidum. 4. S. bulbis faſciculatis, foliis lanceolatis, nectarii labio trifido acuto: lacinia intermedia obtuſa. *Sp. pl. p.* 1338. *Fl. dan. t.* 115.

Orchis radicibus confertis, teretibus, calcare breviſſimo, labello trifido. *Hall. hiſt. n.* 1270. *t.* 26.

Orchis albida. *Scop. carn. tom.* 2. *p.* 201.

Habitat & floret cum præc. Per.

Epipogium. 5. S. bulbis compreſſis dentatis, caule vaginato, corollis reſupinatis, nectarii labio indiviſo. *Sp. pl. p.* 1338. *Iacqu. auſtr. t.* 84.

Epipactis caule aphyllo, flore reſupinato, labello ovato lanceolato, calcare ovato turgido. *Hall. hiſt. n.* 1289.

In ſylvis Auſtriæ, Carnioliæ umbroſis ſubalpinis. Fl. Iulio, Auguſto. Per.

repens. 6. S. bulbis fibroſis, foliis ovatis radicalibus, floribus ſecundis. *Sp. pl. p.* 1339. *Iacqu. auſtr. t.* 369.

Epipactis foliis petiolatis, ovato-lanceolatis, floribus tetrapetalis hirſutis. *Hall. hiſt. n.* 1295. *t.* 22.

In ſylvis montanis, & ſubalpinis. Fl. Iulio, Auguſto. Per.

OPHRYS. Nectarium ſubtus ſubcarinatum.

Nidus avis. 1. O. bulbis fibroſo-faſciculatis, caule vaginato aphyllo, nectarii labio bifido. *Syſt. veget. p.* 812. *Fl. dan. t.* 181. *Scop. carn. tom.* 2. *p.* 206.

Epipactis aphylla, flore inermi, labello bicorni. *Hall. hiſt. n.* 1290. *t.* 37.

In ſylvis. Fl. Iunio. Per.

Corallorhiza. 2. O. bulbis ramoſis flexuoſis, caule vaginato aphyllo, nectarii labio trifido. *Syſt. veget. p.* 812. *Fl. dan. t.* 451. *Hall. hiſt. n.* 1301. *t.* 44.

Corallorhiza Neotia. *Scop. carn. tom.* 2. *p.* 207.

Dentaria coralloide radice, ſive aphyllos tertia. *Cluſ. hiſt. p. CXX.*

In ſylvis ſubalpinis. Fl. Iulio, Auguſto. Per.

ſpiralis. 3. O. bulbis aggregatis oblongis, caule ſubfolioſo, floribus ſpirali-ſecundis, nectarii labio indiviſo crenulato. *Syſt. veget. p.* 812. *Fl. dan. t.* 387.

Epipactis bulbis cylindricis, ſpica ſpirali, labello crenulato. *Hall. hiſt. n.* 1294.

Serapias ſpiralis. *Scop. carn. tom.* 2. *p.* 201.

In ſylvis montanis, & ſubalpinis. Fl. Iulio, Auguſto. Per.

ovata. 4. O. bulbo fibroſo, caule bifolio, foliis ovatis, nectarii labio bifido. *Sp. pl. p.* 1340. *Fl. dan. t.* 137. *Curtis lond. Scop. carn. tom.* 2. *p.* 206.

Epipactis foliis binis ovatis, labello bifido. *Hall. hiſt. n.* 1291. *t.* 37.

In pratis humidis. Fl. Majo, Iunio, Iulio. Per.

cordata. 5. O. bulbo fibroſo, caule bifolio, foliis cordatis. *Sp. pl. p.* 1340. *Iacqu. collect. tom.* 2. p. 265. *Scop. carn. tom.* 2. *p.* 207.

Epipactis foliis binis, cordatis, labello bifido, poſtice bidentato. *Hall. hiſt. n.* 1292. *t.* 22. *f.* 4.

In ſylvis ſubalpinis. Fl. Iulio, Auguſto. Per.

Lœſelii. 6. O. bulbo ſubrotundo: ſcapo nudo trigono, nectarii labio ovato. *Sp. pl. p.* 1341. *Hænke in Iacqu. colleᴐ. tom.* 2. *p.* 53.

Ophrys paludoſa. *Fl. dan. t.* 877.

Ophrys diphyllos bulboſa. *Lœs. pruſs.* 180. *t.* 58.

In Pannonia in paluſtribus ad lacum Peiſonis. *Hænke.* Fl. Iunio. Per.

monophyllos. 7. O. bulbo rotundo, ſcapo nudo, folio ovato, nectarii labio integro. *Sp. pl. p.* 1342. *Wulfen in Iacqu. collect. tom.* 4. *p.* 340. *t.* 13. *f.* 2.

Epipactis folio unico, amplexicauli, ſpica prolixa multiflora. *Hall. hiſt. n.* 1293. *t.* 36.

In Carnioliæ montanis.

myodes. 8. O. bulbis ſubrotundis, ſcapo folioſo, nectarii labio oblongo, quadripartito, laciniis duabus mediis elongatis lanceolatis. *Syſt. veget. p.* 813. *Iacqu. miſcell. tom.* 2. *p.* 373. *ic. t.* 164.

Orchis radicibus ſubrotundis, labello ſericeo biſulco, brachiolis brevibus ſubulatis. *Hall. hiſt. n.* 1265.

Ophrys inſectifera var. myodes. *Sp. pl. p.* 1343.

Orchis muſcaria. *Scop. carn. tom.* 2. *p.* 193.

Orchis muſcæ corpus referens, ſive galea & alis herbidis. *Bauh. pin.* 83. *Vaill. paris.* 147. *t.* 31. *f.* 18.

In Auſtriæ ſubalpinis. In Litoralis, Forijulii vineis, pratis. Fl. Iunio, Iulio. Per.

arachnites. 9. O. bulbis ſubrotundis, ſcapo folioſo, nectarii labio lato emarginato appendiculato. *Syſt. veget. p.* 813. *Curtis lond.*

Orchis radicibus ſubrotundis, labello holoſericeo emarginato, appendiculato. *Hall. hiſt. n.* 1266. *t.* 24.

Ophrys inſectifera var. arachnites. *Sp. pl. p.* 1343.

Orchis arachnites. *Scop. carn. tom.* 2. *p.* 194.

Orchis araneam referens. *Bauh. pin.* 84. *Vaill. paris. t.* 30. *f.* 10, 11, 12, 13.

In pratis. Fl. Majo, Iunio. Per.

Monorchis. 10. O. bulbo globoſo, ſcapo nudo, nectarii labio trifido cruciato. *Sp. pl. p.* 1342. *Fl. dan. t.* 102.

Orchis bulbo unico, ſubrotundo, labello cruciformi. *Hall. hiſt. n.* 1262. *t.* 22.

Orchis Mnorchis. *Scop. carn. tom.* 2. *p.* 195.

In pratis, paſcuis ſubalpinis, montanis. Fl. Iunio, Iulio. Per.

alpina. 11. O. bulbis ovatis, ſcapo nudo, foliis ſubulatis, nectarii labio indiviſo obtuſo utrinque unidentato. *Sp. pl. p.* 1342. *Fl. dan. t.* 452. *Iacqu. en. p.* 295. *t.* 9.

Orchis radicibus ſubrotundis, labello ovato, utrinque denticulo notato. *Hall. hiſt. n.* 1263. *t.* 22.

Orchis alpina. *Scop. carn. tom.* 2. *p.* 196.

Chamæorchis alpina gramineo folio. *Bauh. pin.* 81. *prodr.* 29.

In alpibus. Fl. a Iulio ad Septembrim. Per.

SERAPIAS. Nectarium ovatum, gibbum, labio ovato.

latifolia. 1. S. bulbis fibroſis, foliis ovatis amplexicaulibus, floribus pendulis. *Syſt. veget. p.* 814. *Fl. dan. t.* 811. *Scop. carn. tom.* 2. *p.* 203.

Epipactis foliis amplexicaulibus, ovato-lanceolatis, labello lanceolato. *Hall. hiſt. n.* 1297. *t.* 40.

Helleborine latifolia montana. *Bauh. pin.* 186.

In collibus herbidis, montanis ſylvaticis, & ſubalpinis. Fl. Iunio, Iulio. Per.

longifolia. 2. S. bulbis fibroſis, foliis enſiformibus ſeſſilibus, floribus pendulis: labio obtuſo plicato. *Retzii ſcand. p.* 208. *Syſt. veget. p.* 815. *Fl. dan. t.* 267.

Epipactis foliis enſiformibus caulinis, floribus pendulis, labello obtuſo, oris plicatis. *Hall. hiſt. n.* 1296. *t.* 39.

Serapias paluſtris. *Scop. carn. tom.* 2. *p.* 204.

Helleborine anguſtifolia paluſtris ſeu pratenſis. *Bauh. pin.* 87.

In pratis paluſtribus. Fl. Iunio. Per.

enſifolia. 3. S. bulbis fibroſis; foliis enſiformibus, diſtichis; bracteis minutiſſimis; floribus erectis: nectarii labio obtuſo, petalis breviore. *Syſt. veget. p.* 815. *Fl. dan. t.* 506.

Serapias longifolia. *Scop. carn. tom.* 2. *p.* 202.

Serapias xyphophyllum. *Ehrh. beytr. tom.* 1. *p.* 185.

In montanis ſylvaticis. Fl. Majo, Iunio. Per.

lancifolia. 4. S. bulbis fibroſis; foliis ovato-lanceolatis; bracteis longitudine capſulæ; floribus erectis: nectarii labio obtuſo, petalis breviore. *Syſt. veget. p.* 815. *Hall. hiſt. n.* 41.

Serapias grandiflora. *Scop. carn. tom.* 2. *p.* 203.

Serapias lonchophyllum. *Ehrh. l. c.*

Habitat & floret cum præc. Per.

rubra. 5. S. bulbis fibroſis, foliis enſiformibus, floribus erectis: nectarii labio acuto. *Syſt. veget. p.* 816. *Fl. dan. t.* 345.

Epipactis caule paucifloro, lineis acuti labelli undulatis. *Hall. hiſt. n.* 1299. *t.* 42.

In collibus, montanis ſylvaticis Auſtriæ, Pannoniæ. Fl. Iunio. Per.

Lingua. 6. S. bulbis ſubrotundis, nectarii labio trifido acuminato glabro petalis longiore. *Syſt. veget. p.* 816.

Orchis lingua. *Scop. carn. tom.* 2. *p.* 187.

Orchis montana italica, flore ferrugineo; lingua oblonga. *Bauh. pin.* 84. *prodr.* 29. *Segu. ver. ſuppl. p.* 248. *t.* 8. *f.* 4.

In herbidis agri tergeſtini prope *Tomai. I. Hoſt.* Fl. Iunio. Per.

CYPRIPEDIUM. Nectarium ventricoſum, inflatum, cavum.

Calceolus. 1. C. radicibus fibroſis, foliis ovato-lanceolatis caulinis. *Sp. pl. p.* 1346. *Fl. dan. t.* 999.

Calceolus radicibus fibroſis, foliis ovato-lanceolatis. *Hall. hiſt. n.* 1300. *t.* 43.

Elleborine recentiorum prima. *Clus. hiſt. p.* 272.

In montanis ſylvaticis agri vindobonenſis. Fl. Iunio. Per.

HEXANDRIA.

ARISTOLOCHIA. Calyx nullus. Corolla monopetala, lingulata, integra. Filamenta nulla. Antheræ ſtigmatibus adnatæ. Capſula ſexlocularis, infera.

rotunda. 1. A. foliis cordatis ſubſeſſilibus obtuſis, caule infirmo, floribus ſolitariis. *Sp. pl. p.* 1364. *Scop. carn. tom.* 2. *p.* 208.

Ariſtolochia rotunda prima. *Clus. hiſt. p. LXX.*

In ſylvaticis, ad ſepes Forijulii, Litoralis. Fl. primo vere. Per.

longa. 2. A. foliis cordatis petiolatis integerrimis obtuſiuſculis, caule infirmo, floribus ſolitariis. *Sp. pl. p.* 1364. *Scop. carn. tom.* 2. *p.* 209.

Ariſtolochia longa prima. *Cluſ. hiſt. p. LXX.*

Habitat & floret cum præc. Per.

Clematitis. 3. A. foliis cordatis, caule erecto, floribus axillaribus confertis. *Sp. pl. p.* 1364. *Miller illuſtr. ic. Scop. carn. tom.* 2. *p.* 209. *Hall. hiſt. n.* 1029.

Ariſtolochia Clematitis vulgaris. *Cluſ. hiſt. p. LXXI.*

L' Ariſtoloche Clematite. Regn. bot.

In vineis, ad agrorum margines, fluviorum ripas, & alibi. Fl. Majo, Iunio. Per.

MONOECIA.

MONANDRIA.

ARUM. Spatha monophylla, cucullata. Spadix clavatus, ſimpliciſſimus, ſuperne nudus, in medio antheris inter duplicem cirrhorum ordinem; inferne germinibus obvallatus.

Masc. Calyx, & corolla propria nulla. Anthera ſingula ſeſſilis, tetragona.

Fem. Calyx, & corolla propria nulla. Germen ſingulum obovatum. Stylus nullus. Stigma villis barbatum. Bacca globoſa, unilocularis, polyſperma.

Dracunculus. 1. A. foliis pedatis: foliolis lanceolatis integerrimis æquantibus ſpatham ſpadice longiorem. *Sp. pl. p.* 1367. *Scop. carn. tom.* 2. *p.* 210.

La Serpentaire. Regn. bot.

In Carnia *al lido del Tolmeſo. I. Hoſt.* Fl. Iunio. Per.

maculatum. 2. A. acaule, foliis haſtatis integerrimis, ſpadice clavato. *Sp. pl. p.* 1370. *Fl. dan. t.* 505. *Scop. carn. tom.* 2. *p.* 210. *Curtis lond.*

Arum foliis ſagittatis; ſpatha recta: clava cylindrica. *Hall. hiſt. n.* 1302.

Le Pied de Veau. Regn. bot.

In nemoribus, ad ſepes. Fl. primo vere. Per.

CALLA. Spatha monophylla, ovato-cordata, patens. Spadix digitiformis, fructificationibus tectus.

Maſc. Calyx & corolla propria nulla. Filamentum ſingulum compreſſum, inferne anguſtius, truncatum, perſiſtens. Anthera depreſſa, bilocularis.

Fem. Calyx, corolla nulla. Germen ſingulum ſubrotundum. Stylus ſimplex. Stigma acutum. Bacca tetragono-globoſa, unilocularis, polyſperma.

paluſtris. 1. C. foliis cordatis, ſpatha plana, ſpadice undique tecto ſtaminibus & piſtillis mixtis. *Sp. pl. p.* 1373. *Fl. dan. t.* 422.

In Carinthia in limoſis ad effluxum lacus vertenſis, & oſſiacenſis; ad influxum lacus mileſtadienſis. *Wulfen.* Fl. Iunio. Per.

ZOSTERA. Spatha nulla, niſi folii baſis vaginans, longitudinaliter connivens, ſuperne utrinque emarginata, includens ſpadicem linearem, planum, altero latere ſtaminibus ſuperne, piſtillisque inferne inſtructum.

Maſc. Calyx, & corolla propria nulla. Filamenta breviſſima. Anthera ovato-oblonga, nutans.

Fem. Calyx, & corolla nulla. Germen ovatum, anceps, nutans. Stylus unicus, ſimplex. Stigmata duo capillaria. Capſula ovata, roſtrata, membranacea, diaphana, unilocularis, evalvis. Semen unicum, ſtriatum.

marina. 1. Z. pericarpiis ſeſſilibus. *Syſt. veget. p.* 838. *Fl. dan. t.* 15. *Scop. carn. tom.* 2. *p.* 211.

Alga anguſtifolia vitriariorum. *Bauh. pin.* 364.

In mari adriatico. Per.

ZANNICHELLIA. Maſc. Calyx, & corolla nulla. Anthera ſagittata.

Fem. Calyx monophyllus. Corolla nulla. Germina quatuor ad octo. Capſulæ totidem, uniloculares. Semina ſolitaria.

paluſtris. 1. Zannichellia. *Sp. pl. p.* 1375. *Fl. dan. t.* 67. *Scop. carn. tom.* 2. *p.* 212. *Hall. hiſt. n.* 1604. *Mich. gen. p.* 71. *t.* 34. *f.* 1. 2.

In aquis ſtagnantibus, vel lente fluentibus. Fl. Iulio, Auguſto. Per.

CHARA. Maſc. Calyx, & corolla nulla. Anthera germini ſubjecta.

Fem. Calyx tetraphyllus. Corolla nulla. Stigma qninquefidum.

tomento-ſa. 1. C. aculeis caulinis ovatis. *Sp. pl. p.* 1624.

Equiſetum fragile majus ſubcinereum aquis immerſum. *Moris. hiſt.* 3. *p.* 621. *ſect.* 15. *t.* 4. *f.* 9.

In aquis ſtagnantibus, mari. Fl. Iunio.

vulgaris. 2. C. caulibus lævibus, frondibus interne dentatis. *Sp. pl. p.* 1624. *Fl. dan. t.* 150. *Scop. carn. tom.* 2. *p.* 400. *Hedwig theor. crypt. t.* 32, 33.

Chara caule lævi fragili. *Hall. hiſt. n.* 1681.

In aquis ſtagnantibus. Fl. Iunio.

hiſpida. 3. C. aculeis caulinis capillaribus confertis. *Sp. pl. p.* 1624. *Fl. dan. t.* 154.

Chara caule fragili ſpinoſo. *Hall. hiſt. n.* 1682.

Tergeſti in foſſis *alle Saule.* Fl. Iunio.

DIANDRIA.

LEMNA. Maſc. Calyx monophyllus. Corolla nulla.
Fem. Cal cor. ut in mare. Styllus unicus. Capſula unilocularis.

triſulca. 1. L. foliis petiolatis lanceolatis. *Sp. pl. p.* 1376.

Hydrophace caule filiformi; foliis ellipticis, lanceolatis, viviparis. *Hall. hiſt. n.* 1901.

Lenticula triſulca. *Scop. carn. tom.* 2. *p.* 213.

Lenticularia ramoſa monorhiza, foliis oblongis, pediculis longioribus donatis. *Mich. gen. p.* 16. *t.* 11. *f.* 5.

In aquis ſtagnantibus. Fl. Auguſto.

minor. 2. L. foliis ſeſſilibus utrinque planiuſculis, radicibus ſolitariis. *Sp. pl. p.* 1376.

Hydrophace monorhiza, foliis planis ovatis. *Hall. hiſt. n.* 1900.

Lenticula minor. *Scop. carn. tom.* 2. *p.* 213.

Lenticularia minor, monorhiza, foliis ſubrotundis, utrinque viridibus. *Mich. gen. p.* 16. *t.* 11. *f.* 3.

In rivulis, & aquis ſtagnantibus. Fl. Iunio, Iulio.

polyrhiza. 3. L. foliis ſeſſilibus, radicibus confertis. *Sp. pl. p.* 1377.

Hydrophace polyrhiza, foliis ovatis planis. *Hall. hiſt. n.* 1899.

Lenticularia major polyrhiza, inferne atropurpurea. *Mich. gen. p.* 16. *t.* 11. *f.* 1.

In Auſtriæ, Pannoniæ foſſis. Fl. Iulio, Auguſto.

TRIANDRIA.

TYPHA. Masc. Amentum cylindricum. Calyx obſoletus, triphyllus. Corolla nulla.

Fem. Calyx, & corolla nulla. Germen inſidens ſetæ. Stigma capillare, perſiſtens. Semen unicum. Pappus capillaris, ſetæ ſeminiferæ a baſi ad medium affixus.

latifolia. 1. T. foliis ſubenſiformibus, ſpica maſcula femineaque approximatis. *Sp. pl. p.* 1377. *Scop. carn. tom.* 2. *p.* 214. *Fl. dan. t.* 645. *Curtis lond.*

Typha clava unica. *Hall. hiſt. n.* 1305.

In aquis ſtagnantibus, vel lente fluentibus. Fl. Iulio. Per.

anguſtifolia. 2. T. foliis ſemicylindricis, ſpica maſcula femineaque remotis. *Sp. pl. p.* 1377. *Fl. dan. t.* 815. *Curtis lond. Scop. carn. tom.* 2. *p.* 214.

Typha clava maſcula a feminea remota. *Hall. hiſt. n.* 1306.

Habitat & floret cum præc. Per.

SPARGANIUM. Masc. Amentum ſubrotundum. Calyx triphyllus. Corolla nulla.

Fem. Amentum, calyx, corolla ut in mare. Stigma bifidum. Drupa exſucca, monoſperma.

erectum. 1. S. foliis erectis triquetris. *Sp. pl. p.* 1378. *Curtis lond. Scop. carn. tom.* 2. *p.* 215.

Sparganium caule foliisque erectis. *Hall. hiſt. n.* 1303.

In aquis lente fluentibus, vel ſtagnantibus. Fl. Iulio. Per.

CAREX. Masc. Amentum imbricatum. Calyx monophyllus. Corolla nulla.

Fem. Amentum, calyx, corolla maris. Nectarium inflatum bidentatum. Stigmata duo vel tria. Semen triquetrum intra nectarium.

* *Spica una ſimplici.*

dioica. 1. C. ſpica ſimplici dioica. *Sp. pl. p.* 1379. *Fl. dan. t.* 369.

In udis. Fl. ab Aprili ad Iunium. Per.

Stigmata duo.

pulicaris. 2. C. ſpica ſimplici androgyna, ſuperne maſcula, capſulis ſemininibus teretibus. *Retzii ſcand. p.* 216. *Sp. pl. p.* 1380. *Leers herb. p.* 198. *t.* 14. *f.* 1.

Carex pſyllophora. *Ehrh. beytr. tom.* 1. *p.* 186.

In udis paluſtribus. Fl. Majo, Iunio. Per.

Stigmata duo.

leucoglochin. 3. C. ſpica ſimplici androgyna, ſuperne maſcula, capſulis reflexis, ſemininibus triquetris. *Retzii ſcand. p.* 217. *Ehrh. beytr. tom.* 1. *p.* 186.

Carex pauciflora. *Ligthf. ſcot.* 543. *t.* 6.

In paluſtribus Pannoniæ, Carnioliæ. Fl. cum præc. Per.

Stigmata tria.

Bellardi. 4. C. ſpica ſimplici hermaphrodita, culmo tereti, foliis capillaribus. *Allion. pedem. tom.* 2. *p.* 264. *t.* 92. *f.* 2.

In alpibus ſtyriacis, tyrolenſibus. Fl. Iulio, Auguſto. Per.

cyperoides. 5. C. capitulo terminali ſubrotundo, floribus ſimpliciſſimis ſubulatis, involucro longo. *Syſt. veget. p.* 843.

Carex bohemica. *Schreb. gr. t.* 28. *f.* 3..

Carex bohemica aquatica annua, cyperi facie, caule molli, capitulis in glomeratam ſpicam digeſtis. *Mich. gen. p.* 70. *t.* 33. *f.* 19.

In Auſtriæ paluſtribus prope *Weitra. R. Liborius Miller.* Fl. Iunio. Per.

juncifolia. 6. C. ſpica unica ſubrotunda, glumis ovatis, breviſſime ariſtatis. *Hall. hiſt. n.* 1354. *Allion. pedem. tom.* 2. *p.* 264. *t.* 92. *f.* 4.

Gramen cyperoides, juncifolium, ſpica ſimplici, in ſummitate culmi, atro-fuſca. *Scheuch. agr.* 492. *t.* 11. *f.* 7.

In collibus herbidis, campis, pratis Auſtriæ, Pannoniæ. Fl. Aprili, Majo. Per.

** *Spicis pluribus androgynis.*

leporina. 7. C. ſpica compoſita, ſpiculis ovatis ſeſſilibus approximatis alternis androgynis nudis. *Sp. pl. p.* 1381. *Leers herb. p.* 199. *t.* 14. *f.* 6.

Carex ſpicis nudis ſubrotundis, diſtinctis, in ſummo caule congeſtis. *Hall. hiſt. n.* 1361.

In udis. Fl. Aprili, Majo. Per.

brizoides. 8. C. ſpica compoſita diſticha nuda: ſpiculis androgynis oblongis contiguis, culmo ſuperne nudo. *Sp. pl. p.* 1381. *Scop. carn. tom.* 2. *p.* 231.

Carex ſpicis diſtichis, teretibus repandis. *Hall. hiſt. n.* 1358.

Carex fibrata radice, anguſtifolia, caule exquiſite triangulari, capitulis pulchellis, longioribus, & anguſtioribus, albicantibus, confertim naſcentibus. *Mich. gen. p.* 70. *t.* 33. *f.* 17.

In ſylvaticis. Fl. Majo, Iunio. Per.

præcox. 9. C. ſpica compoſita diſticha nudiuſcula: ſpiculis contiguis androgynis teretibus rectis, culmo ſuperne nudo. *Schreb. ſpic. p.* 63.

Carex anguſtifolia, caule triquetro, ſpicis pluribus elegantibus, parum inter ſe diſtantibus. *Segu. ver.* 1. *p.* 124. *t.* 1. *f.* 3.

In herbidis, pratis Auſtriæ, Pannoniæ. Fl. Aprili, Majo. Per.

diſticha. 10. C. ſpica compoſita ſubdiſticha: ſpiculis ovatis ſubimbricatis androgynis foliolo longiori inſtructis, culmo triquetro. *Huds. angl. ed.* 2. 403. *Schreb. ſpic. p.* 63.

Carex ſpicis in ſummo caule congeſtis, imis foliolo inſidentibus. *Hall. hiſt. n.* 1362.

Carex arenaria. *Leers herb. p.* 198. *t.* 14. *f.* 2.

Carex ſpicata. *Pollich. palat. tom.* 2. *p.* 562.

In pratis paluſtribus Auſtriæ, Pannoniæ. Fl. Majo. Per.

vulpina. 11. C. ſpica ſupradecompoſita inferne laxiore: ſpiculis androgynis ovatis ſeſſilibus glomeratis: ſuperne maſculis. *Sp. pl. p.* 1382. *Fl. dan. t.* 308. *Leers herb. p.* 199. *t.* 14. *f.* 5. *Scop. carn. tom.* 2. *p.* 230.

Carex ſpicis ſubrotundis echinatis, ſupremis confluentibus, capſulis roſtratis bifidis. *Hall. hiſt. n.* 1364.

In paluſtribus, ad ſepes, & alibi. Fl. Majo, Iunio. Per.

ſpicata. 12. C. ſpiculis ſubrotundis androgynis contiguis, capſulis ovatis acutis. *Huds. angl. ed.* 2. 405.

Carex muricata. *Leers herb. p.* 200. *t.* 14. *f.* 8. *Fl. dan. t.* 284.

Gramen cyperoides ſpicatum minus. *Raj. ſyn.* 424.

Carex paluſtris media, radice fibroſa, caule exquiſite triangulari, ſpica brevi compactiori. *Mich. gen. p.* 69. *t.* 33. *f.* 14.

In pratis humidis, ſylvis. Fl. Majo. Per.

murica-ta. 13. C. ſpiculis ſubovatis ſeſſilibus remotis androgynis, capſulis acutis divergentibus ſpinoſis. *Sp. pl. p.* 1382.

Carex canescens. *Leers. herb. p.* 201. *t.* 14. *f.* 3.

Cum præcedente. Fl. Majo. Per.

echina-ta. 14. C. ſpiculis ternis ſubovatis ſeſſilibus, capſulis mucronatis divergentibus. *Sibthorp. oxon. p.* 28.

Carex muricata. *Huds. angl. ed.* 2. 406.

Gramen cyperoides ſpicatum minimum, ſpica divulſa aculeata. *Raj. hiſt.* 1298. *Scheuch. agr.* 485. *t.* 11. *f.* 3.

In udis alpium Iudenburgenſium. Fl. Iunio, Iulio. Per.

remota. 15. C. ſpiculis ovatis ſubſeſſilibus remotis androgynis, foliis caulinis culmum æquantibus. *Sp. pl. p.* 1383. *Leers. herb. p.* 200. *t.* 15. *f.* 1.

Carex ſpicis androgynis, imis in ala prælongi folii ſeſſilibus, ſupremis nudis. *Hall. hiſt. n.* 1357.

Carex anguſtifolia, caule triquetro, capitulis pulchellis, ſtrigoſioribus, compactis, inter ſe diſtantibus, & in foliorum alis ſeſſilibus. *Mich. gen. p.* 70. *t.* 33. *f.* 16.

In ſylvis umbroſis. Fl. Majo, Per.

panicu-lata. 16. C. racemo compoſito, ſpiculis androgynis. *Sp. pl. p.* 1383. *Leers herb. p.* 201. *t.* 14. *f.* 4.

Carex ſpica paniculata. *Hall. hiſt. n.* 1368.

Carex radice repente, caule exquiſite triangulari, ſpica multiplici, fusca. *Mich. gen. p.* 68. *t.* 33. *f.* 7.

In udis Styriæ, Auſtriæ, Pannoniæ. Fl. Majo, Iunio. Per.

**** Spicis ſexu diſtinctis: femineis ſeſſilibus.*

flava. 17. C. ſpicis confertis ſubſeſſilibus ſubrotundis: maſcula lineari, capſulis acutis recurvis. *Sp. pl. p.* 1384. *Leers herb. p.* 202. *t.* 15. *f.* 6.

Carex ſpicis femininis petiolatis, erectis, ovatis, echinatis. *Hall. hiſt. n.* 1380.

Cyperoides paluſtre aculeatum, ſpica breviore. *Tourn. inſt.* 529. *Scheuch. agr.* 426.

In udis. Fl. Aprili, Majo. Per.

digitata. 18. C. ſpicis linearibus erectis: maſcula breviore inferioreque, bracteis aphyllis, capſulis diſtantibus. *Sp. pl. p.* 1384. *Leers herb. p.* 202. *t.* 16. *f.* 4. *Scop. carn. tom.* 2. *p.* 217.

Carex ſpicis femininis petiolatis, raris, maſculam ſuperantibus. *Hall. hiſt. n.* 1376.

In nemoribus. Fl. Martio, Aprili. Per.

montana. 19. C. ſpicis femininis ſeſſilibus ſubſolitariis ovatis maſculæ approximatis, culmo nudo, capſulis pubescentibus. *Sp. pl. p.* 1385. *Leers herb. p.* 203. *t.* 16. *f.* 6.

Carex foliis cariophyllæis, ſpicis ſeſſilibus approximatis, paucifloris, capſulis ovatis, ſubhirſutis. *Hall. hiſt. n.* 1372.

Carex præcox. *Iacqu. auſtr. t.* 446.

In montanis, ſylvaticis. Fl. primo vere. Per.

tomentoſa. 20. C. ſpicis femineis ſubpedunculatis, capſulis ſubgloboſis tomentoſis. *Syſt. veget. p.* 844. *Leers hurb. p.* 203. *t.* 15. *f.* 7.

Carex anguſtifolius, ſpicis ſeſſilibus, teretibus, approximatis, capſulis hirſutis. *Hall. hiſt. n.* 1373.

Cyperoides anguſtifolium, montanum, foliculis ſeminalibus villoſis. *Scheuch. agr.* 423.

In pratis udis. Fl. Majo, Iunio. Per.

filiformis. 21. C. ſpica maſcula oblonga, femineis ſeſſilibus oblongis: inferiore foliolo proprio breviore. *Sp. pl.* *p.* 1385. *Leers herb. p.* 204. *t.* 16. *f.* 5.

In collibus herbidis, montanis ſylvaticis, pratis ſiccioribus. Fl. Aprili. Per.

pilulifera. 22. C. ſpicis terminalibus confertis ſubrotundis, maſcula oblonga. *Sp. pl. p.* 1385. *Fl. dan. t.* 371.

In paluſtribus, pratis humidis Auſtriæ, Pannoniæ. Fl. Majo. Per.

humilis. 23. C. ſpica maſcula oblonga, femineis ſerioribus remotis ſubpedunculatis paucifloris, culmo foliis breviore. *Schreb. ſpic. p.* 65.

Cyperoides longifolium, culmo pumilo veluti folioſo, ſpicis obſeſſo. *Scheuch. agr.* 408.

In collibus apricis, ad ſylvarum margines Auſtriæ, Styriæ, Pannoniæ. Fl. Martio, Aprili. Per.

Michelii. 24. C. ſpica maſcula obovata, femineis ſubpedunculatis alternis: glumis ariſtatis, culmo ſeneſcente proſtrato.

Cyperoides ſylvarum, anguſtifolium, caule trilatero, ſpicis parvis, capſulis rarius diſpoſitis obeſis, in anguſtum collum vix ab una parte, ab altera vero magis bifidum coarctatis. *Mich. gen. p.* 58. *t.* 32. *f.* 5.

In Auſtriæ, Pannoniæ collibus, pratis ſiccioribus, montanis ſylvaticis. Fl. Majo. Per.

**** *Spicis ſexu diſtinctis: femineis pedunculatis.*

atrata. 25. C. ſpicis androgynis terminalibus pedunculatis: florentibus erectis; fructiferis pendulis. *Sp. pl. p.* 1386. *Fl. dan. t.* 158. *Scop. carn. tom.* 2. *p.* 222.

Carex ſpicis ovatis, petiolatis, congeſtis, ſeneſcendo nutantibus. *Hall. hiſt. n.* 1369.

Cyperoides alpinum, pulchrum, foliis caryophyllæis, ſpicis atris & tumentibus. *Scheuch. agr.* 481. *t.* 11. *f.* 1. 2.

In alpibus. Fl. Iunio, Iulio. Per.

limoſa. 26. C. ſpicis ovatis pendulis: maſcula longiore erectiore, radice repente. *Sp. pl. p.* 1386. *Fl. dan. t.* 646.

Carex ſpicis femininis pendulis, capſulis ovatis compreſſis. *Hall. hiſt. n.* 1392.

Carex *elegans*, *Willd prodr.* 104. *t.* 1. *f.* 4.

Cyperoides ſpica pendula, breviore, ſquamis e ſpadiceo, vel fuſco rutilante viridibus. *Scheuch. agr.* 443. *t.* 10. *f.* 13.

In alpium ſtyriacarum, carniolicarum udis. Fl. Iunio, Iulio. Per.

ſylvatica. 27. C. ſpicis pendulis: maſcula erecta: femineis filiformibus pedunculo brevioribus, capſulis acutis diſtantibus. *Schreb. ſpic. p.* 62. *Huds. angl. ed.* 2. 411. *Fl. dan. t.* 404.

Carex ſpicis femineis pendulis raris, prælongis, capſulis roſtratis, bifidis. *Hall. hiſt. n.* 1395.

Carex capillaris. *Leers herb. p.* 205. *t.* 15. *f.* 2.

Carex patula. *Scop. carn. tom.* 2. *p.* 226. *t.* 59.

Cyperoides ſylvarum, tenuius ſpicatum. *Tourn. inſt.* 530. *Scheuch. agr.* 418.

In ſylvis. Fl. Majo.

capillaris. 28. C. ſpicis pendulis: maſcula erecta, femineis oblongis, capſulis nudis acuminatis. *Sp. pl. p.* 1386. *Fl. dan. t.* 168. *Scop. carn. tom.* 2. *p.* 220. *t.* 59.

Carex ſpicis femininis raris, pendulis, capſulis ovato-conicis mucronatis. *Hall. hiſt. n.* 1394.

Cyperoides alpinum ſpicis feminiferis pendulis binis in ſummo caule. *Segu. ver. ſuppl. p.* 83. *t.* 3. *f.* 1.

In alpinis, & ſubalpinis. Fl. Iunio, Iulio. Per.

piloſa. 29. C. culmo triquetro: ſpicis erectis, remotis: femineis linearibus, folio proprio longioribus. *Scop. carn. tom. 2. p.* 226.

Carex ſpicis femineis petiolatis erectis, floribus remotis. *Hall. hiſt. n.* 1379.

Cyperoides latifolium, hirſutum, ſpicis tenuiſſimis. *Scheuch. agr.* 412.

In ſylvis montanis. Fl. Aprili. Per.

palleſcens. 30. C. ſpicis pendulis: maſcula erecta, femineis ovatis imbricatis, capſulis confertis obtuſis. *Sp. pl. p.* 1386. *Fl. dan. t.* 444. *Leers herb. p.* 206. *t.* 15. *f.* 4.

Carex ſpicis femininis pendulis, capſulis ovato-conicis. *Hall. hiſt. n.* 1393.

Cyperoides polyſtachyon, flavicans, ſpicis prope ſummitatem caulis. *Mich. gen. p.* 61. *t.* 32. *f.* 13.

In montanis, & alpinis. Fl. Majo, Iunio. Per.

ferruginea. 31. C. ſpicis erectis pedunculatis lanceolatis: maſcula terminali, femineis remotis tenuioribus. *Roth. germ. tom. 2. p.* 455. *Scop. carn. tom. 2. p.* 225.

Cyperoides anguſtifolium, alpinum, ſpica ſpadicea tenuiore. *Scheuch. agr.* 413. *t.* 10. *f.* 6.

In Carnioliæ, Tyrolis montanis, & alpinis.

firma. 32. C. culmo foliisque firmis erectis, ſpicis femineis, longe petiolatis, erectis: ſquamis retuſis mucronatis. *Mygind. Hall. hiſt. n.* 1383.

In alpibus auſtriacis, ſtyriacis. Fl. æſtate. Per.

panicea. 33. C. ſpicis pedunculatis erectis remotis: femineis linearibus, capſulis obtuſiuſculis inflatis. *Sp. pl. p.* 1387. *Fl. dan. t.* 261. *Leers herb. p.* 206. *t.* 15. *f.* 5.

In udis, paluſtribus. Fl. Aprili, Majo. Per.

Pſeudo-cyperus. 34. C. ſpicis pendulis, pedunculis geminatis. *Sp. pl. p.* 1387.

Carex ſpicis femininis pendulis, aſperis, capſulis longe ariſtatis, retroverſis. *Hall. hiſt. n.* 1397.

In Auſtria in paluſtribus prope *Himberg*; in Pannonia prope Poſonium, *Sövényháza*, alibique, Fl. Majo, Iunio. Per.

alba. 35. C. culmo tereti nudo, ſpicis pedunculatis, erectis, albidis: feminea ſupra marem elata, capſulis lævibus. *Scop. carn. tom.* 2. *p.* 216. *Hanke in Iacqu. collect. tom.* 2. *p.* 81. *Allion. pedem. tom.* 2. *p.* 269.

Carex ſpicis femineis raris, petiolatis, teretibus, acutis, marem æquantibus. *Hall. hiſt. n.* 1377.

Gramen ſylvaticum, anguſtifolium, ſpica alba. *Bauh. pin.* 4. *prodr.* 22. *Scheuch. agr.* 410. *t.* 10. *f.* 4. 5.

In montanis ſylvaticis Auſtriæ, Styriæ, Carnioliæ. Fl. ab Aprili ad Iunium. Per.

diſtans. 36. C. ſpicis remotiſſimis ſubſeſſilibus bractea vaginante, capſulis angulatis mucronatis. *Sp. pl. p.* 1387.

Carex ſpicis breviſſime petiolatis, diſſitis. *Hall. hiſt. n.* 1382.

In uliginoſis, paluſtribus, alibique, Fl. Majo. Per.

cæspitoſa. 37. C. ſpicis erectis cylindricis ternis ſubſeſſilibus: maſcula terminali, culmo triquetro. *Sp. pl. p.* 1388. *Parkinſon theatr.* 1266. *f.* 3.

In uliginoſis, paluſtribus Auſtriæ, Styriæ, Pannoniæ. Fl. Majo, Iunio. Per.

***** *Spicis ſexu diſtinctis: maſculis pluribus.*

flacca. 38. C. ſpicis maſculis geminis rectis, femineis pedunculatis pendulis, capſulis ovato-triquetris confertis. *Retzii ſcand. p.* 222. *Schreb. ſpic. app.* 669.

Carex ſpicis maribus una pluribus, femininis pendulis, capſulis ovato-triquetris confertis. *Hall. hiſt. n.* 1408.

Carex limoſa. *Leers herb. p.* 204. *t.* 15. *f.* 3.

Carex glauca. *Scop. carn. tom.* 2. *p.* 223.

Cyperoides paluſtre, ſpicis purpuro-ſpadiceis tenuibus pediculis inſidentibus. *Scheuch. agr.* 467.

In ſylvaticis, uliginoſis, paluſtribus. Fl. Majo. Per.

laſiocarpa. 39. C. ſpicis maſculis geminis: femineis remotis ſubſeſſilibus, bractea longiſſima, capſulis hirſutis. *Ehrh. beytr. tom.* 3. *p.* 73. *Retzii ſcand. p.* 222.

Carex ſplendida. *Willd prodr. p.* 103. *t.* 1. *f.* 3.

In paluſtribus Auſtriæ prope *Himberg.* Fl. Majo. Per.

pendula. 40. C. ſpicis ſubſeſſilibus pendulis: maſcula erecta; femineis cylindricis longiſſimis, capſulis ſubrotundis acuminatis. *Huds. angl. ed.* 2. 411. *Curtis lond.*

Carex ſpicis femineis pendulis longiſſimis, capſulis mucronatis, ovatis. *Hall. hiſt. n.* 1396.

Carex maxima. *Scop. carn. tom.* 2. *p.* 229.

Cyperoides ſpica pendula longiore & anguſtiore. *Tourn. inſt.* 529. *Scheuch. agr.* 445.

In udis ſylvaticis. Fl. Majo, Iunio. Per.

gracilis. 41. C. ſpicis maſculis & femineis pluribus, ſubfiliformibus, floribus digynis. *Curtis lond. Sibthorp oxon. p.* 32.

Gramen cyperoides majus anguſtifolium. *Raj. ſyn.* 417.

In paluſtribus, ad foſſas, fluviorum ripas. Fl. Majo. Per.

acuta. 42. C. ſpicis maſculis pluribus, obtuſis, ſquamis obtuſiuſculis, caule acutangulo. *Curtis lond. Sibthorp. oxon. p.* 32.

Cum præc. Fl. Aprili. Per.

craſſa. 43. C. ſpicis maſculis pluribus triquetris: femineis pedunculatis, calycibus linearibus acuminatis, capſulis ſubinflatis bicornibus. *Retzii ſcand. p.* 223. *Ehrh. beytr. tom.* 4. *p.* 43.

Carex riparia. *Curtis lond. Sibthorp oxon. p.* 31.

Habitat & floret cum præc. Per.

veſicaria. 44. C. ſpicis maſculis pluribus: femineis pedunculatis, capſulis inflatis acuminatis, culmo acute triquetro. *Retzii ſcand, p.* 223. *Fl. dan. t.* 647.

In Auſtriæ, Pannoniæ paluſtribus, uliginoſis. Fl. Aprili, Majo. Per.

obtuſangula. 45. C. ſpicis maſculis pluribus: femineis pedunculatis, capſulis inflatis acuminatis, culmo obſolete triquetro. *Retzii ſcand. p.* 223.

In Styriæ ſuperioris aquoſis. Fl. Iunio. Per.

hirta. 46. C. ſpicis remotis: maſculis pluribus: femineis ſubpedunculatis erectis, ſquamis longis ariſtatis, capſulis hirtis. *Sp. pl. p.* 1389. *Fl. dan. t.* 379. *Leers herb. p.* 208. *t.* 16. *f.* 3.

Carex caule foliis capſuliſque hirſutis. *Hall. hiſt. n.* 1403.

Cyperoides polyſtachyon, lanuginoſum. *Tourn. inſt.* 529. *Scheuch. agr.* 478.

In pratis, aliiſque ſiccis & humidis locis. Fl. Majo. Per.

TETRANDRIA.

BETULA. Masc. Amentum cylindraceum. Calyx monophyllus, trifidus, triflorus. Corolla quadripartita.

Fem. Amentum cylindraceum. Calyx monophyllus, ſubtrifidus, biflorus. Semen utrinque membrana alatum.

alba. 1. B. foliis deltoidibus acutis duplicato-ſerratis glabris, ſtrobilorum ſquamis lobis lateralibus rotundatis, petiolis glabris pedunculis longioribus. *Ait. Kew. tom.* 3. *p.* 336. *Willd. baumz. p.* 36. *Sp. pl. p.* 1393. *Scop. carn. tom.* 2. *p.* 232. *Reitter und Abel. abbild. t.* 15.

Betula foliis cordato-lanceolatis ſerratis. *Hall. hiſt. n.* 1628.

Le Bouleau. Regn. bot.

In ſylvis. Fl. Majo, Iunio. Arb.

Variat foliis glabris, ſcabris; ramis pendulis, adſcendentibus.

nana. 2. B. foliis orbiculatis crenatis, utrinque glabris. *Willd. baumz. p.* 42. *Sp. pl. p.* 1394. *Fl. lapp. n.* 266. *t.* 6. *f.* 4. *Am. acad. tom.* 1. *t.* 1. *Hall. hiſt. n.* 1629.

In Carinthia in turfoſis alpium reichenavienſium ad S. Laurentium. *Wulfen.* Ego in Styria in alpibus judenburgenſibus ad *Bürgerſee.* Fl. poſt ſolutas nives. Fr.

Alnus. 3. B. pedunculis ramoſis, foliis ſubrotundo-cuneiformibus obtuſiſſimis glutinoſis: axillis venarum ſubtus villoſis. *Ait. Kew. tom.* 3. *p.* 338. *Willd. baumz. p.* 44. *Reitter und Abel abbild. t.* 16.

Alnus foliis glabris, rotundis, rotunde crenatis, ſpongiolis ad nervorum angulos. *Hall. hiſt. n.* 1630.

Betula Alnus, var. glutinoſa. *Sp. pl. p.* 1394.

Alnus vulgaris. *Clus. hiſt. p.* 12.

Alnus rotundifolia glutinoſa viridis. *Bauh. pin.* 428.

Betula laciniata, ſeu Alnus foliis eleganter inciſis. *Zann. iſt. p.* 8. *t.* 277.

L' Aune. Regn. bot.

Ad pagos, fluvios, in montanis, & in alpibus. Varietas foliis laciniatis in Carnia *nella valle del Chiadin.* Fl. a Februario ad Aprilem. Arb.

incana. 4. B. pedunculis ramoſis, foliis ſubrotundo-ellipticis acutis ſubtus pubeſcentibus: axillis venarum nudis, ſtipulis lanceolatis. *Ait. Kew. tom.* 3. *p.* 339. *Willd. baumz. p.* 45. *Reitter und Abel abbild. t.* 16.

Alnus foliis mucronatis, acute ſerratis, ſubtus lanuginoſis. *Hall. hiſt. n.* 1631.

Betula Alnus, var. incana. *Sp. pl. p.* 1394.

Alnus altera. *Clus. hiſt. p.* 12.

Alnus folio incano. *Bauh. pin.* 428.

Habitat & floret cum præc. Arb.

BUXUS. Maſc. Calyx triphyllus. Petala duo. Germinis rudimentum.

Fem. Calyx tetraphyllus. Petala tria. Styli tres. Capſula triroſtris, trilocularis. Semina duo.

ſempervirens. 1. Buxus. *Sp. pl. p.* 1394. *Hall. hiſt. n.* 1610. *Scop. carn. tom.* 2. *p.* 234.

Le Buis. Regn. bot.

Occurrit ad pagos, in hortis neglectis. Fl. primo vere. Fr.

Germini inſident nectaria tria cum ſtylis alternantia. Ehrh. beytr. tom. 4. p. 175.

URTICA. Maſc. Calyx tetraphyllus. Corolla nulla. Nectarium centrale, cyathiforme.

Fem. Calyx bivalvis. Corolla nulla. Semen unicum, nitidum.

urens. 1. U. foliis oppoſitis ovalibus. *Sp. pl. p.* 1396. *Fl. dan. t.* 739. *Scop. carn. tom.* 2. *p.* 234.

Urtica ſexubus ſede disjunctis, foliis ovato-lanceolatis ſerratis, ſpicis oblongis. *Hall. hiſt. n.* 1615.

L' Ortie mineure. Bulliard herb. t. 230.

In cultis, & incultis. Fl. æſtate.

dioica. 2. U. foliis oppoſitis cordatis, racemis geminis. *Sp. pl. p.* 1396. *Scop. carn. tom.* 2. *p.* 234. *Fl. dan. t.* 740.

Urtica ſtirpibus ſexu diſtinctis, foliis ovato-lanceolatis, productis, ſerratis. *Hall. hiſt. n.* 1614.

La grande Ortie. Regn. bot.

In ſylvaticis, ad ſepes, vias. Fl. æſtate. Per.

MORUS. Maſc. Calyx quadripartitus. Corolla nulla.

Fem. Calyx tetraphyllus. Corolla nulla. Styli duo. Calyx maximus, ſucculentus factus. Semen unicum.

alba. 1. M. foliis oblique cordatis lævibus. *Sp. pl. p.* 1398. *Scop. carn. tom.* 2. *p.* 235. *Willd. baumz. p.* 195.

Morus foliis ſcabris, cordatis & ſemitrilobis. *Hall. hiſt. n.* 1611.

Ad vias, pagos. Fl. Majo. Arb.

nigra. 2. M. foliis cordatis ſcabris. *Sp. pl. p.* 1398.

Le Meurier noir. Regn. bot.

In Litorali, Forojulio ad vias, plateas, vineas. Fl. Iunio. Arb.

XANTHIUM. Maſc. Calyx communis imbricatus. Corolla compoſita, in hemisphærium digeſta. Propria monopetala, quinquefida, infundibuliformis. Receptaculum paleaceum.

Fem. flores infra maſculos. Calyx involucrum diphyllum, biflorum. Corolla nulla. Drupa ſicca, undique aculeis uncinatis tecta, apice bifida. Nucleus bilocularis.

ſtrumarium. 1. X. caule inermi, foliis cordatis, trinerviis. *Sp. pl. p.* 1400. *Scop. carn. tom.* 2. *p.* 236.

Xanthium foliis ſemitrilobatis, aculeis uncinatis. *Hall. hiſt. n.* 1621.

Le petit glouteron. Regn. bot.

Ad fimeta, vias, pagos. Fl. Iulio, Auguſto.

ſpinoſum. 2. X. ſpinis ternatis, foliis trilobis. *Syſt. veget. p.* 852.

Xanthium luſitanicum ſpinoſum. *Pluk. alm.* 206. *t.* 239. *f.* 1.

Tergeſti ad latera viarum. Fl. Iunio, Iulio, Auguſto.

AMARANTHUS. Maſc. Calyx tri- ſeu pentaphyllus. Corolla nulla. Stamina tria ſeu quinque.

Fem. Calyx & corolla ut in mare. Styli tres. Capſula unilocularis, circumſciſſa. Semen unicum.

Blitum. 1. A. glomerulis lateralibus trifidis, foliis ovatis retuſis, caule diffuſo. *Sp. pl. p.* 1405. *Scop. carn. tom.* 2. *p.* 237.

Amaranthus procumbens foliis ovatis emarginatis. *Hall. hiſt. n.* 1606.

Blitum album minus. *Raj. hiſt.* 200. *Pluk. phyt.* 212. *f.* 2.

In cultis, ruderatis, ad vias. Fl. æſtate.

viridis. 2. A. glomerulis triandris, floribus masculis trifidis, foliis ovatis emarginatis, caule erecto. *Sp. pl. p.* 1405.

Blitum minus album polyſpermum folio ſubrotundo. *Sloan. jam.* 49. *hiſt.* 1. *t.* 92. *f.* 1.

Habitat & floret cum præc.

POLYANDRIA.

CERATOPHYLLUM. Maſc. Calyx multipartitus. Corolla nulla. Stamina a ſexdecim ad viginti.

Fem. Calyx & corolla ut in mare. Piſtillum unicum. Stylus nullus. Semen unicum, nudum.

demerſum. 1. C. foliis dichotomo-bigeminis, fructibus triſpinoſis. *Sp. pl. p.* 1409. *Scop. carn. tom.* 2. *p.* 238.

Ceratophyllum foliis ſubſpinoſis, capſulis tricornibus. *Hall. hiſt. n.* 1599.

Hydroceratophyllum folio aſpero quatuor cornibus armato. *Vaill. act.* 1719. *p.* 21. *t.* 2. *f.* 1.

In aquis ſtagnantibus, vel lente fluentibus. Fl. Iulio, Auguſto. Per.

MYRIOPHYLLUM. Maſc. Calyx tetraphyllus. Corolla nulla. Stamina octo.

Fem. Calyx & corolla ut in mare. Germina quatuor. Stylus nullus. Stigmata pubeſcentia. Semina quatuor, nuda.

ſpicatum. 1. M. floribus maſculis interrupte ſpicatis. *Sp. pl. p.* 1409. *Scop. carn. tom.* 2. *p.* 238. *Fl. dan. t.* 681.

Pentapteris ſpica nuda. *Hall. hiſt. n.* 993.

Millefolium aquaticum pennatum ſpicatum. *Bauh. pin.* 141. *prodr.* 73.

In aquis ſtagnantibus, vel lente fluentibus, Fl. æſtate. Per.

verticillatum. 2. M. floribus omnibus verticillatis. *Sp. pl. p.* 1410. *Fl. dan. t.* 1046.

Pentapteris floribus alaribus. *Hall. hiſt. n.* 992.

Millefolium aquaticum, floſculis ad foliorum nodos. *Bauh. pin.* 141.

Myriophyllum aquaticum minus. *Cluſ. hiſt. p. CCLII.*

Habitat & floret cum præc. Per.

SAGITTARIA. Maſc. Calyx triphyllus. Corolla tripetala. Filamenta fere viginti quatuor.

Fem. Calyx & corolla maris. Piſtilla numeroſa. Receptaculum globoſum ſemina multa, nuda in globum colligens.

ſagittifolia. 1. S. foliis ſagittatis acutis. *Sp. pl. p.* 1410. *Fl. dan. t.* 172.

Sagitta foliis acuminatis. *Hall. hiſt. n.* 1185.

Sagitta major. *Scop. carn. tom.* 2. *p.* 239.

In aquis ſtagnantibus. Fl. Iulio, Auguſto.

Folia, quæ primo prodeunt, linearia, quæ ſerius, ſagittata ſunt.

POTERIUM. Masc. Calyx tetraphyllus. Corolla quadripartita. Stamina plurima.

Fem. in eadem ſpica ſupra maſculos. Calyx & corolla maris. Piſtilla duo. Bacca e tubo corollæ indurato.

Sangui-ſorba. 1. P. inerme, caulibus ſubanguloſis. *Sp. pl. p.* 1411. *Curtis lond.*

Pimpinella polyſtemon. *Hall. hiſt. n.* 706.

Sanguiſorba minor. *Scop. carn. tom.* 1. *p.* 110.

In pratis, paſcuis, collibus, & alibi. Fl. æſtate. Per.

In flore maſculo germen & ſtigmata; at imperfecta ſteriliaque omnia. Leers herb. p. 212.

QUERCUS. Maſc. Calyx quinquefidus fere. Corolla nulla. Stamina quinque ad decem.

Fem. Calyx monophyllus, integerrimus. Corolla nulla. Styli duo ad quinque. Semen unicum, ovatum.

Ilex. 1. Q. foliis ovato-oblongis indiviſis ſerratisque petiolatis ſubtus incanis, cortice integro. *Sp. pl. p.* 1412. *Scop. carn. tom.* 2. *p.* 240.

Ilex major. *Cluſ. hiſt. p.* 23.

Le Chêne verd. Regn. bot.

Ad litus maris adriatici. Fl. Majo. Arb.

coccifera. 2. Q. foliis ovatis indiviſis ſpinoſo-dentatis, utrinque glabris. *Sp. pl. p.* 1413. *Scop. carn. tom.* 2. *p.* 240.

Ilex coccigera. *Cluſ. hiſt. p.* 24.

In Iſtria, & in inſulis venetis. Fl. Majo. Fr.

Suber. 3. Q. foliis ovato - oblongis indiviſis ſerratis ſubtus tomentoſis, cortice rimoſo fungoſo. *Sp. pl. p.* 1413.

Suber latifolium ſempervirens. *Bauh. pin.* 424.

Le Liege. Regn. bot.

Habitat & floret cum praec. Arb.

Robur. 4. Q. foliis oblongis pedunculatis glabris ſinuatis: lobis rotundatis, glandibus oblongis ſeſſilibus. *Willd. baumz. p.* 277.

Quercus foliis deciduis petiolatis oblongis ſuperne latioribus, ſinubus acutioribus, angulis obtuſis: glandibus ſubſeſſilibus. *Du Roi baumz. ed.* 1. 2. *p.* 237. *Reitter und Abel abbild. t.* 1.

In ſylvis. Fl. Majo. Arb.

pedunculata. 5. Q. foliis oblongis ſubſeſſilibus glabris ſinuatis: lobis rotundatis, glandibus oblongis pedunculatis. *Willd. baumz. p.* 178.

Quercus *femina*, foliis deciduis ſubpetiolatis oblongis obtuſis pinnato-ſinuatis: glandibus pedunculatis ſubſolitariis. *Du Roi baumz. ed.* 1. 2. *p.* 240. *Reitter und Abel abbild. t.* 2.

Habitat & floret cum praec. Arb.

Cerris. 6. Q. foliis ſinuato - pinnatifidis ſubtus pubescentibus: laciniis acutiusculis, ramentis axillaribus filiformibus, calycibus echinato - ramentaceis. *Ait Kew. tom.* 3. *p.* 358. *Sp. pl. p.* 1415. *Willd. baumz. p.* 280. *Du Roi baumz. ed.* 1. 2. *p.* 259. *t.* 5. *f.* 1.

Quercus *crinita*, foliis oblongis lyrato - pinnatifidis ſubtus pubescentibus, glandibus ſeſſilibus, cupula crinito-hiſpida. *Lamarck encycl.* 1. *p.* 710.

In ſylvis. Fl. Majo. Arb.

Varietas foliis leviter ſinuatis, planiusculis, in Auſtria, Pannonia, Carniolia calidiore, Littorali crescens, eſt Quercus Ægilops Scopolii.

IUGLANS. Masc. Calyx monophyllus, ſquamiformis. Corolla ſexpartita. Filamenta octodecim.

Fem. Calyx quadrifidus, ſuperus. Corolla quadripartita. Styli duo. Drupa nucleo ſulcato.

regia. 1. I. foliolis ovalibus glabris ſubſerratis ſubæqualibus. *Sp. pl. p.* 1415. *Scop. carn. tom.* 2. *p.* 241.

Iuglans foliis ſeptenis, ovato-lanceolatis, integerrimis. *Hall. hiſt. n.* 1624.

Le Noyer. Regn. bot.

Ad vias, pagos. Fl. Majo. Arb.

FAGUS. Masc. Amentum. Calyx quinquefidus, campanulatus. Corolla nulla.

Fem. Calyx quadridentatus. Corolla nulla. Stylus trifidus. Capſula muricata, quadrivalvis. Nuces duæ.

Caſtanea. 1. F. foliis lanceolatis acuminato-ſerratis ſubtus nudis. *Sp. pl. p.* 1416. *Willd. baumz. p.* 111.

Caſtanea foliis ovato-lanceolatis ſerratis, dentibus aduncis. *Hall. hiſt. n.* 1623.

Caſtanea ſativa. *Scop. carn. tom.* 2. *p.* 242.

Le Chataignier. Regn. bot.

In Tyroli vinifera, Iſtria, Forojulio, Litorali, Pannonia, occurrit quoque per Auſtriam, reliquasque provincias. Fl. Iunio. Arb.

Capſula ſpinis ramoſis rigidis echinata. Nuces duæ vel tres, hinc convexæ, inde planiusculæ. Linn.

ſylvatica. 2. F. foliis ovatis obſolete ſerratis. *Sp. pl. p.* 1416. *Willd. baumz. p.* 113. *Reitter und Abel abbild. t.* 6.

Fagus foliis ovato-lanceolatis, oris undulatis. *Hall. hiſt. n.* 1622.

Caſtanea fagus. *Scop. carn. tom.* 2. *p.* 242.

In ſylvis. Fl. Majo, Iunio. Arb.

Pericarpium coriaceum, triquetrum, uniloculare, univalve, ſpinis mollibus muricatum. Semen plerumque unicum, triquetrum. Ehrh. beytr. tom. 4. p. 176.

CARPINUS. Maſc. Amentum conſtans ſquamis unifloris, ciliatis. Corolla nulla.

Fem. Amentum conſtans ſquamis unifloris, villoſis. Corolla nulla. Germina duo. Styli ſingulis duo. Nux ovata.

Betulus. 1. C. ſtrobilis foliaceis laxis, ſquamis profunde trifidis, lobis lanceolatis, intermedio longiori. *Lamarck encycl.* 1. *p.* 699. *Sp. pl. p.* 1416. *Scop. carn. tom.* 2. *p.* 243. *Hall. hiſt. n.* 1627. *Willd. baumz. p.* 52. *Reitter und Abel abbild. t.* 12.

Oſtrya ulmo ſimilis, fructu in umbilicis foliaceis. *Bauh. pin.* 427.

In ſylvis. Fl. Majo, Iunio. Arb.

Semen intra baſin ſquamæ amentaceæ concavæ gerit. Linn.

Oſtrya. 2. C. ſquamis ſtrobilorum inflatis, *Sp. pl. p.* 1417.

Oſtria italica, Carpini folio, fructu longiore ſeu breviore habitiore. *Mich. gen. p.* 223. *t.* 104. *f.* 1. 2.

Oſtrya Ulmo ſimilis, fructu racemoſo, lupulo ſimili. *Bauh. pin.* 427.

In Litorali. Fl. Majo. Arb.

Semen intra ſquamam amentaceam inflatam recondit. Linn.

CORYLUS. Maſc. Amentum conſtans ſquamis trifidis. Corolla nulla. Stamina octo.

Fem. Calyx diphyllus, lacerus. Corolla nulla. Styli duo. Nux ovata.

Avellana. 1. C. ſtipulis ovatis obtuſis. *Sp. pl. p.* 1417. *Scop. carn. tom.* 2. *p.* 245.

Corylus ſepium. *Hall. hiſt. n.* 1625.

Le Coudrier. Regn. bot.

In ſylvaticis, ad ſepes. Fl. primo vere. Fr.

Colurna. 2. C. ſtipulis linearibus acutis. *Sp. pl. p.* 1417.

Corylus byzantina. *Herm. Lugd.* 191. *Seb. mus.* 1. *t.* 27. *f.* 2.

Avellana byzantina. *Clus. hiſt. p.* 11.

In Pannoniæ finitimis prope Semlinam. In Auſtria retro *Baden* in *Gainfahren.* Fl. cum præc. Arb.

MONADELPHIA.

PINUS. Maſc. Amentum. Calyx tetraphyllus. Corolla nulla. Stamina plurima. Antheræ nudæ.

Fem. Strobilus conſtans ſquamis bifloris. Corolla nulla. Piſtillum unicum. Nux.

ſylveſtris. 1. P. foliis geminis: primordialibus ſolitariis glabris. *Sp. pl. p.* 1418. *Scop. carn. tom.* 2. *p.* 247. *Pallas roſſic. t.* 2. *f.* 2.

Pinus foliis binis, convexo-concavis, julis maſculis ſolitariis alaribus. *Hall. hiſt. n.* 1660.

In montanis, ſubalpinis, alpinis. Fl. Majo, Iunio. Arb.

Folia ad lentem margine ſerrulata.

Cembra. 2. P. foliis quinis lævibus. *Sp. pl. p.* 1419. *Pallas roſſic. t.* 2.

Pinus foliis quinis triquetris. *Hall. hiſt. n.* 1659.

Pinaſter. *Mich. gen. p.* 223. *t.* 19.

In Styria in alpibus judenburgenſibus. In Tyrolis alpinis, & ſubalpinis. Fl. Iunio. Arb.

Larix. 3. P. foliis faſciculatis mollibus obtuſiuſculis, bracteis extra ſquamas ſtrobilorum extantibus. *Ait. Kew. tom.* 3. *p.* 369. *Sp. pl. p.* 1420. *Scop. carn. tom.* 2. *p.* 248. *Pallas roſſic. t.* 1. *Willd. baumz. p.* 216. *Reitter und Abel abbild. t.* 96.

Pinus foliis faſciculatis deciduis. *Hall. hiſt. n.* 1658.

Larix. *Clus. hiſt. p.* 34.

In ſylvis montanis, & ſubalpinis. Fl. Majo, Iunio. Arb.

Folia, quæ in faſciculum congeſta, obtuſa ſunt; acuta contra, quæ ſolitaria ſunt.

Picea. 4. P. foliis ſolitariis emarginatis. *Sp. pl. p.* 1420. *Scop. carn. tom.* 2. *p.* 245. *Pallas roſſic. t.* 1. *f. F. Willd. baumz. p.* 217. *Reitter und Abel abbild. t.* 98.

Pinnis foliis ſolitariis, planis, pectinatis, emarginatis. *Hall. hiſt. n.* 1657.

Abies. *Clus. hiſt. p.* 34.

In ſylvis. Fl. Majo, Iunio. Arb.

Abies. 5. P. foliis ſolitariis ſubulatis mucronatis lævibus bifariam verſis. *Sp. pl. p.* 1421. *Scop. carn. tom.* 2. *p.* 246. *Fl. dan. t.* 193. *Willd. baumz. p.* 221. *Reitter und Abel abbild. t.* 99.

Pinus foliis ſolitariis tetragonis mucronatis. *Hall. hiſt. n.* 1656.

Picea. *Clus. hiſt. p.* 33.

In ſylvis. Fl. Majo, Iunio. Arb.

SYNGENESIA.

MOMORDICA. Maſc. Calyx quinquefidus. Corolla quinquepartita. Filamenta tria.

Fem. Calyx & corolla maris. Stylus trifidus. Pomum elaſtice diſſiliens.

Elaterium. 1. M. pomis hiſpidis, cirrhis nullis. *Sp. pl. p.* 1434.

Cucumis ſylveſtris, aſininus dictus. *Bauh. pin.* 314. *Zann. iſt. p.* 91. *t.* 63.

Le Concombre ſauvage. Bulliard. herb. t. 81.

Flumine ad litus maris, ad vias. Fl. æſtate.

BRYONIA. Maſc. Calyx quinquedentatus. Corolla quinquepartita. Filamenta tria.

Fem. Calyx & corolla maris. Stylus quadrifidus. Bacca ſubgloboſa, polyſperma.

alba. 1. B. foliis palmatis utrinque calloſo-ſcabris, floribus monoicis. *Syſt. veget. p.* 869. *Scop. carn. tom.* 2. *p.* 250. *Fl. dan. t.* 813.

Bryonia. *Hall. hiſt. n.* 574.

Bryonia alba baccis nigris. *Bauh. pin.* 297.

Ad ſepes Litoralis, Forijulii. Fl. æſtate. Per.

dioica. 2. B. foliis palmatis utrinque colloſo-ſcabris, floribus dioicis. *Syſt. veget. p.* 870. *Iacqu. auſtr. t.* 199.

La Colouvrée. Regn. bot.

Ad ſepes Auſtriæ, Pannoniæ. Fl. æſtate. Per.

Baccæ rubræ.

DIOECIA.

DIANDRIA.

SALIX. Maſc. Amentum conſtans ſquamis unifloris. Corolla nulla. Glandula baſeos nectarifera.

Fem. Amentum, corolla, Glandula maris. Stylus bifidus. Capſula unilocularis, bivalvis. Semina papposa.

* *Foliis ſerratis glabris.*

triandra. 1. S. foliis ſerratis glabris, floribus triandris. *Sp. pl. p.* 1442. *Curtis lond. Scop. carn. tom.* 2. *p.* 259. *Hoffm. ſal.* 1. *p.* 45. *t.* 9. *f.* 1. 2. *t.* 10. *f.* 3. 4. *t.* 23. *f.* 2.

Salix foliis glabris elliptico-lanceolatis, ſerratis, ſtipulis dentatis, julis gracilibus triandris. *Hall. hiſt. n.* 1637.

Ad fluvios, lacus, torrentes. Fl. Majo. Fr.

phylicifolia. 2. S. foliis ſerratis glabris lanceolatis: crenis undatis. *Sp. pl. p.* 1442. *Fl. lapp. n.* 351. *t.* 8. *f. D. Wulfen in Iacqu. collect. tom.* 2. *p.* 139.

In Carinthia in monte *Loibel*; in convallibus taureri malnizenſis; in valle rablenſi per adſcenſum montis *Predil. Wulfen.* In Carnia in monte *Croſtis. I. Hoſt.* Fl. Majo, Iunio. Arb.

vitellina. 3. S. foliis ſerratis, ovato-lanceolatis, acutis, ſupra glabris: ſerraturis cartilagineis. *Hoffm. ſal.* 1. *p.* 57. *t.* 11. *f.* 1. *t.* 12. *f.* 2. 3. *t.* 24. *f.* 1. *Sp. pl. p.* 1442.

Salix lutea tenuior, ſativa, viminea. *Bauh. hiſt.* 1. *p.* 214.

Ad pagos, fluvios, rivulos, in pratis humidis. Fl. Majo. Arb.

fragilis. 4. S. foliis ſerratis glabris ovato-lanceolatis, petiolis dentato-glanduloſis. *Sp. pl. p.* 1443. *Fl. lapp. n.* 349. *t.* 8. *f. b. Willd. baumz. p.* 335. *Reitter und Abel abbild. t.* 27.

Salix folio longo latoque ſplendente, fragilis. *Raj. ſyn.* 448.

In ſylvis, ad fluvios, alibique. Fl. Aprili, Majo. Arb.

monandra. 5. S. foliis ſerratis glabris lineari-lanceolatis, ſuperioribus obliquis. *Hoffm. ſal.* 1. *p.* 18. *t.* 1. 5. *f.* 1. *t.* 13. *f.* 1.

Salix monoſtemon, foliis glabris, lineari-lanceolatis, ſerratis, ſuperne conjugatis, julis tomentoſis. *Hall. hiſt. n.* 1640.

Salyx purpurea, & Salix Helix. *Sp. pl. p.* 1444.

Ad fluvios, foſſas, lacus, alibique. Fl. Aprili. Fr.

myrſinites. 6. S. foliis ſerratis glabris, ovatis ſubdiaphanis. *Hoffm. ſal.* 1. *p.* 71. *t.* 17. *f.* 1. 2. *t.* 18. *f.* 1. 2. *t.* 19. *f.* 2. *t.* 24. *f.* 2. *Sp. pl. p.* 1445. *Fl. lapp. n.* 353. *t.* 8. *f. F. t.* 7. *f.* 6. *Wulfen in Iacqu. collect. tom.* 2. *p.* 136.

Salix foliis ovato-lanceolatis, venoſis, ſerratis, julis tomentoſis. *Hall. hiſt. n.* 1645.

Clagenfurti *am Fuſſe des Schmalbergels dem Walde der Ziegeln gegenüber*, partim extra, partim in foſſa. *Wulfen.* Ego in Styria in alpibus judenburgenſibus ad rivulos. Fl. Aprili, Majo. Fr.

arbuscula. 7. S. foliis ſubſerratis glabris ſubdiaphanis ſubtus glaucis. *Sp. pl. p.* 1445. *Iacqu. auſtr. t.* 408.

Salix foliis ſerratis glabris obovatis. *Fl. lapp. n.* 352. *t.* 8. *f. e.*

Salix foliis integris glabris ovatis confertis pellucidis. *Fl. lapp. n.* 356.

Salix foliis ſerratis glabris lanceolatis utrinque acutis. *Fl. lapp. n.* 360. *t.* 8. *f. M.*

In alpibus. Fl. Iunio. Fr.

herbacea. 8. S. foliis ſerratis glabris orbiculatis. *Sp. pl. p.* 1445. *Fl. lapp. n.* 355. *t.* 7. *f.* 3. 4. *t.* 8. *f.* 9. *Hoffm. ſal.* 1. *p.* 74. *t.* 20. *f.* 1. 2. 3. 4. *Fl. dan. t.* 117.

In alpiqus. Fl. Iulio, Auguſto.

retuſa. 9. S. foliis ſubſerratis glabris obovatis obtuſiſſimis. *Sp. pl. p.* 1445.

Salix foliis glabris ovatis nitentibus, julis pauciſloris. *Hall. hiſt. n.* 1648.

Salix alpina pumila, repens, retuſo ſerpylli folio lucido. *Bocc. mus. p.* 18.

In alpibus, Fl. Iulio, Auguſto. Fr.

** *Foliis glabris integerrimis.*

reticulata. 10. S. foliis integerrimis glabris ovatis obtuſis. *Sp. pl. p.* 1446. *Iacqu. en.* 296. *Fl. dan. t.* 212. *Hoffm. ſal.* 2. *t.* 25. *f.* 1. 3. *t.* 26. 27.

Salix foliis integris glabris ovatis ſubtus reticulatis. *Fl. lapp. n.* 359. *t.* 8. *f. L. t.* 7. *f.* 1. 2.

Salix foliis rotundis, integerrimis, ſubtus tomentoſis, julis gracilibus longe petiolatis. *Hall. hiſt. n.* 1650.

In alpibus. Fl. Iunio, Iulio. Fr.

*** *Foliis integerrimis, villoſis.*

arenaria. 11. S. foliis integerrimis ovatis acutis: ſupra ſubvilloſis, ſubtus tomentoſis. *Sp. pl. p.* 1447.

Salix foliis integris ſubtus villoſis ovatis acutis. *Fl. lapp. n.* 362. *t.* 8. *f.* o. *q.*

Salis foliis integerrimis, ovato-lanceolatis, ſubtus ſericeis, julis tomentoſis. *Hall. hiſt. n.* 1642.

In alpibus ſtyriacis, tyrolenſibus ad torrentes, rivulos. Fl. Iunio, Iulio. Fr.

incubacea. 12. S. foliis integerrimis lanceolatis: ſubtus villoſis nitidis, ſtipulis ovatis acutis. *Sp. pl. p.* 1447.

Salix pumila anguſtifolia ſecunda. *Cluſ. hiſt. p.* 86.

In pratis paluſtribus Auſtriæ, Pannoniæ. Fl. Martio, Aprili. Fr.

Iacquinii. 13. S. foliis integerrimis lanceolatis obovatisque ciliatis: ſubtus villoſis.

Salix fusca. *Iacqu. auſtr. t.* 409.

In alpibus. Fl. Iulio. Fr.

Salix fusca Iacquinii non eſt fusca Linn. ſed nova ſpecies. Willd. baumz. p. 452.

roſmarinifolia. 14. S. foliis integerrimis lanceolato-linearibus ſtrictis ſeſſilibus ſubtus tomentoſis. *Sp. pl. p.* 1448.

In Auſtriæ, Pannoniæ arenoſis ad fluvios, torrentes. Fl. Majo. Fr.

**** *Foliis ſubſerratis, villoſis.*

caprea. 15. S. foliis ovatis rugoſis: ſubtus tomentoſis undatis, ſuperne denticulatis. *Sp. pl. p.* 1448. *Hoffm. ſal.* 1. *t.* 3. *f.* 1. 2. *t.* 5. *f.* 4. *t.* 21. *f.* 1. *Willd baumz. p.* 344.

Salix foliis ovatis, rugoſis, ſubtus reticulatis, tomentoſis, julis ovatis, ſtipulis amplexicaulibus ſerratis. *Hall. hiſt. n.* 1653.

In ſylvaticis. Fl. Martio. Arb.

cinerea. 16. S. foliis ſubſerratis, oblongo-ovatis: ſubtus ſubvilloſis, ſtipulis dimidiato-cordatis. *Sp. pl. p.* 1449.

Salix aurita. *Hoffm. ſal.* 1. *t.* 4. *f.* 1. 2. *t.* 5. *f.* 3. *auct. Cl. Willdenow.*

Ad fluvios, in paluſtribus, in ſylvis montanis, & demiſſis. Fl. Aprili, Majo. Fr.

acuminata. 17. S. foliis ovato-oblongis, ſubtus tomentoſis: ſuperioribus integris, inferioribus crenatis. *Hoffm. ſal.* 1. *t.* 6. *f.* 1. *t.* 22. *f.* 2.

Salix foliis oblongo-ovatis acuminatis apice obtuſe dentatis ſubtus rugoſis tomentoſis. *Willd. baumz. p.* 346.

Ad fluvios, lacus, in pratis paluſtribus. Fl. Martio, Aprili. Fr.

viminalis. 18. S. foliis ſubintegerrimis lanceolato-linearibus longiſſimis acutis ſubtus ſericeis, ramis virgatis. *Sp. pl. p.* 1448. *Hoffm. ſal.* 1. *t.* 2. *f.* 1. 2. *t.* 5. *f.* 2. *t.* 21. *f.* 2.

Salix foliis prælongis, obſcure dentatis, ſubtus tomentoſis & albicantibus. *Hall. hiſt. n.* 1641.

Ad fluvios, in paluſtribus. Fl. Apili, Majo. Fr.

alba. 19. S. foliis lanceolatis acuminatis ſerratis utrinque pubescentibus: ſerraturis infimis glanduloſis. *Sp. pl. p.* 1449. *Hoffm. ſal.* 1. *t.* 7. *f.* 1. *t.* 8. *f.* 2. *t.* 24. *f.* 3. *Scop. carn. tom.* 2. *p.* 258.

Ad pagos, fluvios, ſepes, in ſylvaticis. Fl. Majo. Arb.

TRIANDRIA.

EMPETRUM. Masc. Calyx tripartitus. Corolla tripetala. Stamina longa.

Fem. Calyx & corolla maris. Styli novem. Baccca enneaſperma.

nigrum. 1. E. procumbens. *Sp. pl. p.* 1450. *Fl. dan. t.* 975.

Empetrum procumbens, foliis ovato-lanceolatis, obtuſis. *Hall. hiſt. n.* 1605.

Erica coris folio undecima. *Clus. hiſt. p.* 45.

Erica baccifera procumbens nigra. *Bauh. pin.* 486.

In alpibus. Fl. Iunio, Iulio. Fr.

OSYRIS. Maſc. Calyx trifidus: Corolla nulla.

Fem. Calyx & corolla maris. Stylus nullus. Stigma ſubrotundum. Drupa unilocularis.

alba. 1. O. foliis linearibus. *Syſt. veget. p.* 881. *Scop. carn. tom.* 2. *p.* 260.

Caſia quorundam. *Clus. hiſt. p.* 91.

Oſyris frutescens baccifera. *Bauh. pin.* 212. *Zann. iſt. p.* 53. *t.* 83.

Ad ſepes, maceries Litoralis. Fl. Aprili, Majo. Fr..

TETRANDRIA.

VISCUM. Maſc. Calyx quadripartitus. Corolla nulla. Filamenta nulla. Antheræ calyci adnatæ.

Fem. Calyx tetraphyllus, ſuperus. Corolla nulla. Stylus nullus. Bacca monoſſperma. Semen cordatum.

album. 1. V. foliis lanceolatis obtuſis, caule dichotomo, ſpicis axillaribus. *Sp. pl. p.* 1451. *Miller illuſtr. ic. Scop. carn. tom.* 2. *p.* 261. *Willd. baumz. p.* 406.

Viscum foliis lanceolatis obtuſis, caule dichotomo, ſpicis alaribus. *Hall. hiſt. n.* 1609.

Le Gui de Chêne. Regn. bot.

Arboribus adnaſcitur. Fl. Februario, Martio. Fr.

HIPPOPHAË. Maſc. Calyx bipartitus. Corolla nulla.

Fem. Calyx bifidus. Corolla nulla. Stylus unicus. Bacca monoſperma.

Rhamnoides. 1. H. foliis lanceolatis. *Sp. pl. p.* 1452. *Fl. dan. t.* 265. *Pallas roſſic. t.* 68. *Willd. baumz. p.* 142.

Hippophaë foliis linearibus, ſubtus rubiginoſis. *Hall. hiſt. n.* 1603.

Oſyris Rhamnoides. *Scop. carn. tom.* 2. *p.* 261.

Rhamnus ſecundus. *Clus. hiſt. p.* 110.

In arenoſis ad fluvios, & ad litus maris adriatici. Fl. primo vere. Arb.

PENTANDRIA.

PISTACIA. Maſc. Calyx communis amentum. Calyx proprius quinquefidus. Corolla nulla.

Fem. Calyx communis nullus. Proprius trifidus. Corolla nulla. Styli duo. Drupa monoſperma.

Terebinthus. 1. P. foliis impari pinnatis: foliolis ovato-lanceolatis. *Sp. pl. p.* 1455. *Scop. carn. tom.* 2. *p.* 262.

Terebinthus. *Clus. hiſt. p.* 15.

Le Terebinthe. Regn. bot.

In Litorali. Fl. Aprili, Majo. Arb.

Semina, quæ ob gravem terebinthinæ odorem ſpernunt incolæ, libentiſſime vorat Loxia Coccothrauſtes.

Lentiscus. 2. P. foliis abrupte pinnatis: foliolis lanceolatis. *Sp. pl. p.* 1455.

Lentiscus. *Clus. hiſt. p.* 14.

Le Lentisque. Regn. bot.

In Iſtria, & inſulis venetis. Fl. Iunio. Fr.

CANNABIS. Masc. Calyx quinquepartitus. Corolla nulla.

Fem. Calyx monophyllus, integer, latere hians. Corolla nulla. Styli duo. Nux bivalvis, intra calycem clauſum.

ſativa. 1. Cannabis. *Sp. pl. p.* 1457. *Hall. hiſt. n.* 1616.

Le Chanvre. Regn. bot.

Ad pagos, in ruderatis. Fl. æſtate.

HUMULUS. Masc. Calyx pentaphyllus. Corolla nulla.

Fem. Calyx monophyllus, oblique patens, integer. Corolla nulla. Styli duo. Semen unicum intra calycem foliatum.

Lupulus. 1. Humulus. *Sp. pl. p.* 1457. *Miller illuſtr. ic. Scop. carn. tom.* 2. *p.* 263. *Hall. hiſt. n.* 1618. *Clus. hiſt. p.* 126.

Le Houblon. Regn. bot.

Ad ſepes, in ſylvis. Fl. Iunio. Per.

HEXANDRIA.

TAMUS. Masc. Calyx ſexpartitus. Corolla nulla.

Fem. Calyx & corolla maris. Stylus trifidus. Bacca trilocularis, infera. Semina duo.

communis. 1. T. foliis cordatis indiviſis. *Sp. pl. p.* 1458. *Miller iluſtr. ic. Scop. carn. tom.* 2. *p.* 263. *Hall. hiſt. n.* 1620.

Le Sceau de notre Dame. Regn. bot.

In Croatia, Carniolia, Litorali ad ſepes, in nemoribus. Fl. Martio, Aprili. Per.

Semina a duobus ad quatuor.

SMILAX. Maſc. Calyx hexaphyllus. Corolla nulla.

Fem. Calyx & corolla maris. Styli tres. Bacca trilocularis. Semina duo.

aſpera. 1. S. caule aculeato angulato, foliis dentato-aculeatis cordatis novemnerviis. *Sp. pl. p.* 1458. *Scop. carn. tom.* 2. *p.* 264.

Smilax aſpera rutilo fructu. *Cluſ. hiſt. p.* 112.

In Litorali in aſperis, ad ſepes. Fl. Majo. Per.

OCTANDRIA.

POPULUS. Maſc. Amentum conſtans ſquamis unifloris, margine laceris. Corolla nulla. Nectarium turbinatum, obliquum, integrum.

Fem. Calyx, corolla, nectarium maris. Stigma quadrifidum. Capſula bilocularis. Semina multa, pappoſa.

alba. 1. P. foliis ſubrotundis dentato-angulatis ſubtus tomentoſis. *Sp. pl. p.* 1463. *Scop. carn. tom.* 2. *p.* 265.

Populus foliis anguloſis, ſubtus tomentoſis, ſubrotundis, dentato-angulatis. *Hall. hiſt. n.* 1634.

In ſylvis, ad fluvios. Fl. primo vere. Arb.

tremula. 2. P. foliis ſubrotundis dentato-angulatis utrinque glabris. *Sp. pl. p.* 1464. *Scop. carn. tom.* 2. *p.* 265. *Willd. baumz. p.* 228. *Reitter und Abel abbild. t.* 9.

Populus foliis glabris, orbiculatis, acuminatis, laxe ſerratis. *Hall. hiſt. n.* 1633.

Eſpe. Cram. t. 13.

In ſylvis. Fl. primo vere. Arb.

nigra. 3. P. foliis deltoidibus acuminatis ſerratis. *Sp. pl. p.* 1464. *Scop. carn. tom.* 2. *p.* 265.

Populus foliis glabris cordato-rhomboideis, ſerratis. *Hall. hiſt. n.* 1632.

Le Peuplier noir. Regn. bot.

Ad fluvios, pagos. Fl. Aprili. Arb.

RHODIOLA. Maſc. Calyx quadripartitus. Corolla tetrapetala. Nectaria quatuor. Germina quatuor. Styli & ſtigmata obſoleta. Pericarpium abortiens.

Fem. Calyx & nectaria maris. Corolla nulla. Germina quatuor, in ſtylos ſimplices deſinentia. Stigmata obtuſa. Capſulæ quatuor, polyſpermæ.

roſea. 1. Rhodiola. *Sp. pl. p.* 1465. *Fl. dan. t.* 483.

Sedum ſexu diſtinctum, foliis ſerratis, umbellis denſiſſimis. *Hall. hiſt. n.* 953.

Sedum roſeum. *Scop. carn. tom.* 1. *p.* 326.

Rhodia radix. *Clus. hiſt. p. LXV.*

In alpibus. Fl. Iulio. Per.

ENNEANDRIA.

MERCURIALIS. Maſc. Calyx tripartitus. Corolla nulla. Stamina novem vel duodecim. Antheræ globoſæ, didymæ.

Fem. Calyx & corolla maris. Styli duo. Capſula dicocca, bilocularis, monoſperma.

perennis. 1. M. caule ſimpliciſſimo, foliis ſcabris. *Sp. pl. p.* 1465. *Fl. dan. t.* 400. *Curtis lond.*

Mercurialis caule perenni, indiviſo, foliis hirſutis, ovato-lanceolatis. *Hall. hiſt. n.* 1601.

Mercurialis Cynocrampe. *Scop. carn. tom.* 2. *p.* 266.

In ſylvaticis. Fl. primo vere.

annua. 2. M. caule brachiato, foliis glabris, floribus ſpicatis. *Sp. pl. p.* 1465. *Curtis lond. Scop. carn. tom.* 2. *p.* 267.

Mercurialis caule annuo, brachiato, foliis glabris, conjugatis, ovato-lanceolatis. *Hall. hiſt. n.* 1600.

La Mercuriale. Regn. bot.

In cultis. Fl. a Iunio ad ſerum Autumnum.

HYDROCHARIS. Maſc. Spatha diphylla. Calyx trifidus. Corolla tripetala. Filamenta tria interiora ſtylifera.

Fem. Calyx & corolla maris. Styli ſex. Capſula ſexlocularis, polyſperma, infera.

Morſus ranæ. 1. Hydrocharis. *Sp. pl. p.* 1466. *Fl. dan. t.* 878. *Curtis lond. Hall. hiſt. n.* 1068.

In Auſtriæ, Pannoniæ aquis ſtagnantibus. Fl. Iunio, Iulio. Per.

POLYANDRIA.

STRATIOTES. Maſc. Spatha diphylla. Calyx tripartitus. Corolla tripetala. Nectaria viginti. Stamina duodecim.

Fem. Spatha, Calyx, corolla, nectaria maris. Germen unicum. Styli ſex, bipartiti. Bacca.

Aloides. 1. S. foliis enſiformi-triangulis ciliato-aculeatis. *Sp. pl. p.* 754. *Fl. dan. t.* 337.

In Auſtria in aquis ſtagnantibus ad Danubii ripas. Fl. æſtate. Per.

MONADELPHIA.

IUNIPERUS. Maſc. Amentum. Calycis loco ſquama. Corolla nulla. Stamina tria.

Fem. Calyx tripartitus. Corolla tripetala. Styli tres. Bacca triſperma.

Sabina. 1. I. foliis oppoſitis erectis decurrentibus: oppoſitionibus pyxidatis. *Sp. pl. p.* 1472. *Scop. carn. tom.* 2. *p.* 268.

Iuniperus foliis cauli adpreſſis, lanceolatis, alterne conjugatis. *Hall. hiſt. n.* 1662.

La Sabine. Regn. bot.

In Iſtria, & in inſulis venetis. Fl. primo vere. Fr.

communis. 2. I. foliis ternis patentibus mucronatis bacca longioribus. *Sp. pl. p.* 1470. *Scop. carn. tom.* 2. *p.* 268. *Pallas roſſic. t.* 54. *f. b. c.*

Iuniperus foliis convexo-concavis, ariſtatis, baccis alaribus. *Hall. hiſt. n.* 1661.

Iuniperus vulgaris. *Clus. hiſt. p.* 38.

Le Genevrier. Regn. bot.

In ſylvaticis, montanis, & alpinis. Fl. primo vere. Fr.

Frutex in alpibus creſcens habet ramos proſtratos, folia breviora, incurva, tenuiora, minus pungentia.

Oxycedrus. 3. I. foliis ternis patentibus mucronatis, bacca brevioribus. *Sp. pl. p.* 1470. *Scop. carn. tom.* 2. *p.* 268. *Du Hamel arb. t.* 128.

Oxycedrus. *Clus. hiſt. p.* 39.

Iuniperus major, bacca ruſeſcente. *Bauh. pin.* 489.

In aſperis totius Litoralis. Fl. Majo. Fr.

POLYGAMIA.

MONOECIA.

VERATRUM. Herm. Calyx nullus. Corolla hexapetala. Stamina ſex. Piſtilla tria. Capſulæ tres, polyſpermæ.

Maſc. Piſtillorum rudimenta.

album. 1. V. racemo ſupradecompoſito, corollis erectis. *Sp. pl. p.* 1479. *Iacqu. auſtr. t.* 335. *Scop. carn. tom.* 2. *p.* 272.

Veratrum ſpica paniculata, floribus ſexu diſtinctis. *Hall. hiſt. n.* 1204.

Elleborus albus exalbido flore. *Clus. hiſt. p.* 274.

Le Veratre blanc. Bulliard. herb. t. 155.

In alpinis, & ſubalpinis. In Auſtria & in ſimilibus locis, & in pratis planitiei prope *Erla, Lachſenburg*, *Himberg*, *Gaden*, alibique. Fl. a Iunio ad Auguſtum. Per.

nigrum. 2. V. racemo compoſito, corollis patentiſſimis. *Sp. pl. p.* 1479. *Iacqu. auſtr. t.* 336. *Scop. carn. tom.* 2. *p.* 272.

Elleborus albus alter, flore ex purpura nigricante. *Clus. hiſt. p.* 274.

Le Veratre noir. Bulliard herb. t. 149.

In montoſis ſylvaticis Auſtriæ, Carnioliæ calidioris. Fl. Iulio. Per.

ANDROPOGON. Herm. Calyx. Gluma uniflora. Corolla gluma baſi ariſtata. Stamina tria. Styli duo. Semen unicum.

Maſc. Calyx, Corolla, Stamina prioris.

Gryllus. 1. A. paniculæ pedunculis ſimpliciſſimis trifloris: floſculo hermaphrodito ſeſſili ariſtato ciliato baſi barbato. *Sp. pl. p.* 1480. *Scop. carn. tom.* 2. *p.* 273.

Phœnix ſpica laxa, locuſtis longe petiolatis, quadrifloris, duobus floribus ſeſſilibus, duobus petiolatis. *Hall. hiſt. n.* 1412.

Gramen ſparteum feſtuceum, ſeu Ægilops ſpartea villoſa. *Barr. ic.* 1237. *t.* 18. *n.* 2. *Zann. iſt. p.* 123. *t.* 135.

Ægylops bromoides, juba purpuraſcente. *Bauh. hiſt.* 2. *p.* 436. *Scheuch. agr.* 267. *t.* 6. *f.* 1.

In pratis, aliisque herbidis locis Carnioliæ calidioris, Forijulii, Litoralis. Fl. Iunio, Iulio. Per.

diſtachyon. 2. A. ſpicis binis terminalibus, culmo indiviſo. *Sp. pl. p.* 1481. *Iacqu. collect. tom.* 1. *p.* 165. *ic. t.* 630.

Andropogon ſpicis geminis bifloris, alterne muticis. *Hall. hiſt. n.* 1415.

Gramen dactylon, ſpica gemina. *Tourn. inſt.* 521. *Scheuch. agr.* 95. *Zann. t.* 36.

In herbidis agri fluminenſis, inſularumque venetarum. Fl. Iunio, Iulio. Per.

Iſchæmum. 3. A. ſpicis digitatis plurimis, floſculis ſeſſilibus; ariſtato muticoque, pedicellis lanatis. *Sp. pl. p.* 1483. *Iacqu. auſtr. t.* 384. *Scop. carn. tom.* 2. *p.* 274.

Andropogon ſpicis villoſis ſeptenis, locuſtis bifloris, floribus alternis, ſeſſilibus. *Hall. hiſt. n.* 1414.

In collibus ſiccis, aggeribus, ad vias. Fl. Auguſto, Septembri. Per.

HOLCUS. Herm. Calyx gluma uni- ſeu biflora. Corolla gluma ariſtata. Stamina tria. Styli duo. Semen unicum.

Maſc. Calyx gluma bivalvis. Corolla nulla. Stamina tria.

halepenſis. 1. H. glumis glabris, floribus hermaphroditis muticis, femineo ariſtato. *Sp. pl. p.* 1485. *Schreb. gr. t.* 18.

Andropogon arundinaceum. *Scop. carn. tom.* 2. *p.* 274.

Milium arundinaceum perenne minus, ſemine oblongo, nigro. *Mont. bon. Zann. iſt. p.* 126. *n.* 23.

Gramen arundinaceum, paniculatum; locuſtis partim muticis, partim ariſtatis. *Scheuch. agr.* 509. *t.* 11. *f.* 12.

In vineis, ad ſepes Litoralis, Forijulii. Fl. Iulio. Per.

lanatus. 2. H. glumis bifloris villoſis: hermaphrodito mutico, maſculo ariſta recurva. *Sp. pl. p.* 1485. *Schreb. gr. t.* 20. *f.* 1. *Leers herb. p.* 221. *t.* 7. *f.* 6. *Scop. carn. tom.* 2. *p.* 275. *Curtis lond.*

Avena diantha, floribus ovatis, perfecto mutico, imperfecto ariſtato. *Hall. hiſt. n.* 1484.

In pratis humidis. Fl. Iunio. Per.

odoratus. 3. H. glumis trifloris muticis glabris: hermaphrodito diandro. *Sp. pl. p.* 1485. *Fl. dan. t.* 963.

Avena locuſtis trianthis, paniculatis, hiantibus, floſculo tertio imperfecto. *Hall. hiſt. n.* 1496.

Gramen paniculatum odoratum. *Bauh. pin.* 7. *Scheuch. agr.* 236. *t.* 4. *f.* 26.

Ad margines ſylvarum, in nemoribus Auſtriae. Fl. Martio, Aprili. Per.

CENCHRUS. Involucrum laciniatum, echinatum, biflorum. Calyx gluma biflora: altero maſculo, altero hermaphrodito.

Herm. Corollæ gluma mutica. Stamina tria. Semen unicum.

Masc. Corollæ gluma mutica. Stamina tria.

racemoſus 1. C. panicula ſpicata, glumis muricatis ſetis ciliaribus. *Sp. pl. p.* 1487. *Schreb. gr. t.* 4.

Tragus. *Hall. hiſt. n.* 1413.

Gramen caninum, maritimum aſperum, ſive ſpica echinata. *Bauh. pin.* 2. *Scheuch agr.* 76. *t.* 2. *f.* 7.

In Litoralis, Forijulii aridis. Fl. Iunio, Iulio.

ÆGILOPS. Herm. Calyx gluma ſubtriflora, cartilaginea. Corollæ gluma terminata triplici ariſta. Stamina tria. Styli duo. Semen unicum.

Masc. Calyx & corolla prioris. Stamina tria.

ovata. 1. Æ. ſpica ariſtata, calycibus omnibus triariſtatis. *Sp. pl. p.* 1489.

Phleum Ægilops. *Scop. carn. tom.* 1. *p.* 55.

Gramen ſpicatum durioribus & craſſioribus locuſtis, ſpica brevi. *Tourn. inſt.* 519. *Scheuch. agr.* 11.

In vineis, aſperisque Litoralis, Iſtriæ. Fl. primo vere.

VAILLANTIA. Herm. Calyx nullus. Corolla quadripartita. Stamina quatuor. Stylus bifidus. Semen unicum.

Maſc. Calyx nullus. Corolla tri- ſeu quadripartita. Stamina quatuor vel tria. Piſtillum obſoletum.

cruciata. 1. V. floribus maſculis quadrifidis, pedunculis diphyllis. *Sp. pl. p.* 1491.

Galium foliis quaternis hirſutis, petiolis octifloris. *Hall. hiſt. n.* 709.

Galium Cruciata. *Scop. carn. tom.* 1. *p.* 100.

Cruciata. *Dod. purg. p.* 107. *Blakw. herb. t.* 76.

Ad ſepes, inter frutices, & alibi. Fl. primo vere. Per.

glabra. 2. V. floribus maſculis quadrifidis, pedunculis dichotomis aphyllis, foliis ovalibus ciliatis. *Sp. pl. p.* 1491.

Galium foliis quaternis, glabris, ovato-lanceolatis, racemis breviſſimis, reflexis, æqualibus. *Hall. nom. n.* 720.

Galium vernum. *Scop. carn. tom.* 1. *p.* 99. *t.* 2.

In Carnioliæ, Litoralis herbidis. Fl. primo vere. Per.

PARIETARIA. Herm. Calyx quadrifidus. Corolla nulla. Stamina quatuor. Stylus unus. Semen unicum, ſuperum, elongatum.

Fem. Calyx quadrifidus. Corolla nulla. Stamen nullum. Stylus & ſemen prioris.

officinalis. 1. P. foliis lanceolato-ovatis, pedunculis dichotomis, calycibus diphyllis. *Syſt. veget. p.* 908. *Fl. dan. t.* 521. *Scop. carn. tom.* 2. *p.* 277.

Parietaria foliis hirſutis, elliptico - lanceolatis. *Hall. hiſt. n.* 1612.

La Parietaire. Regn. bot.

In muris, ſylvis, ruderatis. Fl. Iunio. Per.

ATRIPLEX. Herm. Calyx pentaphyllus. Corolla nulla. Stamina quinque. Stylus bipartitus. Semen unicum, depreſſum.

Fem. Calyx diphyllus: foliolis maturo ſemine magnis. Corolla nulla. Stamina nulla. Stylus bipartitus. Semen unicum, compreſſum.

Portulacoides. 1. A. caule fruticoſo, foliis obovatis. *Sp. pl. p.* 1493. *Scop. carn. tom.* 2. *p.* 278.

Halimus ſeu Portulaca marina. *Bauh. pin.* 120. *Zann. iſt. p.* 32. *t.* 127.

In Litorali ad mare. Fl. Auguſto.

hortenſis. 2. A. caule erecto herbaceo, foliis triangularibus. *Sp. pl. p.* 1493.

Atriplex hortenſis alba ſeu pallide virens. *Bauh. pin.* 119.

In ruderatis, ad hortos, pagos, vias. Fl. Iulio.

laciniata. 3. A. caule herbaceo, foliis deltoidibus dentatis ſubtus argenteis. *Sp. pl. p.* 1494. *Scop. carn. tom.* 2. *p.* 278.

In Auſtriæ, Litoralis ruderatis, ad vias. Fl. Iulio.

haſtata. 4. A. caule herbaceo: valvulis femineis magnis deltoidibus ſinuatis. *Sp. pl. p.* 1494. *Curtis lond.*

Atriplex foliis triangularibus, baſi producta, valvulis triangularibus, ſubaſperis. *Hall. hiſt. n.* 1617.

Ad fimeta, in paluſtribus. Fl. Iulio.

M m

patula. 5. A. caule herbaceo patulo, foliis ſubdeltoideo-lanceolatis, calycibus ſeminum disco-dentatis. *Sp. pl. p.* 1494. *Scop. carn. tom.* 2. *p.* 279.

Atriplex foliis imis hamatis, reliquis lingulatis, valvulis ſeminalibus dentatis. *Hall. hiſt. n.* 1616.

Spear Orrach. Pet. t. 7.

In cultis, ad vias, hortos. Fl. Iulio, Auguſto.

litoralis. 6. A. caule herbaceo erecto, foliis omnibus linearibus integerrimis. *Sp. pl. p.* 1494.

Ad litus maris adriatici. Fl. Auguſto.

ACER. Herm. Calyx quinquefidus. Corolla pentapetala. Stamina octo. Piſtillum unicum. Capſulæ duæ ſeu tres, monoſpermæ, ala terminatæ.

Masc. Calyx, corolla, ſtamina prioris.

tataricum. 1. A. foliis cordatis indiviſis ſerratis: lobis obſoletis, floribus racemoſis. *Sp. pl. p.* 1495. *Pallas roſſic. t.* 4.

In Pannonia. Fl. Majo. Arb.

Pſeudo-Platanus. 2. A. foliis quinquelobis inæqualiter ſerratis, floribus racemoſis. *Sp. pl. p.* 1495. *Willd. baumz. p.* 2. *Reitter und Abel abbild. t.* 13. *Du Hamel. arb.* 1. *t.* 36. *f.* 1.

Acer foliis quinquelobis, acute ſerratis, racemis pendulis. *Hall. hiſt. n.* 1029.

Acer latifolium. *Clus. hiſt. p.* 10.

In ſylvis. Fl. Majo, Iunio. Arb.

Platanoides. 3. A. foliis quinquelobis acuminatis acute dentatis glabris, floribus corymboſis. *Sp. pl. p.* 1496. *Miller. ic. t.* 8. *f.* 1. *Willd. baumz. p.* 6. *Reitter und Abel abbild. t.* 14.

Acer foliis quinquelobis, dentibus acutis, intervallis lunatis, racemis erectis. *Hall. hiſt. n.* 1029.

In ſylvis. Fl. Aprili, Majo. Arb.

campeſtre. 4. A. foliis lobatis obtuſis emarginatis. *Sp. pl. p.* 1497. *Scop. carn. tom.* 2. *p.* 280. *Willd. baumz. p.* 8. *Reitter und Abel abbild. t.* 25.

Acer foliis ſemitrilobis obtuſis, lobis lateralibus emarginatis. *Hall. hiſt. n.* 1029.

In ſylvis, ad ſepes. Fl. Aprili. Arb.

monſpeſſulanum. 5. A. foliis trilobis integerrimis glabris annuis. *Syſt. veget. p.* 912. *Scop. carn. tom.* 2. *p.* 280. *Du Hamel. arb.* 1. *p.* 28. *t.* 10. *f.* 8.

In Litorali, Iſtria. Fl. Aprili. Arb.

Flores ab apibus frequentantur. Loxia coccothrauſtes vorat fructus.

CELTIS. Herm. Calyx quinquepartitus. Corolla nulla. Stamina quinque. Styli duo. Drupa monoſperma.

Maſc. Calyx ſexpartitus. Corolla nulla. Stamina ſex.

auſtralis. 1. C. foliis ovato-lanceolatis. *Sp. pl. p.* 1478. *Scop. carn. tom.* 2. *p.* 271. *Delic. ins.* 2. *t.* 18.

Celtis foliis ſcabris ſerratis, longiſſime lanceolatis. *Hall. hiſt. n.* 1619.

In Litorali, Iſtria, Forojulio. Fl. Aprili. Arb.

DIOECIA.

FRAXINUS. Herm. Calyx vel nullus ſeu quadripartitus. Corolla vel nulla vel tetrapetala. Stamina duo. Piſtillum unum. Semen unicum, lanceolatum.

Fem. Piſtillum unum, lanceolatum.

excelſior. 1. F. foliis ſerratis, floribus apetalis. *Sp. pl. p.* 1509. *Fl. dan. t.* 969.

Fraxinus apetala, foliis pinnatis ſerratis. *Hall. hiſt. n.* 528.

Fraxinus Ornus. *Scop. carn. tom.* 2. *p.* 281.

In ſylvis, ad vias, pagos. Fl. Aprili, Majo. Arb.

Folia in Tyroli dant vaccis; in Styria & Pannonia ovibus.

Ornus. 2. F. foliolis ſerratis, floribus corollatis. *Sp. pl. p.* 1510. *Mich. gen. p.* 225. *t.* 107.

Fraxinus florifera. *Scop. carn. tom.* 2. *p.* 282.

In Litorali, Iſtria. Fl. Majo. Fr.

TRIOECIA.

FICUS. Receptaculum commune turbinatum, carnoſum, connivens, occultans flosculos vel in eodem vel diſtincto.

Masc. Calyx tripartitus. Corolla nulla. Stamina tria.

Fem. Calyx quinquepartitus. Corolla nulla. Piſtillum unum. Semen unicum.

Carica. 1. F. foliis palmatis. *Sp. pl. p.* 1513. *Scop. carn. tom.* 2. *p.* 283. *Hall. hiſt. n.* 1607.

Le Figuier. Regn. bot.

In maceriis, muris vetuſtis Litoralis, Forijulii. Arb.

CRYPTOGAMIA.

MISCELLANEÆ.

EQUISETUM. Fructificationes in clavam ovato-oblongam digeſtæ, verticillatæ, ſtipitatæ.

ſylvaticum. 1. E. caule ſpicato, frondibus compoſitis. *Sp. pl. p.* 1516. *Bolt. fil. t.* 33. *Scop. carn. tom.* 2. *p.* 285. *Hoffm. fl.* 2. *p.* 3.

Equiſetum foliis repetito-ramoſis, vaginis læviſſimis. *Hall. hiſt. n.* 1680.

In ſylvis montanis, & ſubalpinis. Fl. Aprili, Majo. Per.

arvenſe. 2. E. ſcapo fructificante nudo, ſterili frondoſo. *Sp. pl. p.* 1516. *Bolt. fil. t.* 34. *Curtis lond. Scop. carn. tom.* 2. *p.* 286. *Hoffm. fl.* 2. *p.* 3.

Equiſetum caule florigero nudo, ſterili verticillato, radiorum duodecim. *Hall. hiſt. n.* 1676.

In arvis, pratis, ad foſſas, fluvios, alibique. Fl. Martio, Aprili. Per.

paluſtre. 3. E. caule angulato, frondibus ſimplicibus. *Sp. pl. p.* 1516. *Bolt. fil. t.* 35. *Scop. carn. tom.* 2. *p.* 286. *Hoffm. fl.* 2. *p.* 3.

Equiſetum caule ſulcato, ramis multifloris, foliis indiviſis. *Hall. hiſt. n.* 1677.

In paludoſis, foſſis. Fl. a Majo ad Iulium. Per.

fluviatile. 4. E. caule ſtriato frondibus ſubſimplicibus. *Sp. pl. p.* 1517. *Scop. carn. tom.* 2. *p.* 286. *Hoffm. fl.* 2. *p.* 2.

In foſſis, ad fluviorum ripas. Fl. Majo, Iunio. Per.

limoſum. 5. E. caule ſubnudo lævi. *Sp. pl. p.* 1517. *Bolt. fil. t.* 38. *Hoffm. fl.* 2. *p.* 2.

In paluſtribus, aquis ſtagnantibus. Fl. Majo, Iunio. Per.

hyemale, 6. E. caule nudo ſcabro, baſi ſubramoſo. *Sp. pl. p.* 1517. *Bolt. fil. t.* 39. *Hoffm. fl.* 2. *p.* 2.

Ad fluvios, in uliginoſis, alibique. Fl. a Majo ad Auguſtum. Per.

LYCOPODIUM. Fructificationes in axillis foliorum, vel ſquamarum, in ſpicas imbricatas digeſtarum, ſeſſiles.

clavatum, 1. L. foliis ſparſis filamentoſis, ſpicis teretibus pedunculatis geminis. *Sp. pl. p.* 1564. *Fl. dan. t.* 126. *Weis crypt. p.* 269. *Hoffm. fl.* 2. *p.* 15.

Lycopodium ramis prolixis, ſpicis petiolatis gemellis, foliis linearibus pilo ariſtatis. *Hall. hiſt. n.* 1722.

Lycopodium officinale. *Neck. meth. p.* 150.

Lycopodium vulgare piloſum, anfragoſum & repens. *Dill. muſc.* 441. *t.* 58. *f.* 1.

Le Pied de Loup. Regn. bot.

In ſylvis. Fl. Iulio, Auguſto. Per.

Selaginoides, 2. L. foliis ſparſis ciliatis lanceolatis, ſpicis ſolitariis terminalibus folioſis. *Sp. pl. p.* 1565. *Fl. dan. t.* 70. *Scop. carn. tom.* 2. *p.* 301. *Neck. meth. p.* 148. *Hoffm. fl.* 2. *p.* 16.

Lycopodium ſpicis ſeſſilibus, foliis ovato-lanceolatis, ſerratis, confertis. *Hall. hiſt. n.* 1717.

Selaginoides foliis ſpinoſis. *Dill. muſc.* 460. *t.* 68. *f.* 1.

In alpinis, & ſubalpinis. Fl. æſtate. Per.

Selago, 3. L. foliis ſparſis octifariis, caule dichotomo erecto faſtigiato, floribus ſparſis. *Sp. pl. p.* 1565. *Fl. dan. t.* 104. *Scop. carn. tom.* 2. *p.* 302. *Neck. meth. p.* 148. *Hoffm. fl.* 2. *p.* 16.

Lycopodium ſpica ſeſſili, foliis lateralibus lanceolatis, confertis. *Hall. hiſt. n.* 1716.

Selago vulgaris abietis rubræ facie. *Dill. musc.* 435. *t.* 56. *f.* 1.

In ſaxoſis, rupeſtribus, herbidis alpinis, & ſubalpinis. Fl. Auguſto, Septembri. Per.

annotinum. 4. L. foliis ſparſis quinquefariis ſubſerratis, ſurculis annotino-articulatis, ſpicis terminalibus glabris erectis. *Sp. pl. p.* 1566. *Fl. dan. t.* 127. *Scop. carn. tom.* 2. *p.* 303. *Neck. meth. p.* 146. *Hoffm. fl.* 2. *p.* 15.

Lycopodium ramis prolixis, ſpicis ſeſſilibus, foliis linearibus, ſubſerratis. *Hall. hiſt. n.* 1720.

Lycopodium elatius juniperinum, clavis ſingularibus ſine pediculis. *Dill. muſc.* 455. *t.* 63. *f.* 9.

In ſylvis montanis, & ſubalpinis. Fl. æſtate. Per.

alpinum. 5. L. foliis quadrifariam imbricatis acutis, ſurculis erectis dichotomis, ſpicis ſeſſilibus teretibus. *Sp. pl. p.* 1567. *Fl. dan. t.* 79. *Neck. meth. p.* 147. *Hoffm. fl.* 2. *p.* 15.

Lycopodium caule repente, ramis tetragonis. *Fl. lapp. n.* 417. *t.* 11. *f.* 6.

Lycopodium ſpicis ſeſſilibus, foliis lanceolatis adpreſſis, utrinque ternatis. *Hall. hiſt. n.* 1719.

Lycopodium ſabinæ facie. *Dill. muſc.* 445. *t.* 58. *f.* 2.

In alpibus ſtyriacis, tyrolenſibus. Fl. Iulio, Auguſto. Per.

complanatum. 6. L. foliis bifariis connatis: ſuperficialibus ſolitariis, ſpicis geminis pedunculatis. *Sp. pl. p.* 1567. *Scop. carn. tom.* 2. *p.* 303. *Web. ſpic. p.* 33. *Neck. meth. p.* 145.

Lycopodium ſpicis petiolatis, quaternis, caulibus complanatis, foliis adpreſſis. *Hall. hiſt. n.* 1723.

Lycopodium digitatum, foliis arboris vitæ; ſpicis bigemellis. *Dill. muſc.* 448. *t.* 59. *f.* 3.

In ſylvis montoſis Litoralis, Forijulii, Carnioliæ. Fl. Iulio, Auguſto. Per.

helveticum. 7. L. foliis bifariis patulis: ſuperficialibus diſtichis, ſpicis geminis pedunculatis. *Sp. pl. p.* 1568. *Iacqu. auſtr. t.* 196. *Scop. carn. tom.* 2. *p.* 304. *Neck. meth. p.* 147.

Lycopodioides denticulatum pulchrum repens, ſpicis pediculis inſidentibus. *Dill. muſc.* 465. *t.* 64. *f.* 2.

In montanis, ſubalpinis, alpinis. In Auſtria & in ſimilibus locis, & in herbidis inſularum danubialium. Fl. Iulio, Auguſto. Per.

SALVINIA. Fructificationes quatuor ad novem, glomeratæ, inter radices verticillatas.

natans. 1. S. foliis oppoſitis ſimplicibus. *Sp. pl. p.* 1562. ſub Marſilea.

Salvinia vulgaris aquis innatans, foliis ſubrotundis punctatis læte virentibus. *Mich. gen. p.* 107. *t.* 58.

Lenticula paluſtris latifolia punctata. *Bauh. pin.* 362.

In aquis ſtagnantibus Forijulii, Iſtriæ.

F I L I C E S.

OPHIOGLOSSUM. Capſulæ numeroſæ, membrana in ſpicam diſticham connatæ, ſubgloboſæ, maturæ tranſverſim dehiſcentes, annulo nullo. Semina plurima.

vulgatum. 1. O. fronde ovata. *Sp. pl. p.* 1518. *Fl. dan. t.* 147. *Bolt. fil. t.* 3. *Scop. carn. tom.* 2. *p.* 287. *Hoffm. fl.* 2. *p.* 14.

Ophiogloſſum folio unico, ovato lanceolato obtuſo. *Hall. hiſt. n.* 1685.

La Langue de Serpent. Regn. bot.

In pratis humidis. Fl. Majo, Iunio. Per.

OSMUNDA. Capſulæ diſtinctæ in racemum diſpoſitæ, ſubgloboſæ, transverſim dehiscentes, annulo nullo. Semina plurima, minutiſſima.

Lunaria. 1. O. fronde pinnata, ſcapo ſolitario, racemo laterali. *Hoffm. ſl.* 2. *p.* 14. *Sp. pl. p.* 1519. *Bolt. ſil. t.* 4. *Fl. dan. t.* 18. *Scop. carn. tom.* 2. *p.* 288.

Osmunda foliis pinnatis, flabelliformibus, lunatis. *Hall. hiſt. n.* 1686.

Lunaria botrytis, minor. *Cluſ. hiſt. p. CXVIII.*

In pratis, pascuis alpinis, & ſubalpinis, Fl. Iunio, Iulio. Per.

regalis. 2. O. fronde bipinnata, racemo terminali ſupradecompoſito. *Hoffm. ſl.* 2. *p.* 15. *Sp. pl. p.* 1521. *Fl. dan. t.* 217.

In Tyroli in frigidis alpium convallibus. Fl. Iunio, Iulio. Per.

ACROSTICHUM. Fructificationes diſcum totum frondis tegentes.

ſeptentrionale. 1. A. frondibus bifurcatis laciniatis linearibus. *Hoffm. ſl.* 2. *p.* 12. ſub Asplenio. *Sp. pl. p.* 1524. *Bolt. ſil. t.* 8. *Fl. dan. t.* 60. *Scop. carn. tom.* 2. *p.* 289.

Acroſtichum caule bis bifurcato. *Hall. hiſt. n.* 1714.

In rupeſtribus, maceriis Tyrolis, Carinthiæ, Carnioliæ, Styriæ, Auſtriæ prope *Glocknitz* & in vicinis locis. Fl. æſtate. Per.

Marantæ. 2. A. frondibus ſubbipinnatis: pinnis oppoſito-coadunatis ſubtus hirſutiſſimis: baſi ſubdentatis. *Sp. pl. p.* 1527. *Hænke in Iacqu. collect. tom.* 2. *p.* 5.

Acroſtichum pinnis ſubtus pinnatis, ſquamoſis, obtuſis: baſi utrinque appendiculata. *Hall. hiſt. n.* 1715.

Polypodium Marantæ. *Hoffm. fl.* 2. *p.* 5.

Lonchitis altera Marantæ. *Cluſ. hiſt. p. CCXII.*

In rupeſtribus alpium ſtyriacarum, tyrolenſium. Fl. Iulio, Auguſto. Per.

ONOCLEA. Capſulæ ſub frondis pinnis pinnulisque recurvato - contractis.

Spicant. 1. O. frondibus ſterilibus pinnatifidis, fructificante pinnata diſticha. *Hoffm. fl.* 2. *p.* 11.

Osmunda Spicant. *Sp. pl. p.* 1522. *Fl. dan. t.* 99. *Curtis lond.*

Acroſtichum Spicant. *Sibthorp. oxon. p.* 267.

Struthiopteris Spicant. *Scop. carn. tom.* 2. *p.* 288. *Hall. hiſt. n.* 1687.

Lonchitis altera Neotheoreticorum. *Cluſ. hiſt. p. CCXIII.*

In montanis, & ſubalpinis. Fl. Iunio, Iulio. Per.

PTERIS. Capſulæ digeſtæ in lineam, ſubtus cingentem marginem frondis.

aquilina. 1. P. frondibus ſupradecompoſitis: foliis pinnatis: pinnis lanceolatis: infimis pinnatifidis; ſuperioribus minoribus. *Sp. pl. p.* 1533. *Bolt. fil. t.* 10. *Scop. carn. tom.* 2. *p.* 290.

Filix foliis triplicato - pinnatis, pinnis nervoſis integerrimis, ultimis lanceolatis. *Hall. hiſt. n.* 1688.

Le Pteris aquilin. Bulliard. herb. t. 207.

In montanis, ſubalpinis. Fl. æſtate. Per.

ASPLENIUM. Fructificationes in lineolis disci frondis ſparſis.

Scolopendrium. 1. A. frondibus ſimplicibus cordato-lingulatis integerrimis, ſtipitibus hirſutis. *Sp. pl. p.* 1537. *Curtis lond. Bolt. fil. t.* 11. *Scop. carn. tom.* 2. *p.* 291. *Hoffm. fl.* 2. *p.* 13.

Aſplenium petiolis hirſutis, folio longe lineari-lanceolato, integerrimo, poſtice cordato. *Hall. hiſt. n.* 1695.

Phyllitis vulgaris. *Clus. hiſt. p. CCXIII.*

Le Scolopendre. Regn. bot.

In montanis, ſubalpinis; Fl. Auguſto, Septembri. Per.

Ceterach. 2. A. frondibus pinnatifidis: lobis alternis confluentibus. *Sp. pl. p.* 1538. *Bolt. fil. t.* 12. *Scop. carn. tom.* 2. *p.* 291. *Hoffm. fl.* 2. *p.* 13.

Aſplenium foliis pinnatis, pinnis in baſi dilatatis, obtuſis, ſubtus muſcoſis. *Hall. hiſt. n.* 1694.

Le Ceterac. Regn. bot.

In rupeſtribus, muris Forijulii, Litoralis. Fl. æſtate, Autumno. Per.

Trichomanes. 3. A. frondibus pinnatis: pinnis ſubrotundis crenatis. *Sp. pl. p.* 1540. *Fl. dan. t.* 119. *Bolt. fil. t.* 13. *Scop. carn. tom.* 2. *p.* 292. *Hoffm. fl.* 2. *p.* 13.

Aſplenium foliis pinnatis, pinnis ovatis, crenatis. *Hall. hiſt. n.* 1693.

La Doradille Politric. Bulliard herb. t. 185.

In muris, rupibus, maceriis. Fl. toto anno. Per.

Ruta muraria. 4. A. frondibus alternatim decompoſitis: foliolis cuneiformibus crenulatis. *Sp. pl. p.* 1541. *Fl. dan. t.* 190. *Bolt. fil. t.* 16. *Scop. carn. tom.* 2. *p.* 293. *Hoffm. fl.* 2. *p.* 12.

Aſplenium fronde laxa ramoſa, ramis ſecundis trifoliis, ſuperioribus ſemitrilobis rhombeis, circumſerratis. *Hall. hiſt. n.* 1691.

La Doradille des murs. Bulliard. herb. t. 195.

Habitat & floret cum præc. Per.

Breynii. 5. A. frondibus ſubbipinnatis, foliolis oblanceolatis alternis, apice ſerrulatis. *Retzii ſcand. p.* 249. *obs. bot. faſc.* 1. *p.* 32.

Aſplenium caule pinnato, pinnis imis trifoliatis, ſuperioribus ſimplicibus dentatis. *Hall. hiſt. n.* 1690.

Aſplenium germanicum. *Weis crypt. p.* 299. *Hoffm. fl.* 2. *p.* 13.

Aſplenium alternifolium. *Wulfen in Iacqu. miſcell. tom.* 2. *p.* 51. *t.* 5. *f.* 2. *Syſt. veget. p.* 933.

Aſplenium novum germanicum Rutæ murariæ facie. *Breyn. cent. p.* 189. *f.* 97.

Habitat & floret cum Acroſticho ſeptentrionali. Per.

Adiantum nigrum. 6. A. frondibus ſubtripinnatis: foliolis alternis: pinnis lanceolatis inciſo - ſerratis. *Sp. pl. p.* 1541. *Scop. carn. tom.* 2. *p.* 292. *Hoffm. fl.* 2. *p.* 12.

Aſplenium foliis triangularibus, pinnis pinnatis, pinnulis ſemipinnatis, lobulis ovatis, ſerratis. *Hall. hiſt. n.* 1692.

Le Capillaire. Regn. bot.

In rupeſtribus Litoralis, Carnioliæ. Fl. toto anno. Per.

POLYPODIUM. Fructificationes in punctis ſubrotundis, ſparſis per discum frondis.

vulgare. 1. P. frondibus pinnatifidis: pinnis oblongis ſubſerratis obtuſis, radice ſquamata. *Sp. pl. p.* 1544. *Curtis lond. Bolt. fil. t.* 18. *Scop. carn. tom.* 2. *p.* 293. *Hall. hiſt. n.* 1696. *Hoffm. fl.* 2. *p.* 4.

Le Polypode vulgaire. Regn. bot.

In muscoſis, ſaxoſis, rupeſtribus. Fl. æſtate. Per.

Lonchitis. 2. P. frondibus pinnatis: pinnnis lunulatis ciliato-ſerratis declinatis, ſtipitibus ſtrigoſis. *Sp. pl. p.* 1548. *Fl. dan. t.* 497. *Bolt. fil. t.* 19. *Hoffm. fl.* 2. *p.* 4.

Polypodium foliis pinnatis, ciliatis, dentatis, appendiculatis. *Hall. hiſt. n.* 1711.

In alpinis, ſubalpinis. Fl. æſtate. Per.

Phegopteris. 3. P. frondibus ſub-bipinnatis: foliolis infimis reflexis: paribus pinnula quadrangulari coadunatis. *Sp. pl. p.* 1550. *Bolt. fil. t.* 20. *Hoffm. fl.* 2. *p.* 5.

Polypodium foliis pinnatis, reflexis, pinnis ovatis, hirſutis, primis cum nervo confluentibus. *Hall. hiſt. n.* 1698.

In ſylvis Styriæ, Tyrolis montanis, ſubalpinis. Fl. æſtate. Per.

criſtatum. 4. P. frondibus ſub-bipinnatis: foliolis ovato-oblongis: pinnis obtuſiuſculis apice acute ſerratis. *Sp. pl. p.* 1551. *Fl. dan. t.* 707. *Scop. carn. tom.* 2. *p.* 296. *Bolt. fil. t.* 23. *Hoffm. fl.* 2. *p.* 8.

Polypodium frondibus ſub-bipinnatis: foliolis ovato-oblongis: pinnis ovatis inciſis: laciniis apice acute ſerratis. *Müll. fridr. p.* 193. *t.* 2. *f.* 2.

In montanis, ſubalpinis. Fl. æſtate. Per.

Oreopteris. 5. P. frondibus ſubbipinnatis: pinnis alternis; pinnulis integerrimis lanceolatis obtuſiuſculis, fructificationibus marginalibus, radice ſquamoſa. *Sibthorp oxon. p.* 270. *Bolt. fil. t.* 24. *Hoffm. fl.* 2. *p.* 5.

Polypodium montanum. *Willd. prodr. p.* 291.

In Tyrolis ſylvis ſubalpinis. Fl. æſtate. Per.

Filix mas. 6. P. frondibus bipinnatis: pinnis obtuſis crenulatis, ſtipite paleaceo. *Sp. pl. p.* 1551. *Bolt. fil. t.* 24. *Scop. carn. tom.* 2. *p.* 296.

Polypodium pinnis pinnatis, obtuſis, dentatis. *Hall. hiſt. n.* 1701.

La Fougere mâle. Regn. bot.

In ſylvis. Fl. æſtate. Per.

Filix femina. 7. P. frondibus bipinnatis: pinnulis lanceolatis pinnatifidis acutis. *Sp. pl. p.* 1551. *Bolt. fil. t.* 25. *Scop. carn. tom.* 2. *p.* 295. *Hoffm. fl.* 2. *p.* 6.

Habitat & floret cum præc. Per.

aculeatum. 8. P. frondibus bipinnatis: pinnis lunulatis ciliato-dentatis, ſtipite ſtrigoſo. *Sp. pl. p.* 1552. *Bolt. fil. t.* 26. *Miller illuſtr. ic. Scop. carn. tom.* 2. *p.* 295. *Hoffm. fl.* 2. *p.* 8.

Polypodium pinnis pinnatis, ciliatis, ſerratis, appendiculatis. *Hall. hiſt. n.* 1712.

In montanis, ſubalpinis, & alpinis. Fl. æſtate. Per.

rhæticum. 9. P. frondibus bipinnatis: foliolis pinnisque remotis lanceolatis: ſerraturis acuminatis. *Sp. pl. p.* 1552. *Scop. carn. tom.* 2. *p.* 297.

Polypodium pedicularifolium. *Hoffm. fl.* 2. *p.* 10.

Filix rhætica tenuiſſime denticulata. *Bauh. hiſt.* 3. *p.* 740.

In Carnioliæ, Forijulii, Carniæ montanis, & ſubalpinis. Fl. æſtate. Per.

fragile. 10. P. frondibus bipinnatis: foliolis remotis: pinnis ſubrotundis inciſis. *Sp. pl. p.* 1553. *Fl. dan. t.* 401. *Bolt. fil. t.* 27. *Scop. carn. tom.* 2. *p.* 297. *Hoffm. fl.* 2. *p.* 9.

Polypodium pinnis pinnatis, laxe diviſis, pinnulis ſemipinnatis, lobulis ſubrotundis dentatis. *Hall. hiſt. n.* 1707.

Filix pumila ſaxatilis ſecunda. *Clus. hiſt. p. CCXII.*

In muris, rupeſtribus. Fl. æſtate. Per.

alpinum. 11. P. frondibus bipinnatis: pinnulis remote lobatis: lobulis obtuſis dentatis. *Lam. fl. franc. tom.* 1. *p.* 22. *Allion. pedem. tom.* 2. *p.* 187. *Wulfen in Iacqu. collect. tom.* 2. *p.* 171. *ic. t.* 642. *Hoffm. fl.* 2. *p.* 10.

Polypodium pinnis pinnarum pinnatis laxiſſime diviſis, lobulis obtuſis dentatis. *Hall. hiſt. n.* 1709.

In rupeſtribus alpinis, & ſubalpinis. Fl. æſtate. Per.

Dryopteris. 12. P. frondibus ſupradecompoſitis: foliolis ternis bipinnatis. *Sp. pl. p.* 1555. *Bolt. fil. t.* 28. *Hoffm. fl.* 2. *p.* 10.

Polypodium pinnis pinnatis, conjugatis, pinnulis ovatis, obtuſis, crenatis. *Hall. hiſt. n.* 1699.

Filix pumila ſaxatilis prima. *Clus. hiſt. p. CCXII.*

In muris, rupeſtribus. Fl. æſtate. Per.

montanum. 13. P. frondibus ſupradecompoſitis: foliolis alternis tripinnatis. *Lam. fl. franc. tom.* 1. *p.* 23. *Allion. pedem. tom.* 2. *p.* 287. *Hænke in Iacqu. collect. tom.* 2. *p.* 46. *Hoffm. fl.* 2. *p.* 10.

Polypodium triplicato-pinnatum, pinnulis tertiis ſemipinnatis, lobulis bifidis. *Hall. hiſt. n.* 1710.

In alpinis, & ſubalpinis. Fl. æſtate. Per.

ADIANTUM. Fructificationes congeſtæ in maculas ovales, apicibus frondium reflexis ſubjectas.

Capillus veneris. 1. A. frondibus decompoſitis: foliolis alternis: pinnis cuneiformibus lobatis pedicellatis. *Sp. pl. p.* 1558. *Scop. carn. tom.* 2. *p.* 299. *Wulfen in Iacqu. miscell. tom.* 2. *p.* 77. *t.* 7.

Adiantum petiolis ramoſis, foliis flabelliformibus lobatis. *Hall. hiſt. n.* 1713.

Le Capillaire a feuilles de coriandre. Bulliard. herb. t. 247.

In Litorali, Forojulio ad ciſternas, in rupibus humidis. Fl. æſtate. Per.

TRICHOMANES. Capſulæ e ſquama turbinata, ſolitaria, efflorescentes.

tunbrigenſe. 1. T. frondibus pinnatis: pinnis oblongis dichotomis decurrentibus dentatis. *Sp. pl. p.* 1561.

Adiantum radicoſum humi ſparſum ſeu Filicula pellucida noſtras, coriandri foliis mollicellis, globuliferum. *Pluk. alm.* 10. *t.* 3. *f.* 5. 6.

Muscus montanus italicus, adianti foliis. *Bocc. mus.* 2. *p.* 24. *t.* 2. *f.* 1.

In Carnia in paluſtribus prope *Artegna*, alibique. *I. Hoſt.* Per.

MUSCI.

PHASCUM. Capſula ovata, ſubſeſſilis, ſeu ſetæ brevi inſidens, undique clauſa, interdum cum rudimento operculi, nunquam dehiscente.

Masc. ſubdiscoidei terminales, vel gemmacei axillares.

cuspidatum. 1. P. caulescens, foliis ovatis cuspidatis patulis: ſummis erectis conniventibus. *Schreb. phasc. p.* 8. *t.* 1. *f.* 2. *ſpic. p.* 72. *Curtis lond. Web. ſpic. p.* 125. *Hoffm. fl.* 2. *p.* 19.

Phascum acaulon. *Sp. pl. p.* 1570. *Weis crypt. p.* 266.

Bryum bulbiforme. *Neck. meth. p.* 230.

Sphagnum acaulon bulbiforme majus. *Dill. musc.* 251. *t.* 32. *f.* 11.

In agris, hortis, muris vetuſtis, alibique.

ſubulatum. 2. P. caulescens, foliis lanceolato-linearibus patulis. *Schreb. phasc. p.* 8. *ſpic. p.* 71. *Sp. pl. p.* 1570. *Hedw. musc.* 1. *t.* 35. *Curtis lond. Weis crypt. p.* 268. *Web. ſpic. p.* 127. *Hoffm. fl.* 2. *p.* 19.

Bryum ericetorum. *Neck. meth. p.* 231.

Sphagnum acaulon trichodes. *Dill. musc.* 251. *t.* 32. *f.* 10.

Cum præc.

SPHAGNUM. Capſula ſeſſilis in orbiculo pedunculum terminante; ſæpe calyptræ parte inferiore abrupta cincta. Periſtomium nullum.

Masc. in ramis ſuperioribus, axillares.

paluſtre. 1. S. ramis deflexis. *Sp. pl. p.* 1569. *Fl. dan. t.* 474. *Scop. carn. tom.* 2. *p.* 305.

Sphagnum cauliferum ramis teretibus pendulis. *Hall. hiſt. n.* 1724.

In paluſtribus, aliisque udis locis montanis, ſubalpinis, & alpinis.

GYMNOSTOMUM. Capſula ovata. Periſtomium nullum.

Maſc. gemmacei in eadem, vel discoidei in diſtincta planta.

pyriforme. 1. G. foliis explanatis ovato - lanceolatis patentibus, capſula obovata. *Hoffm. fl.* 2. *p.* 27.

Bryum caule breviſſimo, foliis ovato - lanceolatis, capſulis erectis, ovatis,, operculo brevi. *Hall. hiſt. n.* 1832.

Bryum pyriforme. *Sp. pl. p.* 1580. *Weis crypt. p.* 183. *Neck. meth. p.* 197. *Scop. carn. tom.* 2. *p.* 318.

Bryum ſerpyllifolium pellucidum, capſulis pyriformibus. *Dill. musc.* 345. *t.* 44. *f.* 6.

Ad foſſas, in hortis, aliisque umbroſis, & humidis locis.

truncatulum. 2. G. foliis explanatis mucronatis, capſulis truncatis. *Sibthorp oxon. p.* 274. *Hedw. musc.* 1. *t.* 5. *Hoffm.* 2. *p.* 27.

Bryum truncatulum. *Sp. pl. p.* 1584. *Curtis lond. Fl. dan. t.* 537.

Bryum exiguum, creberrimis capſulis rufis. *Dill. musc.* 347. *t.* 45. *f.* 7.

Ad aggeres, in campis, agris, hortis, & alibi.

ovatum. 3. G. foliis ovatis concavis piliferis, capſulis ovatis. *Sibthorp oxon. p.* 274. *Hedw. musc.* 1. *t.* 6. *Hoffm. fl.* 2. *p.* 27.

Bryum capſulis erectis, ovatis; ſurculis ſimplicibus; foliis piliferis, oblongis. *Scop. carn. ed.* 1. *p.* 139. *n.* 6.

Bryum ovatum. *Dicks. faſc.* 2. *p.* 4

Bryum truncatulum. *Weber ſpic. p.*

Cum præc.

Heimii. 4. G. caule erecto, foliis ſpathulato-acun ce denticulatis, capſulis oblongis ere culo oblique roſtrato. *Hedw. musc Hoffm. fl.* 2. *p.* 28.

In agri vindobonenſis umbroſis ſyl uliginoſis.

æſtivum. 5. G. antheris erectis ſubrotundis axillar ſubulatis diſtantibus. *Sp. pl. p.* 1585.

Bryum paluſtre æſtivum, conferva f *musc.* 375. *t.* 47. *f.* 36.

In Styria in alpium judenburgenſiu

Hedwigia. 6. G. caule ramoſo, foliis ovato-lance cavis: apice diaphano denticulato; lium ciliato. *Hoffm. fl.* 2. *p.* 28.

Hypnum caulibus teretibus ramoſis, to-lanceolatis, pilo ariſtatis, opercu *Hall. hiſt. n.* 1792.

Hedwigia ciliata. *Hedw. musc.* 1. *t.*

Hedwigia Anodon. *Ehrh. beytr. tom.*

Bryum ſphagnoides. *Wulfen in Iacq tom.* 2. *p.* 222.

Fontinalis albicans. *Web. ſpic. p.* 38.

Sphagnum nodoſum, hirſutum & incan *musc.* 246. *t.* 32. *f.* 5.

Ad ſaxa Styriæ, Carinthiæ.

riparium. 7. G. caule ramoſo, foliis lanceolato- tentibus, capſulis ovatis: operculo bre

Bryum hypnoides alpinum, operculi *Dill. musc.* 371. *t.* 47. *f.* 30.

In agro vindobonenſi in ſaxis, pal nubii ripas.

aquaticum. 8. G. caule tenui declinato, extremitatibus foliisque linearibus acuminatis reclinatis, capſulis oblongis. *Hoffm. fl.* 2. *p.* 29.

Hedwigia aquatica. *Hedw. muſc.* 3. *t.* 11.

Hypnum aquaticum. *Iacqu. auſtr. t.* 290.

Hypnum aquaticum nigricans, foliis anguſtis falcatis. *Dill. muſc.* 334. *t.* 43. *f.* 70.

In fluviis, torrentibus Auſtriæ, Styriæ, Tyrolis ad pedes alpium; copioſiſſime in agro fluminenſi in *Rezzine* ſuper ſaxa torrentis.

TETRAPHIS. Capſula oblonga. Periſtomium ſimplex: dentibus quatuor pyramidalibus.

Maſc. discoidei, terminales, in diſtincta planta.

pellucida. 1. T. ſurculo ſimplici, foliis ovatis acutis, capſulis cylindricis. *Sibthorp. oxon. p.* 275. *Hedw. fund.* 2. *t.* 7. *f.* 32. *Hoffm. fl.* 2. *p.* 30.

Mnium caule ſimpliciſſimo, foliis ovato-lanceolatis, ſeta fungifera folioſa. *Hall. hiſt. n.* 1853. *t.* 46. *f.* 8.

Mnium pellucidum. *Sp. pl. p.* 1574. *Weis crypt. p.* 161. *Neck. meth. p.* 233.

Bryum diaphanum. *Web. ſpic. p.* 121.

Mnium ſerpylli foliis tenuibus pellucidis. *Dill. muſc.* 232. *t.* 31. *f.* 2.

In ſylvis montanis, & ſubalpinis.

SPLACHNUM. Capſula cylindracea apoph braculiformi, vel ſubgloboſæ inti inſidens. Periſtomium ſimplex: octo, geminatis.

Maſc. gemmaceo - discoidei; pe foli fertiles.

ampullaceum. 1. S. caule erecto ſubſimplici, apophyſi cea obconica. *Sibthorp oxon. p.* 275 *musc.* 2. *t.* 14. *Sp. pl. p.* 1572. *t.* 822. *Scop. carn. tom.* 2. *p.* 307. 2. *p.* 23.

Bryum ampullaceum, foliis Thymi collo ſtrictiore. *Dill. musc.* 343. *t.* 44

In pratis palluſtribus ſtercore vac ſpurcatis. Primo vere.

vasculoſum. 2. S. foliis ſpathulatis integris, obtuſiusc phyſi ſubgloboſa ſanguinea. *Hedw. m* 15. *Sp. pl. p.* 1572. *Hoffm. fl.* 2. *p.* 2

In Styria in humidis alpium jud ſium, præcipue in aſcenſu celſiſſimi *ſcharfe Eck* dicti.

mnioides. 3. S. foliis lanceolatis cuspidatis, erectiusc phyſi ſubgloboſa ſanguinea. *Hedw. mus*

Phascum pedunculatum. *Syſt. veget. p. dan. t.* 192.

Cum præc.

frœlichianum. 4. S. caule erecto ſubſimplici, foliis in linguæformibus, ſuperioribus ſpathulat ſis, apophyſi obliqua inverſe conica *musc.* 3. *t.* 40.

In Auſtriæ alpibus. *Frœlich.* Ego c

ſerratum. 5. S. caulibus erectis ramoſis, foliis ovati apophyſi ſubcylindrica. *Hedwig in litt.*

In alpibus judenburgenſibus cum

GRIMMIA. Capſula ovata. Periſtomium ſimplex: dentibus ſedecim, latiuſculis.

Maſc. gemmacei, vel capituliformes in eadem vel diſtincta planta.

apocarpa, 1. G. foliis ovatis acuminatis carinatis; perigonialibus apice marceſcentibus, capſulis ſubſeſſilibus terminalibus. *Sibthorp oxon. p.* 276. *Hedw. muſc.* 1. *t.* 39.

Hypnum caulibus ramoſis, foliis lanceolatis, hirſutis, operculis ariſtatis. *Hall. hiſt. n.* 1793.

Grimmia Polyodon. *Ehrh. beytr. tom.* 1. *p.* 168.

Fontinalis apocarpa. *Web. ſpic. p.* 38.

Bryum apocarpum. *Sp. pl. p.* 1579. *Hoffm. fl.* 2. *p.* 30. *Scop. carn. tom.* 2. *p.* 321. *Weis crypt. p.* 179.

Ad arborum truncos, ſaxa.

controverſa. 2. G. foliis lanceolatis acutis, ſiccitate contortis, capſulis ovatis. *Sibthorp. oxon. p.* 277.

Weiſſia controverſa. *Hedw. muſc.* 3. *t.* 5.

Bryum controverſum. *Hoffm. fl.* 2. *p.* 34.

Bryum capillaceum breve, pallide & læte virens, capſulis ovatis. *Dill. muſc. t.* 380. *t.* 48. *f.* 43.

In arenoſis, ad foſſas, fluvios, alibique.

recurvata. 3. G. foliis ſetaceis, pedunculis arcuatis, capſulis pyriformibus. *Hedw. muſc.* 1. *t.* 38.

Bryum recurvatum. *Dicks. faſc.* 2. *p.* 7. *Hoffm. fl.* 2. *p.* 31.

Bryum ſetaceum. *Wulfen in Iacqu. miſcell. tom.* 2. *p.* 96. *t.* 12. *f.* 1.

In ſylvis obducit ſaxa humida.

obliquata. 4. G. foliis ſetaceis, pedunculis erectis, oblongis, operculo obliquo.

In Styria in alpibus judenburgenſib

cirrhata. 5. G. capſulis ovato-oblongis, operculis tis, foliis congeſtis lanceolatis patulis, ribus, ſiccitate convolutis. *Hoffm. fl.* ſub Bryo.

Hypnum foliis capillaceis, congeſtis, centibus, capſulis ovato-cylindricis, b ciliatis. *Hall. hiſt. n.* 1786.

Mnyum cirrhatum. *Sp. pl. p.* 1576.

Hypnum cirrhatum. *Weis crypt. p.* 20

Bryum cirrhatum & ſtellatum tenuiorib *Dill. muſc.* 379. *t.* 48. *f.* 42.

In alpinis, ſubalpinis.

ENCALYPTA. Capſula cylindracea. Periſtomi plex: dentibus ſexdecim anguſtis.

Maſc. gemmacei axillares in eode viduo.

vulgaris. 1. E. calyptra laxa conica margine integr nulæ membranula pileiformi. *Hedw. mu* 18. ſub Leerſia. *Hoffm. fl.* 2. *p.* 27. *oxon. p.* 277.

Bryum caule breviſſimo, calyptra cylindr ge ariſtata. *Hall. hiſt. n.* 1829.

Bryum extinctorium, var, minor. *Sp. pl.*

Bryum calyptra extinctorii forma, mini *muſc.* 349. *t.* 45. *f.* 8.

Ad muros vetuſtos, ad terram.

ciliata. 2. E. calyptra laxa conica margine lacini ginulæ membranula cylindrica. *Hedw.* *t.* 19. ſub Leerſia. *Hoffm. fl.* 2. *p.* 27.

Bryum caule folioſo, capſula cylindric ta, longe ariſtata. *Hall. hiſt. n.* 1828.

Bryum extinctorium, var. major. *Sp. pl. p.* 1581.

Bryum contortum. *Wulfen in Iacqu. collect. tom.* 2. *p.* 226.,

Bryum calyptra extinctorii figura majus & ramoſum. *Dill. muſc.* 350. *t.* 45. *f.* 9.

In nemorenſibus, ſylvaticis ad terram, rupes.

lanceolata. 3. E. ſurculo ſubſimplici, foliis lanceolatis mucronatis concavis, capſulis ovatis. *Sibthorp oxon. p.* 278.

Leerſia lanceolata. *Hedw. muſc.* 2. *t.* 23.

Bryum lanceolatum. *Hoffm. fl.* 2. *p.* 31.

In collibus, montanis ſylvaticis Auſtriæ ad terram, ſaxa.

pulvinata. 4. E. foliis oblongis piliferis, capſulis ſubrotundis nutantibus: calyptris minutis. *Sibthorp. oxon. p.* 278.

Bryum foliis lanceolatis, pilo terminatis, capſulis pendulis, ovatis, ariſtatis. *Hall. hiſt. n.* 1822.

Bryum pulvinatum. *Sp. pl. p.* 1586. *Hoffm. fl.* 2. *p.* 31. *Web. ſpic. p.* 112. *Weis crypt. p.* 201. *Neck. meth. p.* 200.

Bryum orbiculare pulvinatum, hirſutie caneſcens, capſulis immerſis. *Dill. muſc.* 395. *t.* 50. *f.* 65.

Ad muros, tecta, ſaxa.

DICRANUM. Capſula ovato-oblonga. Periſtomium ſimplex: dentibus ſexdecim latiuſculis, bifidis.

Maſc. gemmacei, ſeu capituliformes in eadem ſeu diſtincta planta, axillares, ſeu terminales.

bryoides. 1. D. fronde ſimplici pinnata: pinnis oblongis acutis, apice peduncalifera, capſulis oblongis erectis. *Sibthorp. oxon. p.* 279.

Hypnum foliis planis, pinnarum paribu ſetis ex ſummitate. *Hall. hiſt. n.* 1776.

Fiſſidens bryoides. *Hedw. muſc.* 3. *t.*

Hypnum bryoides. *Sp. pl. p.* 1588. *I* 473. *f.* 1. *Weis cript. p.* 217. *Web. J* *Neck. meth. p.* 152. *Hoffm. fl.* 2. *p.* 5

Hypnum taxiforme exiguum, verſus ſu capſuliferum. *Dill. muſc.* 262. *t.* 34. *f.*

In umbroſis.

taxifolium. 2. D. fronde ſimplici pinnata: pinnis obl tis, baſi pedunculifera, capſulis oblongi *Sibthorp. oxon. p.* 279.

Hypnum foliis planis, pinnarum parib nis, ſetis radicalibus. *Hall. hiſt. n.* 177

Hypnum taxifolium. *Sp. pl. p.* 1587. *t.* 473. *f.* 2. *Web. ſpic. p.* 42. *Weis* 216. *Neck. meth. p.* 151. *Hoffm. fl.* 2.

Hypnum taxiforme minus, baſi capſuli *muſc.* 263. *t.* 34. *f.* 2.

Cum præc.

adiantoides. 3. D. fronde ramoſa pinnata: pinnis obl tis, medio pedunculifera, capſulis obl nutantibus. *Sibthorp oxon. p.* 280.

Hypnum foliis planis, pinnarum paribu ginta. *Hall. hiſt. n.* 1774.

Fiſſidens adiantoides. *Hedw. muſc.* 3. *t*

Hypnum adiantoides. *Sp. pl. p.* 1588. *p.* 47. *Neck. meth. p.* 153. *Hoffm. fl.*

Hypnum taxiforme paluſtre ramoſum erectum. *Dill. muſc.* 264. *t.* 34. *f.* 3.

In rupium fiſſuris; in paluſtribus,

ſciuroides. 4. D. ſurculis erectis incurvis, foliis adp ce piliferis, capſulis oblongis: opercul *Sibthorp oxon. p.* 280.

Hypnum caulibus teretibus ramoſis, foliis adpreſſis, pilo ariſtatis, operculis conicis. *Hall. hiſt. n.* 1749.

Hypnum ſciuroides. *Sp. pl. p.* 1596. *Weis crypt. p.* 258. *Neck. meth. p.* 175. *Hoffm. fl.* 2. *p.* 66.

Hypnum arboreum ſciuroides. *Dill. muſc.* 319. *t.* 41. *f.* 54.

Ad arborum truncos.

purpureum. 5. D. ſurculo ramoſo, foliis lanceolatis acutis carinatis, pedunculis axillaribus, capſulis cylindricis. *Sibthorp oxon. p.* 281.

Bryum foliis lanceolatis, ſubulatis, capſulis erectis, operculis conicis. *Hall. hiſt. n.* 1816.

Bryum purpureum. *Weis crypt. p.* 198. *Web. ſpic. p.* 100. *Schreb. ſpic. p.* 79. *Neck. meth. p.* 211. *Hoffm. fl.* 2. *p.* 36.

Mnium purpureum. *Sp. pl. p.* 1575.

Bryum tenue ſtellatum, ſetis purpureis. *Dill. muſc.* 386. *t.* 49. *f.* 51.

In ſylvis, hortis ad terram, ad vias.

ſcoparium. 6. D. capſulis oblongo-cylindricis, foliis lineari-lanceolatis ſecundis recurvis, ſurculo ramoſo declinato. *Sibthorp oxon. p.* 281.

Bryum ſcoparium. *Sp. pl. p.* 1582. *Curtis lond. Neck. meth. p.* 223. *Hoffm. fl.* 2. *p.* 39.

Hypnum ſcoparium. *Weis crypt. p.* 213. *Web. ſpic. p.* 71.

Bryum reclinatum, foliis falcatis, ſcoparum effigie. *Dill. muſc.* 357. *t.* 46. *f.* 16.

In ſylvaticis.

rugoſum. 7. D. capſulis ovatis cernuis, foliis lanceolatis apice ſerratis rugoſo-undulatis ſubſecudis. *Hoffm. fl.* 2. *p.* 39. ſub Bryo. *Schrader ſpic. p.* 59.

In turfoſis alpium tyrolenſium, ſtyriacarum.

Surculi tomento denſo ferrugineo obducti. Hoffm. l. c.

glaucum, 8. D. capſulis oblongis: operculo acuminato vo, foliis erectis imbricatis, ſurculis ramo tis. *Sibthorp oxon. p.* 281.

Hypnum caulibus erectis, teretibus, folii catis, fragilibus, capſulis inclinatis. *Ha n.* 1785.

Hypnum glaucum. *Weis crypt. p.* 208. *W p.* 75.

Bryum glaucum. *Sp. pl. p.* 1582. *Neck. r* 226. *Hoffm. fl.* 2. *p.* 38.

Bryum albidum & glaucum fragile majus erectis, ſetis brevibus. *Dill. muſc.* 362. *f.* 20.

In ſylvis montanis, ſubalpinis.

hetero-mallum. 9. D. capſulis ovatis: operculo acuminato r foliis ſetaceis recurvis ſecundis. *Sibthorp o* 282. *Hedw. muſc.* 1. *t.* 26.

Bryum foliis capillaribus heteromallis, ovatis, operculis ariſtatis. *Hall. hiſt. n.*

Bryum heteromallum. *Dill. muſc.* 375. *t.* 37. *Sp. pl. p.* 1582. *Hoffm. fl.* 2. *p.* 38. *meth. p.* 229.

Hypnum heteromallum. *Weis crypt. p.* 215 *ſpic. p.* 72.

In ſylvaticis ad terram, arborum rac

flexuo-ſum. 10. D. capſulis operculisque erectis, foliis ſ pedunculis flexuoſis. *Sp. pl. p.* 1583. ſul *Hoffm. fl.* 2. *p.* 38.

Bryum foliis capillaceis, capſulis cylindricis culis ariſtatis. *Hall. hiſt. n.* 1804.

Bryum pilosum molle ſetis intortis. *Dill* 273. *t.* 47. *f.* 33.

In Styriæ ſubalpinis ad terram.

puſil-lum. 11. D. capſulis ovatis, operculis roſtratis, pe lis ſiccis flexuoſis, foliis ſetaceis rigidis *Sibthorp oxon. p.* 282. *Hedw. muſc.* 2. *t*

Bryum parvulum. *Dicks. faſc.* 3. *p.* 7. *Hoffm. fl.* 2. *p.* 34.

In agri vindobonenſis uliginoſis, ad Danubii ripas.

varium. 12. D. capſulis ovato-oblongis cernuis: operculo conico, foliis ſubulatis ſparſis. *Hedw. muſc.* 2. *t.* 34.

Dicranum ſimplex. *Sibthorp oxon. p.* 282.

Bryum ſimplex. *Sp. pl. p.* 1587. *Hoffm. fl.* 2. *p.* 35.

Bryum rubrum. *Huds. angl. ed.* 1. 413.

Bryum trichodes, capſulis rubris cernuis. *Dill. muſc.* 390. *t.* 50. *f.* 59.

In argilloſis, ad fluviorum ripas, alibique.

ſtrumiferum. 13. D. foliis linearibus tortilibus, capſulis cernuis baſi interiore ſtrumoſis. *Hedw. muſc.* 2. *t.* 32. ſub Fiſſidente.

Bryum ſtrumiſerum. *Hoffm. fl.* 2. *p.* 36. *Dicks. faſc.* 3. *p.* 8.

In Styriæ, Tyrolis alpinis, ſubalpinis.

polycarpum. 14. D. capſulis ovatis erectis ſubobliquis, operculo incurvo, foliis linearibus ſiccitate tortilibus. *Hoffm. fl.* 2. *p.* 37. ſub Bryo.

Fiſſidens policarpus. *Hedw. muſc.* 2. *t.* 31.

In alpinis, ſubalpinis.

longifolium. 15. D. capſulis obliquis, foliis e lanceolata concava baſi longe canaliculatim capillaribus, arcuatim heteromallis. *Hedw. muſc.* 3. *t.* 9.

In alpibus judenburgenſibus.

flagellare. 16. D. capſulis erectiuſculis ſubcilindraceis: operculo longe roſtrato incurvo, foliis ſubulatis canaliculatis ſecundis, caule ramoſo interrupte flexuoſo. *Dicks. faſc.* 3. *p.* 6. *Hoffm. fl.* 2. *p.* 37. ſub Bryo.

Dicranum flagellare. *Hedw. muſc.* 3. *t.* 1.

Cum præc.

pelluci-dum. 17. D. capſulis erectiusculis, foliis recurvis caule hirſuto. *Sp. pl. p.* 1583. ſub Bryo.

Bryum palluſtre pellucidum, capſulis & brevibus recurvis. *Dill. musc.* 364. *t.* 46.

In alpium judenburgenſium riguis.

ambigu-um. 18. D. caule ſimplici recto, foliis lanceolatis ariſtatis quadrifariam imbricatis, capſula ſc loſo-longicolla. *Hedw. musc.* 3. *t.* 36.

In Styriæ ſubalpinis.

TRICHOSTOMUM. Capſula oblonga. Periſt ſimplex: dentibus ſexdecim, bipa laciniis ſetaceis, rectis, longis.

Masc. gemmacei axillares.

canes-cens. 1. T. capſulis oblongo-ſubulatis: operculo l mo aciformi, foliis lanceolatis carinatis e in apicem canum argute denticulatum pr *Hedw. musc.* 3. *t.* 3.

Bryum canescens. *Hoffm. fl.* 2. *p.* 41.

Bryum hypnoides hirſutie canescens, v *Dill. musc.* 368. *t.* 47. *f.* 27.

In montoſis ſylvaticis, collibus herbi

lanugi-noſum. 2. T. caule procumbente, ramis lateralibus fertilibus, foliis lineari-lanceolatis in api num longum eroſo-denticulatum productis *musc.* 3. *t.* 2.

Bryum lanuginoſum. *Hoffm. fl.* 2. *p.* 41.

Bryum hypnoides, polycephalon lanu montanum. *Dill. musc.* 372. *t.* 47. *f.* 32.

In alpibus judenburgenſibus ad terr pes humidas.

heteroſti-chum. 3. T. capſulis oblongis, foliis lanceolatis, in canum denticulatum productis. *Hedw.* *t.* 25.

Bryum heteroſtichum. *Hoffm. fl.* 2. *p.* 40.

In alpibus judenburgenſibus in rupeſtribus riguis.

glaucum. 4. T. caule erecto ſubramoſo: foliis acuminato-lanceolatis, glaucescentibus, capſulis rectis. *Hedw. musc.* 3. *t.* 37.

In Styria in aſcenſu alpium judenburgenſium, & prope *Eiſenärtz* in monte *Reicheſtein.*

fontinalioides. 5. T. caule longo, ramoſo, foliis lanceolatis patulis, capſulis lateralibus. *Hedw. musc.* 3. *t.* 14.

In Carinthiæ, Styriæ, Carnioliæ aquis ad ſaxa, arborum radices.

DIDYMODON. Capſula oblonga. Periſtomium ſimplex: dentibus triginta duobus, filiformibus, per paria approximatis.

Masc. axillares, ſeu terminales.

capillaceum. 1. D. capſulis erectis oblongis, pedunculis terminalibus, foliis ſetaceis patulis decurrentibus. *Hedw. musc.* 2. *t.* 26. ſub Swartzia.

Bryum capillaceum. *Hoffm. fl.* 2. *p.* 42. *Dicks. fasc.* 1. *p.* 4. *t.* 1. *f.* 6.

In montanis, ſubalpinis, alpinis Auſtriæ, Styriæ.

trifarium. 2. D. foliis imis lanceolatis, ſuperioribus ſubulatis carinatis, trifariis; capſula ovata erecta. *Hedw. musc.* 2. *t.* 28. ſub Swartzia.

In Styriæ montanis, ſubalpinis.

TORTULA. Capſula oblonga, ſeu cylindrac
ſtomium ſimplex: dentibus capillar
raliter convolutis.

Maſc. gemmacei, ſeu capituliform
dem vel diverſo individuo.

ſubulata. 1. T. capſulis cylindricis, operculo coni
foliis ovato-oblongis acutis, ſurculis ſi
breviſſimis. *Sibthorp oxon. p.* 284.

Bryum cauliculo breviſſimo, foliis l
capſulis & operculis longiſſimis. *Hall*
1826.

Bryum ſubulatum. *Sp. pl. p.* 1581. *Cu*
Web. ſpic. p. 99. *Weis crypt. p.* 18
meth. p. 194. *Hoffm. fl.* 2. *p.* 46.

Bryum capſulis longis ſubulatis. *Dill. n*
t. 45. *f.* 10.

In umbroſis ſylvaticis.

muralis. 2. T. capſulis oblongis erectis, foliis ova
piliferis, ſurculis ſubſimplicibus cæſpit
thorp. oxon. p. 284.

Bryum cauliculis breviſſimis, foliis o
ceolatis, patulis, pilo terminatis, ope
ſtato. *Hall. hiſt. n.* 1825.

Bryum murale. *Sp. pl. p.* 1581. *We*
100. *Weis crypt. p.* 189. *Neck. met*
Hoffm. fl. 2. *p.* 45.

Bryum tegulare humile piloſum & inca
muſc. 355. *t.* 45. *f.* 14.

Ad ſaxa, tecta, muros.

ruralis. 3. T. capſulis cylindricis erectiuſculis, f
vatis, carinatis piliferis, ſurculo ramoſo
oxon. p. 285.

Hypnum caulibus erectis, foliis lance
lo fluiante terminatis, capſulis erectis,
ſtratis. *Hall. hiſt. n.* 1789.

Bryum rurale. *Sp. pl. p.* 1581. *Hoffm. fl.* 2. *p.* 45. *Neck. meth. p.* 225.

Bryum rurale unguiculatum hirſutum, elatius & ramoſius. *Dill. muſc.* 352. *t.* 45. *f.* 12.

Ad tecta, muros, in pratis, campis, & alibi.

tortuoſa. 4. T. capſulis erectis, foliis ſetaceis inermibus areſactione retortis cincinatis. *Sp. pl. p.* 1583. ſub Bryo.

Hypnum foliis ſubulatis, cincinatis, capſulis cylindricis, longiſſime ciliatis. *Hall. hiſt. n.* 1787. *t.* 45. *f.* 2.

Bryum flaveſcens. *Scop. carn. tom.* 2. *p.* 321. *t.* 62.

Hypnum tortuoſum. *Web. ſpic. p.* 89.

Bryum cirrhatum, ſetis & capſulis longioribus. *Dill. muſc.* 377. *t.* 48. *f.* 40.

In montanis, alpinis, ſubalpinis.

unguiculata. 5. T. capſulis ſubcylindricis, operculo acuminato obliquo, pedunculis incraſſatis, foliis lineari-lanceolatis carinatis. *Sibthorp oxon. p.* 285.

Bryum unguiculatum & barbatum, ſurculis in ſummitate craſſioribus. *Dill. muſc.* 383. *t.* 48. *f.* 47.

In muris, locis arenoſis, montanis ſylvaticis, & alibi.

decipiens. 6. T. capſulis obliquis, operculo conico, pedunculis æqualibus, foliis lineari-lanceolatis carinatis. *Sibthorp oxon. p.* 285.

Barbula unguiculata. *Hedw. muſc.* 1. *t.* 23.

In ericetis, locis arenoſis, muris.

ſetacea. 7. T. capſulis erectis oblongo-ſubulatis: operculis filiformibus longitudine fere capſularum. *Sp. pl. p.* 1575. ſub Mnio.

Bryum ſetaceum. *Hoffm. fl.* 2. *p.* 43.

Bryum ſtellare nitidum pallidum, capſuli: ſimis. *Dill. muſc.* 381. *t.* 48. *f.* 44.

In inſulis danubialibus.

rigida. 8. T. capſulis oblongis erectis, ſurculis ſim breviſſimis, foliis oblongis rigidis margir lutis. *Sibthorp oxon. p.* 286.

Barbula rigida. *Hedw. muſc.* 1. *t.* 25.

Bryum rigidum. *Huds. angl. ed.* 2. 477.

Bryum acaulon, ericæ tenuifoliæ Gerard *Dill. muſc.* 388. *t.* 49. *f.* 55.

In arenoſis, collibus, montanis ſylv:

WEISIA. Capſula ovata. Periſtomium duple: terius dentibus ſexdecim latiuſculis; rius dentibus totidem latiuſculis erofi nullum. Calyptra conica, pilis erecti

Maſc. gemmacei axillares, & termin:

ſtriata. 1. W. capſulis ovato-cylindricis ſtriatis, fo ceolatis, margine revoluto, patulis: ſumı ce eroſis. *Sibthorp oxon. p.* 287.

Orthotrichum *ſtriatum*, caule ramoſiſſimo lanceolatis carinatis patulis, ſummis apice capſulis oblongis ſubpedunculatis. *Hoffm p.* 25. *Hedw. muſc.* 2. *t.* 36.

Bryum ſtriatum, var. major. *Sp. pl. p.* 1

Polytrichum Bryi ruralis facie, capſulis ſe majus. *Dill. muſc.* 430. *t.* 55. *f.* 8

Ad arborum truncos.

criſpa. 2. W. capſulis ovatis pedunculatis, peduno craſſatis, foliis lineari-lanceolatis ſiccita pis. *Sibthorp oxon. p.* 287.

Orthotrichum *criſpum*, foliis lanceolato-bus confertis ſiccitate criſpis, capſulis ped tis ex ampliata ſeta obovatis. *Hoffm. fl.* 2 *Hedw. muſc.* 2. *t.* 35.

Bryum ſtriatum, var. criſpa. *Sp. pl. p.* 1579.

Polytrichum capillaceum criſpum, calyptris acutis piloſiſſimis. *Dill. muſc.* 433. *t.* 55. *f.* 11.

Ad arborum truncos.

anomala. 3. W. capſulis ovatis ſtriatis, periſtomate interno nullo, foliis lanceolatis carinatis, margine revoluto, patulis. *Sibthorp oxon. p.* 287.

Orthotrichum *anomalum*, caule ramoſo, foliis lanceolatis carinatis patulis, capſulis pedunculatis oblongis ſtriatis. *Hoffm. fl.* 2. *p.* 25. *Hedw. muſc.* 2. *t.* 37.

Bryum ſtriatum, var. minor. *Sp. pl. p.* 1579.

Polytrichum Bryi ruralis facie, capſulis ſeſſilibus, minus. *Dill. muſc.* 431. *t.* 55. *f.* 9.

Ad ſaxa, arborum truncos.

POHLIA. Capſula ovato-oblonga, apophyſi obconicæ, anguſtiori inſidens. Periſtomium duplex: exterius dentibus ſexdecim latiuſculis; interius membrana ſedecimpartita.

Maſc. gemmacei in diſtincta planta.

elongata. 1. Pohlia. *Hedw. muſc.* 1. *t.* 36.

Bryum *elongatum*, foliis lineari-lanceolatis, capſulis oblique erectiuſculis, baſi elongatis. *Dicks. faſc.* 2. *p.* 8.

Mnium Pohlia. *Hoffm. fl.* 2. *p.* 48.

In Styria in umbroſis humidis in aſcenſu alpium judenburgenſium.

FUNARIA. Capſula obconica. Periſtomium exterius dentibus ſedecim obliquis a hærentibus; interius ciliis totidem naceis planis.

Maſc. diſcoidei in diverſa planta.

hygrometrica. 1. F. capſulis ſubovatis nutantibus: opercu foliis ovatis acuminatis concavis, ſurcul plici. *Sibthorp oxon. p.* 288.

Mnium foliis ovato-lanceolatis, conni capſulis ovatis, reflexis. *Hall. hiſt. n.* 1

Mnium hygrometricum. *Sp. pl. p.* 157 *crypt. p.* 113. *Hoffm. ſt.* 2. *p.* 50.

Bryum hygrometricum. *Web. ſpic. p.* 11 *meth. p.* 221.

Bryum bulbiforme aureum, calyptra qu lari, capſulis pyriformibus nutantibus. *D.* 407. *t.* 52. *f.* 75.

* Ad terram, muros.

Setæ, quæ humida tempeſtate arcuat ſunt, ſicca ingruente eriguntur, aut ſalt reflexæ obſervantur. Dill. l. c.

BRYUM. Capſula ovato-oblonga. Periſtomi plex: exterius dentibus ſedecim l lis acutis; interius membranaceum carinatum, laciniatum: laciniis lati capillaribusque alternis.

Maſc. capitati ſeu diſcoidei ſeu ge in eadem ſeu diverſa planta.

fontanum. 1. B. foliis lanceolato-ſubulatis ſubſecunc denticulatis, caule geniculis inflexo ſup moſo, capſulis ſubrotundis obliquis. *oxon. p.* 289.

Mnium caulibus teretibus, erectis, ex multifidis. *Hall. hiſt. n.* 1851.

Mnium fontanum. *Sp. pl. p.* 1574. *Hoffm. fl.* 2. *p.* 54. *Weis crypt. p.* 145.

Bryum fontanum. *Web. ſpic. p.* 120. *Neck. meth. p.* 209.

Bryum paluſtre, ſcapis teretibus ſtellatis, capſulis magnis ſubrotundis. *Dill. musc.* 340. *t.* 44. *f.* 2.

In udis, præcipue ſubalpinis, alpinisque.

ventricoſum. 2. B. capſulis cylindricis cernuis, foliis ſubulatis carinatis, caule ramoſo dichotomo. *Sibthorp oxon. p.* 289.

Mnium caulibus rubiginoſis bipartitis, foliis mollibus, lanceolatis, capſulis cylindricis acuminatis. *Hall. hiſt. n.* 1855.

Mnium paluſtre, *Sp. pl. p.* 1574. *Hoffm. fl.* 2. *p.* 47. *Weis crypt. p.* 165.

Bryum paluſtre. *Web. ſpic. p.* 122. *Neck. meth. p.* 210.

Mnium majus, ramis longioribus bifurcatis. *Dill. musc.* 233. *t.* 31. *f.* 3.

In udis, paluſtribus.

hornum. 3. B. capſulis oblongis cernuis, pedunculo curvato, foliis oblongo - lanceolatis leviter ſerratis, ſurculo ſubſimplici. *Sibthorp oxon. p.* 290.

Mnium foliis lanceolatis, imbricatis, capſulis pendulis, cylindricis obtuſis. *Hall. hiſt. n.* 1846.

Mnium hornum. *Sp. pl. p.* 1576. *Hoffm. fl.* 2. *p.* 48.

Bryum hornum. *Web. ſpic. p.* 115. *Neck. meth. p.* 215.

Bryum ſtellare hornum ſylvarum, capſulis magnis nutantibus. *Dill. musc.* 402. *t.* 51. *f.* 71.

In umbroſis, ſylvaticis.

crudum. 4. B. caule ſimplici, foliis ovato - lanceolatis, ſummis elongato - linearibus apice denticulatis, capſulis pendulis.

Mnium crudum. *Sp pl. p.* 1576. *Hedw. musc.* 1. *t.* 37. *Hoffm. fl.* 2. *p.* 48.

Bryum pendulum hornum molle, foli ceolatis & gramineis. *Dill. musc.* 40 *f.* 70.

In umbroſis ſubalpinis.

pſeudo-triquetrum. 5. B. foliis ovato-lanceolatis acutis, ca longo-ovatis pendulis. *Hedw. musc.* ſub Mnio.

Bryum annotinum paluſtre, capſulis v pendulis. *Dill. musc.* 404. *t.* 72.

In humidis ſubalpinis Styriæ, Ty

turbinatum. 6. B. foliis rubentibus ovato-lanceolati capſulis pyriformibus pendulis. *Hedw.* *t.* 8. ſub Mnio.

Bryum paluſtre complicatum rubens turbinatis pendulis. *Dill. musc.* 406. *t.*

In udis alpium judenburgenſium.

annotinum. 7. B. foliis ovatis acuminatis pellucidis, lis ſubradicalibus, capſulis nutantibus *p.* 1576. ſub Mnio.

Bryum annotinum lanceolatum pelluci ſulis oblongis pendulis. *Dill. musc.* 3 *f.* 68.

In humidis ſubalpinis.

nutans. 8. B. foliis lanceolatis ſubulatis, capſulis nutantibus. *Schreb. ſpic. p.* 81.

Bryum *ſericeum*, capſulis oblongis n foliis lanceolatis imbricatis carinatis. *oxon. p.* 292.

Webera nutans. *Hedw. musc.* 1. *t.* 4

Bryum trichodes læte virens, capitul oblongis. *Dill. musc.* 391. *t.* 50. *f.* 6

In Auſtria *in der Brein* in ascen *Weterkogel.*

pyriforme. 9. B. caule ſimpliciſſimo, foliis perichæt ribus longiſſimis recurvis, capſula *Sp. pl. p.* 1576. ſub Mnio.

Bryum foliis capillaceis, capitulis pyriformibus pendulis. *Hall. hiſt. n.* 1813. *t.* 46. *f.* 7.

Webera pyriformis. *Hedw. muſc.* 1. *t.* 3.

Bryum aureum. *Schreb. ſpic. p.* 81.

Bryum trichodes aureum, capſulis pyriformibus nutantibus, *Dill. muſc.* 391. *t.* 50. *f.* 60.

In agro vindobonenſi in umbroſis ad vetuſtos hortorum muros.

cæſpiticium. 10. B. foliis lanceolatis acuminato-ſetaceis, capſulis pendulis, pedunculis longiſſimis. *Sp. pl. p.* 1586. *Sibthorp oxon. p.* 290.

Mnium cæſpiticium. *Hoffm. fl.* 2. *p.* 50.

Bryum pendulum ovatum, cæſpiticium & piloſum, ſeta bicolori. *Dill. muſc.* 396. *t.* 50. *f.* 66.

In muris, tectis, in ſylvaticis ad terram.

capillare. 11. B. foliis ovatis ſetiferis carinatis, capſulis pendulis, pedunculis longis. *Sp. pl. p.* 1586. *Sibthorp oxon. p.* 290.

Mnium capillare. *Hoffm. fl.* 2. *p.* 50.

Bryum foliis latiuſculis congeſtis, capſulis longis nutantibus. *Dill. muſc.* 398. *t.* 50. *f.* 67.

In ſylvaticis.

alpinum. 12. B. capſulis pendulis oblongis, foliis ovatis acuminatis carinatis, ſurculis ramoſis, axillis pedunculiferis. *Syſt. veget. p.* 949.

Bryum hypnoides pendulum ſericeum, coma inſigni atro-rubente. *Dill. muſc.* 394. *t.* 50. *f.* 64.

In Auſtria *in der Brein* in monte *Weterkogel.*

argenteum. 13. B. foliis ovatis concavis imbricatis conniventibus ſetiferis, ſurculis teretibus, capſulis pendulis. *Sp. pl. p.* 1586. *Sibthorp oxon. p.* 293. *Web. ſpic. p.* 111. *Weis crypt. p.* 199. *Neck. meth. p.* 201.

Bryum cauliculis teretibus, capſulis ovatis acuminatis pendulis. *Hall. hiſt. n.* 1821.

Mnium argenteum. *Hoffm. fl.* 2. p. 51.

Bryum pendulum julaceum, argenteum
ceum. *Dill. muſc.* 392. *t.* 50. *f.* 62.

In muris, tectis, alibique ad terra

carneum. 14. B. foliis lanceolatis alternis pellucidis,
ovatis pendulis. *Sp. pl. p.* 1587.

Bryum foliis lanceolatis rarioribus, cap
tis pendulis. *Hall. hiſt. n.* 1834. *t.* 46.

Mnium carneum. *Hoffm. fl.* 2. p. 51.

Bryum delicatulum. *Hedw. muſc.* 1. *t.*

Bryum lanceolatum pellucidum, capſul
dis, pendulis, carneis. *Dill. muſc.* 40
f. 69.

In umbroſis, humidis.

ſtellare. 15. B. foliis ovatis ariſtatis ſerratis, pedun
leque ſimplicibus, capſulis nutantibus.
meg. 846. *Hoffm. fl.* 2. *p.* 51. ſub Mni

Bryum ſtellare roſeum pendulum minu
lis & ſetis longioribus. *Dill. muſc.* 41
f. 78.

In umbroſis humidis in aſcenſu al
denburgenſium.

cuſpidatum. 16. B. foliis ovatis acutis ſerratis, ſurculis
tibus, caule ſimplici, pedunculis aggre
ſolitariis, capſulis nutantibus. *Schreb.*
84. *Web. ſpic. p.* 117.

Mnium foliis ovato-lanceolatis, ſerra
ſertis, capſulis nutantibus obtuſis. *H*
n. 1841.

Bryum polymorphum. *Neck. meth. p.*

Mnium cuſpidatum. *Hoffm. fl.* 2. *p.* 5

Mnium ſerpyllifolium, var. cuſpidata.
p. 1577. *Weis crypt. p.* 153.

Bryum pendulum, foliis variis pellucid
ſulis ovatis. *Dill. muſc.* 413. *t.* 53. *f.*

In umbroſis ſylvaticis.

punctatum. 17. B. capſulis cernuis, pedunculis aggregatis vel ſolitariis, foliis obovatis integerrimis obtuſis punctatis. *Schreb. ſpic. p.* 89. *Web. ſpic. p.* 116.

Mnium foliis ovatis integerrimis, capſulis pendulis, ovatis, ariſtatis. *Hall. hiſt. n.* 1845.

Bryum ſerpyllifolium. *Neck. meth. p.* 216.

Mnium ſerpyllifolium. *Hoffm. fl.* 2. *p.* 52.

Mnium ſerpyllifolium, var. punctata. *Sp. pl. p.* 1577. *Weis crypt. p.* 153.

Bryum pendulum, ſerpylli folio rotundiore pellucido, capſulis ovatis. *Dill. muſc.* 416. *t.* 53. *f.* 81.

Cum praec.

roſeum. 18. B. capſulis nutantibus, pedunculis aggregatis, foliis roſaceo-congeſtis lanceolatis acutis. *Schreb. ſpic. p.* 84. *Web. ſpic. p.* 118. *Neck. meth. p.* 219.

Mnium foliis ovato-lanceolatis congeſtis, patulis, ſetis numeroſis, capſulis ovatis, acuminatis, pendulis. *Hall. hiſt. n.* 1839.

Mnium roſeum. *Weis crypt. p.* 157. *Hoffm. fl.* 2. *p.* 53.

Mnium ſerpyllifolium, var. prolifera. *Sp. pl. p.* 1577.

Bryum ſtellare roſeum majus, capſulis ovatis pendulis. *Dill. muſc.* 411. *t.* 52. *f.* 77.

Cum praec.

undulatum. 19. B. foliis oblongis undulatis ſerratis, pedunculis aggregatis, capſulis nutantibus. *Schreb. ſpic. p.* 84. ſub B. ligulato.

Mnium foliis cordatis, ellipticis, ſetis aggregatis, capſulis ovatis, obtuſis, pendulis. *Hall. hiſt. n.* 1844.

Bryum dendroides. *Web. ſpic. p.* 117. *Neck. meth. p.* 218.

Mnium undulatum. *Weis crypt. p.* 158. *Hoffm. fl.* 2. *p.* 53.

Mnium ſerpyllifolium, var. undula
p. 1577.

Bryum dendroides polycephalon, Ph
lio undulato pellucido, capſulis ovat
Dill. muſc. 410. *t.* 52. *f.* 76.

Cum præc.

pomiforme. 20. *B.* foliis lanceolato-ſetaceis patulis
capſulis ſphæricis erectis. *Sp. pl. p.*

Bryum foliis ſubulatis, ſetis vertica
ſulis ſphæricis. *Hall. hiſt. n.* 1803.

Mnium pomiforme. *Hoffm. fl.* 2. *p.*

Bryum capillaceum, capſulis ſphæ
muſc. 339. *t.* 44. *f.* 1.

In montanis, ſubalpinis, alpinis

MEESIA. Capſula oblonga. Periſtomium du
terius dentibus ſedecim, brevib
ſis; interius ciliis totidem, acut
tis, vel opere reticulato conne

Maſc. femineis approximati, ve
in diverſa planta.

longiſeta. 1. M. foliis trifariis patulis ſubulato-
carinatis, capſulis obliquis. *Hedw.*
21. 22.

Mnium triquetrum. *Sp. pl. p.* 1578.
2. *p.* 47.

In palluſtribus Pannoniæ ad lac
nis. In ſpongioſis Carniæ in monte

trichodes. 2. M. foliis lineari-lingulatis obtuſis,
liquis. *Sp. pl. p.* 1585. ſub *Bryo.*

Bryum foliis capillaceis erectis, ſe
mis, capſulis ſurſum lateſcentibus
breviſſimo. *Hall. hiſt. n.* 1812.

Meeſia uliginoſa. *Hedw. muſc.* 1. *t.*

Hypnum trichodes. Web. ſpic. p. 87.

Mnium trichodes. *Hoffm. fl.* 2. *p.* 47.

Bryum trichodes aureum, capſulis incurvis obtuſis, in ſetis longis. *Dill. muſc.* 389. *t.* 49. *f.* 58.

In alpinis.

BARTHRAMIA. Capſula ſphærica, ſtriata. Periſtomium duplex: exterius dentibus ſedecim, cuneiformibus; interius membranaceum, plicato-carinatum, apice lacinulatum.

Maſc. Femineis approximati.

halleriana. 1. *B.* foliis lanceolato-ſetaceis ſubſecundis denticulatis, capſulis ovatis pedunculatis lateralibus. *Hedw. muſc.* 2. *t.* 40.

Bryum foliis mollibus ſubulatis, ſetis breviſſimis alaribus, capſulis ovatis. *Hall. hiſt. n.* 1802. *t.* 45. *f.* 8.

Bryum laterale. *Huds. angl.* 2. 483.

Bryum recurvum. *Wulfen in Iacqu. collect. tom.* 2. *p.* 224.

Mnium laterale. *Hoffm. fl.* 2. *p.* 64.

In Carinthia in alpibus ſpitalenſibus retro *Hintereggen. Wulfen.* Ego in Styria in umbroſis in aſcenſu alpium judenburgenſium.

FONTINALIS. Capſula inſidens ſetæ brevi, perichætio imbricato obvolutæ. Periſtomium duplex: exterius dentibus ſedecim, latiuſculis; interius reticulatum.

Maſc. gemmacei axillares.

antipyretica. 1. F. foliis complicato-carinatis trifariis acutis, capſulis lateralibus. *Sp. pl. p.* 1571. *Weis crypt. p.* 261. *Hoffm. fl.* 2. *p.* 79.

Hypnum foliis complicatis, ovato-lanceolatis, capſulis in alis petiolatis, ovatis. *Hall. hiſt. n.* 1794.

Fontinalis triangularis major complicata, e foliorum alis capſulifera. *Dill. muſc.* 254. *t.* 33. *f.* 1.

In aquis.

PTERIGYNANDRUM. Periſtomium ſimplex ſedecim dentatum.

Flos maſculus femineusque alaris.

gracile. 1. P. prolongationibus extantibus faſciculatim ramificatis recurvis, foliis lanceolatis imbricatis, capſulis oblongis erectis. *Hedw. muſc.* 4. *t.* 6.

Hypnum *gracile*, ſurculis repentibus, ramis faſciculatis teretibus erectiuſculis, antheris erectis ovatis. *Huds. angl.* 430. *Syſt. veget. p.* 952.

Hypnum gracile ornithopodioides. *Dill. muſc.* 320. *t.* 41. *f.* 55.

In agri fluminenſis ſylvaticis.

filiforme. 2. P. caule debili ſubramoſo decumbente, foliis lanceolatis concavis, imbricatis, capſulis erectis. *Hedw. muſc.* 4. *t.* 7.

Hypnum *filiforme*, caule ſurculisque filiformibus, foliis adpreſſis concavis, capſulis erectis. *Hoffm. fl.* 2. *p.* 71. *Timm. megap.* 817.

In Styriæ ſubalpinis.

HYPNUM. Capſula oblonga. Periſtomium duplex: exterius dentibus ſedecim latiuſculis; interius membranaceum, æquale, laciniatum: laciniis latiuſculis, cum interjectis capillaribus.

Maſc. gemmacei in diverſis plantis.

* *Fronde ſimplici, ſubramoſa, pinnata.*

denticulatum. 1. H. fronde ſimplici pinnata duplicata, baſi pedunculifera. *Sp. pl. p.* 1588. *Hoffm. fl.* 2. *p.* 56. *Web. ſpic. p.* 43. *Neck. meth. p.* 154. *Schreb. ſpic. p.* 86.

Hypnum foliis laxis, pinnatis, ovato-lanceolatis, ariſtatis, retroverſis. *Hall. hiſt. n.* 1772.

Hypnum denticulatum pennatum, pinnulis duplicatis recurvis.

In umbroſis montanis, ſubalpinis.

ſylvaticum. 2. H. fronde ſubramoſa pinnata, ſupra baſin pedunculifera. *Huds. angl.* 2. 419. *Syſt. veget. p.* 950. *Hoffm. fl.* 2. *p.* 56.

Hypnum denticulatum, pinnatum, pinnis ſimplicibus rectioribus. *Dill. muſc.* 267. *t.* 34. *f.* 6. *Hall. hiſt. n.* 1773.

Cum præc.

lucens. 3. H. fronde ſubramoſa pinnata imbricata, foliis reticulatis planis, ſupra baſin pedunculifera. *Sp. pl. p.* 1589. *Hoffm. fl.* 2. *p.* 57. *Scop. carn. tom.* 2. *p.* 331. *Web. ſpic. p.* 52.

Hypnum pennatum, aquaticum, lucens, longis latisque foliis. *Dill. muſc.* 270. *t.* 34. *f.* 10.

In humidis ſylvaticis Carnioliæ, Croatiæ ad terram.

Pinnæ duplici, interdum triplici verſu diſpoſitæ, luci obverſæ tenuiter & eleganter reticulatæ & punctatæ. Dill. l. c.

** *Caulibus vel ſurculis ramoſ natis.*

undulatum. 4. H. frondibus ramoſis vage pinnatis, latis complicatis nitidis. *Sp. pl. p.* 15 *ſl.* 2. *p.* 57. *Web. ſpic. p.* 55. *Neck* 156. *Weis crypt. p.* 220. *Scop. car p.* 331.

Hypnum caule plano, foliis diſtich rugoſis, nitidis, operculo conico. *n.* 1770.

Hypnum pennatum undulatum, lycop ſparſum. *Dill. musc.* 271. *t.* 36. *f.* 1

In Carnioliæ alpium convallibus pinis ad terram, ſaxa.

purum. 5. H. ſurculis pinnato-ſparſis teretibus, tis concavis mucronatis conniventibus 1594. *Curtis lond. Hoffm. ſl.* 2. *p. ſpic. p.* 78. *Weis crypt. p.* 249. *tom.* 2. *p.* 335.

Hypnum ramis teretibus, foliis ovat latis, ſetis prælongis, capſulis inclina tis. *Hall. hiſt. n.* 1743.

Hypnum cupreſſiforme vulgare, fol *Dill. musc.* 309. *t.* 40. *f.* 45.

In pratis, ſylvis.

compreſſum. 6. H. ſurculis plano-pinnatis apice atte liis ovato-lanceolatis. *Schreb. ſpic. p. ſl.* 2. *p.* 58.

Hypnum ramis teretibus, foliis ova latis, ſetis prælongis, capſulis craſſi lis conicis. *Hall. hiſt. n.* 1742.

Hypnum cupreſſiforme tenuius & c *Dill. musc.* 312. *t.* 40. *f.* 47.

In pratis Auſtriæ palluſtribus p *Himberg.*

cuſpidatum. 7. H. ſurculis pinnato-ramoſis: apice volutis acuminatis. *Sp. pl. p.* 1595. 2. *p.* 58.

Hypnum foliis ovato-lanceolatis, ramis teretibus, ſupremis conicis, acuminatis. *Hall. hiſt. n.* 1741.

Hypnum paluſtre, extremitatibus cuſpidatis & pungentibus. *Dill. muſc.* 300. *t.* 39. *f.* 34.

In udis, paluſtribus.

trique-trum. 8. H. ſurculis pinnato-ſparſis recurvis, foliis triangularibus patulis. *Sp. pl. p.* 1589. *Hoffm. fl.* 2. *p.* 59. *Web. ſpic. p.* 77. *Scop. carn. tom.* 2. *p.* 337. *Neck. meth. p.* 185. *Weis crypt. p.* 222.

Hypnum foliis hiſpidis, ovato-lanceolatis, ariſtatis. *Hall. hiſt. n.* 1729.

Hypnum vulgare triangulum maximum & pallidum. *Dill. muſc.* 293. *t.* 38. *f.* 28.

In ſylvaticis.

nitens. 9. H. ſurculis erectis, ramulis breviſſimis, foliis lanceolatis appreſſis ſtriatis, perichætiis longiſſimis, capſulis obliquis. *Schreb. ſpic. p.* 92. *Hoffm. fl.* 2. *p.* 59.

Hypnum paluſtre erectum trichodes, ramulis crebris, luteo- & rufo-virentibus glabris. *Dill. muſc.* 303. *t.* 39. *f.* 37.

In paluſtribus.

abieti-num. 10. H. ſurculis ſimpliciter pinnatis, pinnis teretiuſculis, remotis inæqualibus. *Sibthorp oxon. p.* 296. *Sp. pl. p.* 1591. *Hoffm. fl.* 2. *p.* 59. *Web. ſpic. p.* 63. *Neck. meth. p.* 163. *Weis crypt. p.* 236. *Scop. carn. tom.* 2. *p.* 338.

Hypnum ramis duris, teretibus, foliis appreſſis, ovato-lanceolatis, pilo ariſtatis. *Hall. hiſt. n.* 1762.

Hypnum luteſcens, alis ſubulatis tenacibus. *Dill. muſc.* 280. *t.* 35. *f.* 17.

In ſylvaticis, & alibi.

prolife-rum. 11. H. ſurculis proliferis triplicato-pinnatis, capſulis incurvis. *Hoffm. fl.* 2. *p.* 60. *Sp. pl. p.* 1590. *Curtis lond.* *Sibthorp oxon. p.* 294.

Hypnum filicinum, Tamariſci foliis non ſplendentibus. *Dill. muſc.* 276. *t*

In ſylvis.

umbratum. 12. H. ſurculis proliferis bipinnato-ſparſis, capſulis obliquis. *Hoffm. fl.* 2. *p.*

In Auſtriæ ſylvis umbroſis.

Folia cordato-ovata, acuminata, lentem ſerrata; ſetæ vix unciales. Ho

ſplendens. 13. H. ſurculis ſubtriplicato-pinnatis prc liis lanceolatis confertis ſplendentibus lis aggregatis. *Sibthorp oxon. p.* 295.

Hypnum *parietinum*, ſurculis proli bipinnatis, capſulis obliquis. *Hoffm.* *Sp. pl. p.* 1590.

Hypnum filicinum, tamariſci foliis ſplendentibus. *Dill. muſc.* 274. *t.* 35.

In ſylvaticis.

*** *Foliis ſecundis vel reflexi*

commutatum. 14. H. caule procumbente, ſurculis ram tis ſetiferis, foliis ſecundis. *Hedw. mu* *Hoffm. fl.* 2. *p.* 60.

Muſcus filicinus paluſtris. *Vaill. pa* 29. *f.* 9.

In udis alpium judenburgenſium.

affine. 15. H. caule procumbente ramoſo pinna foliis ſubſecundis nervoſis: nervo ducto. *Hoffm. fl.* 2. *p.* 61. *Dill. mus* 19. *A.*

In ſylvis Auſtriæ, Styriæ ſubalpi ram, marceſcentes arborum truncos.

filicinum. 16. H. caule erecto ſimplici pinnato ſetif uncinatis ſecundis ſtriatis. *Hoffm. fl* *Dill. muſc. t.* 36. *f.* 19. *B. F.*

In alpium ſtyriacarum udis.

criſtà ca- ſtrenſis. 17. H. ſurculis pinnatis: ramulis approximatis, apicibus recurvis. *Sp. pl. p.* 1591. *Hoffm. fl.* 2. *p.* 63.

Hypnum filicinum, criſtam caſtrenſem repræſentans. *Dill. muſc.* 284. *t.* 36. *f.* 20.

In ſylvis montanis, ſubalpinis.

extricatum. 18. H. caule decumbente ſetifero, foliis uncinatis, ſurculis erectis pinnatis ſenſim attenuatis apice convoluto - ſeuncdis. *Hoffm. fl. p.* 62.

In alpium judenburgenſium udis.

uncinatum. 19. H. caule proſtrato, foliis ſubulato - carinatis ſtriatis, heteromalle uncinatis, perichætiis gracilibus longis. *Hedw. muſc.* 4. *t.* 15. *Hoffm. fl.* 2. *p.* 62.

Cum præc.

molluſcum. 20. H. procumbens, ramis ſimplicibus curvatis, foliis confertis varie diſtortis incurvis ductulorum faſciculo deſtitutis. *Hedw. muſc.* 4. *t.* 22.

Hypnum filicinum ſericeum, molle & pallidum mucronibus aduncis. *Dill. muſc.* 286. *t.* 36. *f.* 22.

Cum præc.

cupreſſiforme. 21. H. caule & ſurculis vage pinnatis, foliis ovato-acuminatis incurvatis ſecundis enerviis. *Sp. pl. p.* 1592. *Hoffm. fl.* 2. *p.* 63. *Hedw. muſc.* 4. *t.* 23. *Neck. meth. p.* 166. *Weis crypt. p.* 238. *Scop. carn. tom.* 2. *p.* 331.

Hypnum caule compreſſo, foliis imbricatis, falcatis, pilo ariſtatis. *Hall. hiſt. n.* 1737.

Hypnum crispum cupreſſiforme, foliis aduncis. *Dill. muſc.* 287. *t.* 37. *f.* 23.

In ſylvis.

rugoſum. 22. H. caule & ſurculis vage pinnatis erectiuſculis incraſſatis recurvis, foliis ſecundis recurvatis baſi rugoſis. *Syſt. veget. p.* 950. *Hoffm. fl.* 2. *p.* 63.

Hypnum luteſcens crispum, Lycopodii facie. *Dill. muſc.* 289. *t.* 37. *f.* 24.

In pratis, ſylvis.

aduncum. 23. H. caule erectiuſculo ramoſo-pinnato, patulis recurvis, foliis ovato-acuminat vatis, foliis perichætii apiculatis. *Hoff. p.* 64. *Sp. pl. p.* 1592. *Hedw. muſc.* 4.

Hypnum paluſtre erectum, ſummitatib cis. *Dill. muſc.* 292. *t.* 37. *f.* 26.

In udis, paluſtribus.

Halleri. 24. H. caule proſtrato, ſurculis erectiuſculi cibus, foliis lanceolatis faſciculo deſtit dique e baſi imbricata recurvis. *Hedw. t.* 21. *Syſt. veget. p.* 952. *Hoffm. fl.* 2

Hypnum caulibus proſtratis, foliis la reflexis, operculo conico. *Hall. hiſt. n.*

In alpibus judenburgenſibus.

ſquarroſum. 25. H. ſurculis vagis erectiuſculis, foliis c tis ovato-lanceolatis apice ſubulatis unc flexis. *Sp. pl. p.* 1593. *Hoffm. fl.* 2. *p.*

Hypnum foliis cincinnatis, patulis, *Hall. hiſt. n.* 1733.

Hypnum repens, triangularibus reflexi majus. *Dill. muſc.* 303. *t.* 39. *f.* 38.

Hyypum repens, triangularibus reflexi majus. *Dill. muſc.* 304. *t.* 39. *f.* 39.

In ſylvarum humidis.

loreum. 26. H. ſurculis vagis erectiuſculis recurvis, fc ſecundis. *Sp. pl. p.* 1593. *Hoffm. fl.* 2.

Hypnum loreum montanum, capſulis ſ dis. *Dill. muſc.* 305. *t.* 49. *f.* 40.

In ſylvis ſubalpinis.

**** *Foliis rectis, ſurculis faſcic tis, vel dendroidibus.*

myoſuroides. 27. H. ſurculis erectis, ramis faſciculatis a incurvis, foliis ovatis, capſulis erectis. *muſc.* 4. *t.* 8. *Schreb. ſpic. p.* 97. *Hoffm p.* 69.

Hypnum myoſuroides craſſius, capſulis erectis. *Dill. muſc.* 316. *t.* 41. *f.* 50.

In ſylvis.

alopecu-rum. 28. H. caule erecto, ramis faſciculatis terminalibus ſubdiviſis, capſulis ſubnutantibus. *Sp. pl. p.* 1594. *Hoffm. fl.* 2. *p.* 70. *Web. ſpic. p.* 74. *Weis crypt. p.* 246.

Hypnum caule nudo, confertim ramoſo, ramis inclinatis, capſulis ariſtatis. *Hall. hiſt. n.* 1748.

Hypnum dendroides obſcurius, ſetis & capſulis brevioribus nutantibus. *Dill. muſc.* 315. *t.* 41. *f.* 49.

In ſylvaticis.

ſerpens. 29. H. caule repente, ſurculis filiformibus ſimplicibus, ſubramoſis, foliis lanceolatis acuminatis remoto-patulis, capſulis oblongis cernuis. *Sp. pl. p.* 1596. *Hoffm. fl.* 2. *p.* 71. *Hedw. muſc.* 4. *t.* 18.

Hypnum trichodes ſerpens, ſetis & capſulis longis erectis. *Dill. muſc.* 329. *t.* 42. *f.* 64.

In ſylvaticis ad terram, ſaxa, arborum truncos.

dubium. 30. H. caule ſubdecumbente, ramis adproximatis filiformibus erectiuſculis vagis ſubpinnatis, foliis lanceolatis acutis erecto-patulis, capſulis oblongis incurvis. *Neck. meth. p.* 161. *Hoffm. fl.* 2. *p.* 71. *Dicks. faſc.* 3. *p.* 10.

Hypnum repens filicinum, trichodes palluſtre. *Dill. muſc.* 286. *t.* 36. *f.* 21.

In inſulis danubialibus ad terram.

prælon-gum. 31. H. caule decumbente, ſurculis ſubpinnatis: ramulis remotis, foliis ovato-lanceolatis patentibus, operculis roſtratis. *Sp. pl. p.* 1591. *Hoffm. fl.* 2. *p.* 72. *Hedw. muſc.* 4. *t.* 29.

Hypnum repens filicinum, triangularibus parvis foliis, prælongum. *Dill. muſc.* 378. *t.* 35. *f.* 15.

In umbroſis ſylvaticis.

rotundifolium. 32. H. ſurculis repentibus ramoſis, fc
patentibus diſtichis, capſulis ov
Dicks. faſc. 2. *p.* 10. *Scop. car*
339. *t.* 62.

In Carniolia ad muros. *Scop.*

intricatum. 33. H. ſurculis repentibus, ramis bre
lanceolatis acuminatis patulis, capſu
ſis obliquis cernuis, operculo con
ſpic. p. 99. *Hedw. muſc.* 4. *t.* 28.
p. 72.

Muscus terreſtris omnium minimu
majusculis oblongis erectis. *Vaill.*
t. 28. *f.* 6. 7.

Ad arborum truncos.

velutinum. 34. H. caule repente, ſurculis confertis
liis ovato-lanceolatis denticulatis,
bus interioribus crinitis, capſulis ol
exaſperatis. *Sp. pl. p.* 1595. *Hoffm.*
Hedw. muſc. 4. *t.* 27.

Hypnum velutinum, capſulis ov
Dill. muſc. 326. *t.* 42. *f.* 61.

Ad terram, arbores.

plumoſum. 35. H. ſurculis pinnatis repentibus, ran
foliis imbricatis ſubulatis, capſulis
pl. p. 1592. *Hedw. muſc.* 4. *t.* 15.

Hypnum repens filicinum plumoſum,
280. *t.* 35. *f.* 16.

Ad ſaxa, arborum truncos.

luteſcens. 36. H. caule ſurculisque vagis depreſſis
foliis ovato-lanceolatis acuminatis
ſulis cernuis, operculis oblique c
exaſperatis. *Schreb. ſpic. p.* 98. *He*
t. 16. *Hoffm. fl.* 2. *p.* 75.

Hypnum ſericeum, ſurculis longior
tioribus, capſulis incurvis. *Dill. m*
42. *f.* 60.

In pratis, ſylvis.

pilife-rum. 37. H. caule decumbente, ramis vagis rectiuſculis, foliis ovatis concavis; extremitate in cuſpidem piliformem protenſa, pedunculo muriculato, operculo roſtrato. *Hedw. muſc.* 4. *t.* 14. *Schreb. ſpic. p.* 91. *Hoffm. fl.* 2. *p.* 75.

In umbroſis ſylvaticis.

ſtriatum. 38. H. caule decumbente, ramis vagis recurvis; foliis patentibus plicato-ſtriatis, operculis mucronatis, ſetis glabris. *Schreb. ſpic. p.* 91. *Hedw. muſc.* 4. *t.* 13. *Hoffm. fl.* 2. *p.* 75.

Hypnum vulgare dentatum, operculis cuſpidatis. *Dill. muſc.* 297. *t.* 38. *f.* 30.

In ſylvis montanis, ſubalpinis.

murale. 39. H. caule declinato ſubramoſo, ramis compreſſiuſculis, foliis ovatis acutis teneris, operculo roſtrato. *Hedw. muſc.* 4. *t.* 30. *Hoffm. fl.* 2. *p.* 76.

Hypnum myoſuroides brevius & craſſius, capſulis cernuis. *Dill. muſc.* 318. *t.* 41. *f.* 52.

In ſylvaticis.

rutabulum. 40. H. caule procumbente vario, ſurculis ſubramoſis erectis curvatis, foliis lanceolato-acuminatis patulis, capſulis cernuis, operculo conico, ſetis muricato-exaſperatis. *Sp. pl. p.* 1590. *Hedw. muſc.* 4. *t.* 12. *Hoffm. fl.* 2. *p.* 77.

Hypnum dentatum vulgatiſſimum, operculis obtuſis. *Dill. muſc.* 295. *t.* 38. *f.* 29.

In ſylvis umbroſis, ad fluvios.

riparium. 41. H. caule proſtrato varie diviſo, ſurculis radicantibus, foliis ovato-acutis diſtichis, capſulis cernuis, operculo acuto, ſetis glabris. *Hedw. muſc.* 4. *t.* 3. *Sp. pl. p.* 1595. *Hoffm. fl.* 2. *p.* 77.

Hypnum aquaticum, flagellis & teretibus & pennatis. *Dill. muſc.* 308. *t.* 4. *f.* 44.

Ad foſſas, fluvios.

riparioi- 42. H. caule ſurculisque elongatis, folii
des. tis concavis, capſulis ovatis, cern
acuminatis. *Hedw. muſc.* 4. *t.* 4.

Hypnum ruſciſolium. *Neck. meth.*

Hypnum foliis ruſciformibus, capſi
dis. *Dill. muſc.* 298. *t.* 38. *f.* 31.

In ſylvis umbroſis, ad fluvio
rivulos.

LESKEA. Capſula oblonga. Periſtomium
terius dentibus ſedecim, acut
membranaceum, partitum in laci

Maſc. gemmacei in diverſis indi

trichoma- 1. L. fronde ramoſa, foliis diſtichis i
noides. naciformibus obtuſis. *Schreb. ſpic.*
fl. 2. *p.* 56. ſub Hypno.

Hypnum pennatum trichomanoides
moſum. *Dill. muſc.* 264. *t.* 34. *f.*

In ſylvis ſubalpinis Styriæ,
arborum radices.

compla- 2. L. fronde ramoſa pinnata, foliis d
nata. catis acinaciformibus acuminatis. *S*
87. *Hoffm. fl.* 2. *p.* 57. *Sp. pl. p.*

Hypnum pennatum compreſſum & ſ
ſulis ovatis. *Dill. muſc.* 268. *t.* 34

In ſylvis ad arborum truncos.

paluſtris. 3. L. caule decumbente repente, ſur
erectiusculis ſetis brevioribus, foliis
concavis ſecundis. *Hedw. muſc.* 4.

Hypnum paluſtre. *Sp. pl. p.* 1593.
p. 64.

Hypnum heterophyllum aquaticum
repens. *Dill. muſc.* 293. *t.* 37. *f.*

In alpium judenburgenſium ud

curtipendula, 4. L. ſurculis vagis ramoſo-pinnatis, foliis ovatis acuminatis patulis, perichætiis longiſſimis. *Sp. pl.* p. 1594. *Hoffm. fl.* 2. p. 66. ſub Hypno.

Hypnum dentatum curtipendulum, viticulis rigidis. *Dill. muſc.* 333. t. 43. f. 69.

In ſylvis ad arborum truncos.

attenuata. 5. L. ſurculis ramoſiſſimis: ramis incurvatis apice gracileſcentibus incraſſatisque, foliis ovato-acuminatis ſubſecundis. *Hedw. muſc.* I. t. 12.

Hypnum attenuatum. *Schreb. ſpic.* p. 100. *Hoffm. fl.* 2. p. 67.

Hypnum repens trichodes arboreum majus, cauliculis ramoſis. *Dill. muſc.* 331. t. 42. f. 66.

Ad arborum radices.

polycarpa. 6. L. caule ſurculisque decumbentibus filiformibus ramoſis, foliis adpreſſis nervoſis, capſulis erectis. *Hoffm. fl.* 2. p. 70. ſub Hypno.

Ad terram, arborum truncos.

ſubtilis. 7. L. caule decumbente reptante, ramulis erectiuſculis confertis, foliis exiguis patulis, capſulis erectis. *Hedw. muſc.* 4. t. 9.

Hypnum ſubtile. *Hoffm. fl.* 2. p. 70.

Ad arborum truncos.

polyantha. 8. L. caule repente ramoſiſſimo multifloro, ramulis capſulisque erectis. *Hedw. muſc.* 4. t. 2.

Hypnum *filifolium*, ſurculis vagis ramoſiſſimis: ramis filiformibus, antheris obliquis. *Huds. angl.* 497. *Syſt. veget.* p. 952.

Hypnum polyanthos. *Schreb. ſpic.* p. 97. *Hoffm. fl.* 2. p. 74.

Hypnum ſericeum ramoſius & tenuius, capſulis acuminatis. *Dill. muſc.* 327. t. 42. f. 62.

Ad arborum truncos.

dendroides. 9. L. caule erecto ſimplici, ramis faſciculati: nalibus ſimpliciuſculis, capſulis erectis. p. 1593. *Hoffm. fl.* 2. p. 69. *Web. ſpic* *Neck. meth.* p. 187. *Scop. carn. tom.* 2. *Weis crypt.* p. 245. ſub Hypno.

Hypnum dendroides ſericeum, ſetis & longioribus erectis. *Dill. muſc.* 313. *t.* 4

In ſylvis.

ſericea. 10. L. caule repente, ſurculis confertis erect lanceolato-acuminatis ſtriato-nervoſis, erectis, ſetis tuberculatis. *Hedw. muſc.*

Hypnum ſericeum. *Sp. pl.* p. 1595. *Ho* p. 74.

Hypnum vulgare ſericeum recurvum, erectis cuſpidatis. *Dill. muſc.* 323. *t.* 4

Ad terram, ſaxa, arborum truncos

NECKERA. Capſula oblonga. Periſtomium exterius dentibus ſedecim, acutis; ciliis ſedecim diſtinctis, cum den terioribus alternantibus.

Maſc. gemmacei in diſtinctis plan

pennata. 1. N. fronde ramoſa pinnata, foliis bifari tis reſime acuminatis, capſula perichæ ſa. *Hedw. muſc.* 3. *t.* 19.

Hypnum caulibus diſtichis, ramoſis, transverſim undoſis. *Hall. hiſt. n.* 1797

Hypnum pennatum. *Hoffm. fl.* 2. p.

Fontinalis pennata. *Sp. pl.* p. 1571.

Sphagnum pennatum undulatum, vag ſa. *Dill. muſc.* 250. *t.* 32. *f.* 9.

Ad arborum truncos.

criſpa. 2. N. frondibus ramoſis pinnatis: fol planis. *Sp. pl.* p. 1589. *Hoffm. fl.* Hypno.

Hypnum caule plano, foliis diſtichis, ovatis, transverſim undulatis, operculis ariſtatis. *Hall. hiſt. n.* 1769. *t.* 45. *f.* 5.

Hypnum pennatum & undulatim crispum, ſetis & capſulis brevibus. *Dill. musc.* 273. *t.* 36. *f.* 12.

Ad rupes, arborum truncos.

viticuloſa. 3. N. ſurculis erectis ſubramoſis flexuoſis, foliis patulis. *Sp. pl. p.* 1592. *Hoffm. fl.* 2. *p.* 67. ſub Hypno.

Hypnum ſubhirſutum, viticulis gracilibus erectis, capſulis teretibus. *Dill. musc.* 307. *t.* 39. *f.* 43.

In ſylvaticis ad arborum truncos, ſaxa.

BUXBAUMIA. Capſula ovata, obliqua, baſi hinc ventricoſa. Periſtomium duplex: exterius dentibus ſedecim, truncatis; interius membranaceum, plicatum.

Masc. discoidei.

folioſa. 1. B. acaulis, capſula ſubſeſſili foliis cincta. *Syſt. veget. p.* 945. *Web. ſpic. p.* 128. *Fl. dan. t.* 249. *f.* 3. *Hedw. fund. t.* 9. *f.* 51. *Hoffm. fl.* 2. *p.* 21.

Sphagnum ſeſſile, foliis radicalibus obtuſis, centralibus ciliatis. *Hall. hiſt. n.* 1725. *t.* 46. *f.* 3.

Bryum phascoides. *Wulfen in Iacqu. collect. tom.* 2. *p.* 220.

Bryum hallerianum. *Neck. meth. p.* 233.

Webera Diphiscium. *Ehrh. beytr. tom.* 1. *p.* 189.

Sphagnum acaulon maximum, foliis in centro ciliaribus. *Dill. musc.* 253. *t.* 32. *f.* 13.

In Carinthiæ, Tyrolis montanis, alpinis, & in agro vindobonenſi in ſylva montis *Gallitzinberg.*

aphylla. 2. B. aphylla, pedunculo elongato, capſula latere depreſſa marginata. *Hoffm. fl.* 2. *Syſt. veget. p.* 945. *Hedw. fund. t.* 9. *f.* 3. *f.* 10. *Fl. dan. t.* 44. *Wulfen in Iacq* *lect. tom.* 2. *p.* 213. *Hall. hiſt. n.* 1728.

In Carinthia in ſylvis prope *Clagenf* in monte Calvariæ, prope *Keſſin*, alibiqu umbroſo. *Wulfen.*

POLYTRICHUM. Capſula oblonga, interdum drangula, apophyſi quadrangulæ in Periſtomium duplex: exterius dentibı ginta duobus, brevibus, baſi coalit curvis; interius membrana plana, verſa, apicibus dentium exterioris tinata. Calyptra piloſa.

Masc. discoidei.

commune. 1. P. capſulis parallelepipedis, calyptris foliis lineari-lanceolatis ſerrulatis, caule ci. *Sibthorp. oxon. p.* 305. *Sp. pl. p.* 1573. *ſpic. p.* 73. *Hoffm. fl.* 2. *p.* 23.

Polytrichum quadrangulare vulgare, ync ſerratis. *Dill. musc.* 420. *t.* 54. *f.* 1.

In ſylvis.

juniperinum. 2. P. capſulis parallelepipedis, calyptris foliis lineari-lanceolatis integerrimis mu caule ſimplici. *Sibthorp oxon. p.* 306.

Polytrichum quadrangulare juniperi foliis ribus & rigidioribus. *Dill. musc.* 424. *t.* *Schreb. ſpic. p.* 73.

In ſterilibus ſylvaticis.

piliferum. 3. P. capſulis parallelepipedis, calyptris foliis integerrimis piliferis, caule ſimpl *thorp. oxon. p.* 306. *Schreb. ſpic. p.* 74 *fl.* 2. *p.* 24.

Polytrichum quadrangulare minus juniperi foliis piloſis. *Dill. muſc.* 426. *t.* 54. *f.* 3.

In montanis ſylvaticis, alpinis.

alpinum. 4. P. caule ramoſo, foliis ſerratis, capſula ovata obliquata. *Hoffm. fl.* 2. *p.* 24. *Sp. pl. p.* 1573.

Polytrichum alpinum ramoſum, capſulis e ſummitate ellipticis. *Dill. muſc.* 427. *t.* 55. *f.* 4.

In alpinis, & ſubalpinis.

urnigerum. 5. P. caule ramoſo, foliis rigidis ſubſerratis, capſula oblonga erecta. *Hoffm. fl.* 2. *p.* 24. *Sp. pl. p.* 1573.

Polytrichum ramoſum, ſetis ex alis urnigeris. *Dill. muſc.* 427. *t.* 55. *f.* 5.

Cum præc.

aloides. 6. P. foliis lineari-lanceolatis planis, apice ſerratis, capſula cylindrica obliquata. *Hedw. muſc.* 1. *t.* 14. *Hoffm. fl.* 2. *p.* 24. *Sibthorp oxon. p.* 307.

Polytrichum aloes folio ſerrato, capſulis oblongis. *Dill. muſc.* 429. *t.* 55. *f.* 7.

Cum præc.

nanum. 7. P. foliis linearibus carinatis ſiccitate incurvis convexis obſcure ſerratis, capſula ſubrotunda demum nutante. *Hedw. muſc.* 1. *t.* 13. *Sibthorp oxon. p.* 306.

Polytrichum ericoides. *Hoffm. fl.* 2. *p.* 24.

Polytrichum ſubrotundum. *Curtis lond.*

Polytrichum nanum capſulis ſubrotundis galeritis, aloes folio non ſerrato. *Dill. muſc.* 428. *t.* 55. *f.* 6.

In montanis ſylvaticis.

hercynicum. 8. P. foliis lanceolato-linearibus concavis incurvis, capſula erecta urceolata. *Hedw. muſc.* 1. *t.* 15.

Orthotrichum hercynicum. *Hoffm. fl.* 2. *p.* 25.

In alpibus judenburgenſibus.

undulatum. 9. B. capſula cylindrica cernua, calyptris gla foliis lanceolatis undulatis ſerratis. *Hedw.* 1. *t.* 16. 17. *Sibthorp oxon. p.* 307.

Bryum foliis lanceolatis, ſerratis, capſuli lindricis inclinatis, ariſtatis. *Hall. hiſt. n.*

Bryum undulatum. *Sp. pl. p.* 1582. *Hoffm. p.* 40. *t.* 1. *Weis crypt. p.* 196. *Web. p.* 101.

Bryum phyllitidifolium. *Neck. meth. p.* 20

Bryum Phyllitidis folio rugoſo acuto, ca incurvis. *Dill. muſc.* 360. *t.* 46. *f.* 18.

In ſylvis.

HEPATICÆ.

MARCHANTIA. Receptaculum peltatum, infern miniferum multicapſulare. Capſula qu valvis. Semina filo inſtructa.

polymorpha. 1. M. receptaculo communi decemfido. *Sib oxon. p.* 313. *Sp. pl. p.* 1603.

Lichen fontanus major, ſtellatus æque, ac bellatus, & cyathophorus. *Dill. muſc.* 5 76. *f.* 6.

In umbroſis, ad aquas.

hemisphærica. 2. M. calyce communi quinquefido hemisph perichætio nullo. *Sp. pl. p.* 1604. *Fl. t.* 762.

Lichen pileatus parvus, foliis crenatis, *muſc.* 519. *t.* 75. *f.* 2.

Cum præc.

conica. 3. M. receptaculo communi quinque capſul nico. *Sibthorp. oxon. p.* 314. *Sp. pl. p.*

Lichen vulgaris major pileatus & verr *Dill. muſc.* 516. *t.* 75. *f.* 1.

In umbroſis humidis, ad fontes, riv

IUNGERMANNIA. Capſula ſolitaria, quadrivalvis, ſemina filo inſtructa.

* *Frondibus pinnatis ſecundis.*

aſplenioides. 1. I. frondibus ſimpliciter pinnatis: foliolis ovatis ſubciliatis. *Sp. pl. p.* 1597. *Sibthorp oxon. p.* 308. *Hoffm. fl.* 2. *p.* 87. *Fl. dan. t.* 1061.

Lichenaſtrum aſplenii facie, pinnis laxioribus. *Dill. muſc.* 482. *t.* 69. *f.* 5.

Lichenaſtrum aſplenii facie, pinnis conferticribus. *Dill. muſc.* 483. *t.* 69. *f.* 6.

In umbroſis ſylvaticis.

viticuloſa. 2. I. frondibus ſimpliciter pinnatis, foliolis ſubrotundis integerrimis. *Sibthorp oxon. p.* 308. *Sp. pl. p.* 1597. *Hoffm. fl.* 2. *p.* 88.

Lichenaſtrum trichomanis facie, e baſi & medio florens. *Dill. muſc.* 484. *t.* 69. *f.* 7.

In humidis ſylvaticis.

polyanthos. 3. I. frondibus ſimpliciter pinnatis, foliolis integerrimis imbricatis convexis. *Sp. pl. p.* 1597. *Hoffm. fl.* 2. *p.* 88.

Lichenaſtrum Trichomanis facie, polyanthemum, breve & repens. *Dill. muſc.* 486. *t.* 70. *f.* 9.

In ſylvis umbroſis.

bidentata. 4. I. frondibus ſimpliciter pinnatis apice floriferis, foliolis bidentatis. *Sp. pl. p.* 1598. *Hoffm. fl.* 2. *p.* 89.

Lichenaſtrum pinnulis acutioribus & concavis bifidis, majus. *Dill. muſc.* 487. *t.* 70. *f.* 11.

In umbroſis, ad fluvios.

bicuſpidata. 5. I. frondibus ſimpliciter pinnatis medio floriferis, foliolis bidentatis. *Sp. pl. p.* 1598. *Hoffm. fl.* 2. *p.* 89.

Lichenaſtrum pinnulis acutiſſime bifidis, mi
mum. *Dill. muſc.* 288. *t.* 70. *f.* 13.

In humidis.

ſcalaris. 6. I. frondibus ſimpliciter pinnatis: foliolis integ
rimis ovatis, extremo caule diſtichis alteri
globulifero. *Hoffm. fl.* 2. *p.* 89.

Mnium Trichomanis facie, foliolis integ
Dill. muſc. 236. *t.* 31. *f.* 5.

In aſcenſu alpium judenburgenſium ad
borum radices.

*** Frondibus pinnatis: pinnulis au
culatis ſuperne tectis.*

undulata. 7. I. frondibus ſupra bipinnatis apice floriferis,
liolis ſubrotundis integerrimis. *Sp. pl. p.* 1
Hoffm. fl. 2. *p.* 85.

Lichenaſtrum pinnulis auriculatis majoribu
non crenatis. *Dill. muſc.* 490. *t.* 71. *f.* 17

In ſylvis montanis, ſubalpinis, ad flu

nemoroſa. 8. I. frondibus ſupra bipinnatis apice floriferi
liolis ciliatis. *Sp. pl. p.* 1598. *Hoffm.*
p. 85.

Lichenaſtrum auriculatum, pinnis minoribu
natis. *Dill. muſc.* 490. *t.* 71. *f.* 18.

In ſylvis umbroſis.

albicans. 9. I. frondibus ſupra bipinnatis apice florifer
liolis linearibus recurvatis. *Sp. pl. p.*
Hoffm. fl. 2. *p.* 84.

Lichenaſtrum auriculatum, pinnulis anguſt
nis recurvis. *Dill. muſc.* 492. *t.* 71. *f.* 2

In umbroſis humidis.

trilobata. 10. I. frondibus ſubtus bipinnatis, foliolis qu
obſolete trilobis. *Sp. pl. p.* 1599.

Lichenaſtrum pinnulis obtuſe trifidis, nervo geniculato. *Dill. muſc.* 493. *t.* 71. *f.* 22.

In ſylvis, Styriæ, Carnioliæ montanis, ſubalpinis.

reptans. 11. I. frondibus ſubtus bipinnatis apice reptantibus, foliolis quadridentatis. *Sp. pl. p.* 1599. *Hoffm. fl.* 2. *p.* 86.

Lichenaſtrum multifidum exiguum ad baſin florens per ſiccitatem imbricatum. *Dill. muſc.* 494. *t.* 71. *f.* 24.

In humidis ſylvaticis, ad fluvios.

*** *Frondibus imbricatis.*

complanata. 12. I. ſurculis repentibus: foliolis inferne auriculatis duplicato-imbricatis, ramis æqualibus. *Sp. pl. p.* 1600. *Hoffm. fl.* 2. *p.* 85.

Lichenaſtrum imbricatum majus, ſquamis compreſſis & planis. *Dill. muſc.* 496. *t.* 72. *f.* 26.

Ad arborum truncos.

dilatata. 13. I. ſurculis repentibus, foliis inferne auriculatis duplicato-imbricatis, ramis apice latioribus. *Sp. pl. p.* 1600. *Hoffm. fl.* 2. *p.* 85.

Lichenaſtrum imbricatum minus, ſquamis convexo-concavis. *Dill. muſc.* 497. *t.* 72. *f.* 27.

Ad arborum truncos.

tamariscifolia. 14. I. foliis imbricatis ſerie duplici: ſuperioribus ſubrotundis convexis obtuſis quadruplo-majoribus. *Sp. pl. p.* 1600. *Hoffm. fl.* 2. *p.* 86.

Lichenaſtrum imbricatum, tamariſci narbonenſis facie. *Dill. muſc.* 499. *t.* 72. *f.* 31.

Ad arborum truncos, rupes.

platyphylla. 15. I. ſurculis procumbentibus ſubtus imbricatis, foliolis cordatis acutis. *Sp. pl. p.* 1600. *Hoffm. fl.* 2. *p.* 87.

Lichenaſtrum arboris vitæ facie, fc
rotundis. *Dill. muſc.* 501. *t.* 72. *f.*

In ſylvis.

Tomentella. 16. I. frondibus decompoſitis bipinnatis: mentoſis ciliato-multifidis. *Hoffm. fl*

Lichenaſtrum filicinum pulchrum vill *muſc.* 503. *t.* 73. *f.* 35.

In aſcenſu alpium judenburgenſi borum truncos.

pulcherrima. 17. I. frondibus ſurculisque pinnatis: foli lisque imbricatis apice bi- trifidisv *Web. ſpic. p.* 214. *Hoffm. fl.* 2. *p.*

Lichenaſtrum ſcorpioides pulchrum *Dill. muſc.* 481. *t.* 69. *f.* 3.

Cum præc.

ciliaris. 18. I. frondibus decompoſitis: foliolis du ne imbricatis ciliatis, perianthiis ter *Hoffm. fl.* 2. *p.* 84. *t.* 2.

Ad ſemiputridos arborum truncos.

varia. 19. I. ſurculis erectiuſculis bifariam imbri liolis bipartitis. *Sp. pl. p.* 1601.

Lichenaſtrum foliis variis. *Dill. mu* 73. *f.* 36.

In ſylvis Styriæ ſubalpinis.

**** *Surculis undique imbric foliolis ſparſis.*

emarginata. 20. I. frondibus caulescentibus teretibus ſt ſis: foliolis diſtichis imbricatis conca obtuſe emarginatis. *Hoffm. fl.* 2. *p.*

Cum præc.

julacea. 21. I. frondibus caulescentibus, teretibu preſſis: foliolis arcte imbricatis adpre argenteis. *Hoffm. fl.* 2. *p.* 82. *Sp. pl.*

Lichenaſtrum alpinum Bryi julacei argentei facie. *Dill. muſc.* 506. *t.* 73. *f.* 38.

In alpibus judenburgenſibus.

tricophylla. 22. I. ſurculis teretibus, foliolis capillaceis æqualibus. *Sp. pl. p.* 1601. *Hoffm. fl.* 2. *p.* 82.

Iungermannia multiflora. *Linn. mant. p.* 130. *auct. Cl. Ehrharti.*

Lichenaſtrum trichodes minimum in extremitate florens. *Dill. muſc.* 505. *t.* 73. *f.* 37.

Cum præc.

⁂⁂⁂ *Acaules frondibus ſimplicibus.*

epiphylla. 23. I. acaulis, foliolo frondi innato. *Sp. pl. p.* 1602. *Hoffm. fl.* 2. *p.* 92. *t.* 4.

Lichenaſtrum capitulis rotundis e foliorum medio enaſcentibus. *Dill. muſc.* 508. *t.* 74. *f.* 41.

In umbroſis, ad rivulos, fluvios.

pinguis. 24. I. acaulis, fronde oblonga ſinuata pingui. *Sp. pl. p.* 1602. *Hoffm. fl.* 2. *p.* 92.

Lichenaſtrum capitulis oblongis juxta foliorum diviſuras enaſcentibus. *Dill. muſc.* 509. *t.* 74. *f.* 42.

In udis paluſtribus.

furcata. 25. I. acaulis, fronde lineari-ramoſa extremitatibus furcatis obtuſiuſculis. *Sp. pl. p.* 1602. *Hoffm. fl.* 2. *p.* 90.

Lichenaſtrum tenuifolium furcatum, thecis globoſis piloſis. *Dill. muſc.* 512. *t.* 74. *t.* 45.

Ad arborum truncos, rupes humidas.

BLASIA. Masc. Anthera ſubgloboſa frondi i

Fem. Calyx lagenæformis. Semi
per tubulum exeuntia.

puſilla. 1. Blaſia. *Sp. pl. p.* 1605. *Fl. dan. t.* 45
fl. 2. *t.* 3.

Blaſia puſilla, lichenis pyxidati facie. *M*
p. 14. *t.* 7.

Mnium lichenis facie. *Dill. muſc.* 237. *t*

In umbroſis udis, Styriæ ſuperiori
ad foſſas, fluvios.

RICCIA. Granula frondi innata in quibus (
corolla nulla. Anthera cylindrica
germini turbinato impoſita. Stylus
mis antheram perforans. Pericarpiu
ricum coronatum anthera emarcida,
na hemiſphærica, pedicellata.

cryſtalli- 1. R. frondibus ſuperficie papilloſis. *Sp. pl.*
na.

Riccia minima & minor pinguis aſper
ſtallina perfuſa. *Mich. gen. p.* 107. *t.* 57

Lichen paluſtris, rutæ folio. *Dill. mu*
t. 78. *f.* 12.

In humidis.

fluitans. 2. R. frondibus dichotomis lineari - filif
Sp. pl. p. 1606.

Lichenaſtrum aquaticum fluitans tenuifol
catum. *Dill. muſc.* 213. *t.* 74. *f.* 47.

In aquis ſtagnantibus, vel lente fl

Fructificatio eſt capſula, inferiori paginæ non procul ab apice inſidens, globoſa, evalvis, continens ſemina a vi triginta, alba vel fuſca. Ehrh. beytr. p. 81.

ANTHOCEROS. Maſc. Calyx ſeſſilis, cylindricus, integer. Anthera ſubulata, longiſſima, bivalvis.

Fem. Calyx ſexpartitus. Semina tria.

lævis. 1. A. frondibus indiviſis ſinuatis lævibus. *Sp. pl.* p. 1606.

Anthoceros foliis majoribus laciniatis. *Dill. musc.* 476. *t.* 68. *f.* 2.

In umbroſis, humidis Carniæ. *I. Hoſt.*

A L G Æ.

LICHEN. Maſc. Veſiculæ conglomeratæ, minutiſſimæ, farinæ inſtar congeſtæ, ſeu ſparſæ in diſco, margine, vel apicibus frondium.

Fem. Receptaculum ſubrotundum, planiuſculum, convexum, (*Tuberculum*) concavum (*Scutella*) ſubrevolutum, margine adfixum (*Pelta*) ſæpe colore a fronde diverſum, intus continens ſemina in ſeries digeſta.

* *Pulverulenti.*

antiquitatis. 1. L. pulverulentus ater. *Sp. pl. p.* 1638. ſub Byſſo.

Ad muros vetuſtos, ſaxa.

ſaxatilis. 2. L. pulverulentus cinereus. *Sp. pl. p.* 1638. ſub B.

Obducit ſaxa.

flavus. 3. L. pulverulentus flavus. *Sibthorp oxon. p.* 315.

Byſſus candelaris. *Sp .pl. p.* 1639.

Byſſus pulverulenta flava lignis adnaſcens. *Dill. musc.* 3. *t.* 1. *f.* 5.

In muris, arborum cortice.

botryoides. 4. L. pulverulentus viridis. *Sp. pl. p*
Byſſo.

Byſſus botryoides ſaturate virens. *Di*
t. 1. *f.* 5.

Obducit terram humidam, umbr

incanus. 5. L. pulverulentus incanus. *Sp. pl. p.* 1

Byſſus pulverulenta incana, farinæ i
Dill. muſc. 3. *t.* 1. *f.* 3.

Ad foſſas, lacus, & in aliis hum

lacteus. 6. L. pulverulento-cruſtaceus albiſſimus
1639. ſub B.

Byſſus candidiſſima, calcis inſtar muſ
Dill. muſc. 2. *t.* 1. *f.* 2.

In muſcis, arborum corticibus.

** *Leproſi tuberculati.*

ſcriptus. 7. L. leproſus albicans, lineolis nigris r
racteriformibus. *Sp. pl. p.* 1606.

Lichenoides cruſta tenuiſſima, pereg
litteris inſcripta. *Dill. muſc.* 125. *t.* 1

In arborum cortice.

geographicus. 8. L. leproſus flaveſcens, lineolis nigr
referens. *Sp. pl. p.* 1607.

Verrucaria geographica. *Hoffm. lich.*

Lichenoides nigro-flavum, tabulæ g
inſtar pictum. *Dill. muſc.* 126. *t.* 18.

Obducit rupes alpinas.

ſanguinarius. 9. L. leproſus cinereo-viteſcens, tuber
Sp. pl. p. 1607.

Verrucaria ſaguinaria. *Hoffm. lich. t.*

In arborum cortice.

fuſco-ater. 10. L. leproſus fuſcus, tuberculis atris. *Sp. pl. p.* 1607. *Wulfen in Iacqu. collect. tom.* 2. *p.* 230. *t.* 14. *f.* 3.

Verrucaria fuſco-atra. *Hoffm. lich. t.* 54. *f.* 1.

In rupibus.

calcareus. 11. L. leproſus candidus, tuberculis atris. *Sp. pl. p.* 1607.

Lichenoides tartareum tinctorium candidum, tuberculis atris. *Dill. muſc.* 128. *t.* 18. *f.* 8.

In rupibus calcareis Litoralis.

immerſus. 12. L. calcareus albidus, verrucis immerſis nigris. *Hoffm. lich. t.* 12. *f.* 2. 3. 4. ſub Verrucaria. *Web. ſpic. p.* 188.

Lichen ſcutis in ſaxum immerſis, nigris. *Hall. hiſt. n.* 2076.

In rupibus.

petræus. 13. L. cruſtaceus ſubcinereus ſcutellis nigris depreſſis concentricis. *Hoffm. lich. t.* 50. *f.* 1. 2. ſub V. *Wulfen in Iacqu. collect. tom.* 3. *p.* 116. *t.* 6. *f.* 2.

Ad ſaxa.

albo-ater. 14. L. leproſus albus, verrucis nigris. *Hoffm. lich. t.* 15. *f.* 2. ſub V.

In arborum cortice.

purpuraſcens. 15. L. tenuiſſimus læte purpureus lineis ſanguineis terminatus; verrucis immerſis concoloribus. *Hoffm. lich. t.* 15. *f.* 1. ſub V.

Lichen marmoreus. *Scop. carn. tom.* 2. *p.* 367. *Wulfen in Iacqu. collect. tom.* 2. *p.* 178. *t.* 13. *f.* 1.

In rupibus calcareis Carnioliæ, Carinthiæ.

ventoſus. 16. L. leproſus flavus, tuberculis rubris. *Sp. pl. p.* 1607.

Verrucaria ventoſa. *Hoffm. lich. t.* 27 *veget. p.* 957.

Lichen flaveſcens. *Wulfen in Iacqu. m* 2. *p.* 79. *t.* 9. *f.* 1. *Syſt. veget. p.* 9

Lichen cruentus. *Web. ſpic. p.* 184.

In rupibus alpinis.

carpine-us. 17. L. leproſus cinereus, tuberculis albi *Sp. pl. p.* 1608.

Ad arborum truncos.

ærugi-noſus. 18. L. leproſus æruginoſus, tuberculis fe nis carneis. *Sibthorp oxon. p.* 321. *I t.* 275. *Scop. carn. tom.* 2. *p.* 361.

Lichen cruſta verrucoſa, cinerea, ſ neis planis. *Hall. hiſt. n.* 2044.

In ſylvis ſubalpinis ad putrid truncos.

ericeto-rum. 19. L. leproſus candidus, tuberculis inca *pl. p.* 1608.

Lichen cruſta tenace verrucoſa albida carnatis. *Hall. hiſt. n.* 2042.

Coralloides fungiforme carneum baſi le *muſc.* 76. *t.* 14. *f.* 1.

In ſylvaticis ad terram.

*** *Leproſi ſcutellati.*

jolithus. 20. L. leproſus ruber, peltis minutiſſim binatis albidis. *Retzii ſcand. p.* 274.

Byſſus jolithus. *Sp. pl. p.* 1638.

Byſſus germanica minima ſaxatilis odorem ſpirans. *Mich. gen. p.* 210

In montanis, ſubalpinis ad ſa

cinereo-fuscus. 21. L. cruſtaceus cinereus, patellis complanatis rufescentibus tenuiſſime marginatis. *Hoffm. lich. t.* 12. *f.* 1. ſub Patellaria.

In arborum cortice.

albo-cœrulescens. 22. L. calcareus, patellis albo cœrulescentibus, nigro-marginatis. *Hoffm. lich. t.* 14. *f.* 2. ſub P. *Wulfen in Iacqu. collect. tom.* 2. *p.* 184. *t.* 15. *f.* 1.

In Carinthiæ rupibus. *Wulfen.*

flavo-virescens. 23. L. cruſtaceus, flavo-virescens, patellis ochroleucis flavo-marginatis. *Hoffm. lich. t.* 20. *f.* 1. ſub P. *Wulfen in Iacqu. collect. tom.* 2. *p.* 230. *t.* 13. *f.* 4.

In Carinthia in rupibus prope *Schmalzberg. Wulfen.*

tartareus. 24. L. cruſtaceus ex albido virescens, ſcutellis flaveſcentibus: margine albo. *Sp. pl. p.* 1608. *Wulfen in Iacqu. collect. tom.* 4. *p.* 241. *t.* 8. *f.* 2.

Lichenoides tartareum farinaceum, ſcutellarum umbone fusco. *Dill. musc.* 132. *t.* 18. *f.* 12.

Ad ſaxa.

rimoſus. 25. L. cruſtaceus albicans, fruſtulis ſubrotundo-angulatis, ſcutellis glaucis albo-marginatis. *Retzii ſcand. p.* 275. *Fl. dan. t.* 468. *f.* 3.

Patellaria Glaucoma. *Hoffm. lich. t.* 52. *f.* 1. 2. 3. *t.* 53. *f.* 4. 5. 6.

In rupibus alpium judenburgenſium.

ſubfuscus. 26. L. cruſtaceus albicans, ſcutellis ſubfuscis: margine cinereo ſubcrenato. *Sp. pl. p.* 1609.

Lichenoides cruſtaceum & leproſum, ſcutellis ſubfuscis. *Dill. musc.* 134. *t.* 18. *f.* 16.

Ad arborum corticem, muros, ſaxa.

muscorum. 27. L. cruſtaceus incanus, ſcutellis atris connexis. *Retzii ſcand. p.* 275. *Fl. dan. t.* 1003.

Patellaria muſcorum. *Hoffm. lich. t.* 2:

Muſcis adnaſcitur.

Parellus. 28. L. cruſtaceus albus, peltis concavis obtı dis. *Syſt. veget. p.* 958.

Patellaria Perella. *Hoffm. lich. t.* 12. *f.*

In rupibus Litoralis.

upſalienſis. 29. L. cruſtaceus, foliis ſubulatis ſtriatis, *p.* 1609.

Patellaria *upſalienſis*, cruſta ex alienis foliolis, verrucoſa albida, patellæ c punctatæ. *Hoffm. lich. t.* 21. *f.* 2.

In alpibus judenburgenſibus adnaſci tuis graminum ceſpitibus.

ſcruposus. 30. L. cruſtaceus cinereus granoſus, ſcutell ſis atris, margine crenulato. *Schreb. ſpi*

Patellaria ſcrupoſa. *Hoffm. lich. t.* 11.

Lichen cruſta tenace alba, ſcutellis atris. *Hall. hiſt. n.* 2051.

Ad ſaxa.

veſicularis. 31. L. polymorphus, tuberculis fungoſis ni reisque mixtis. *Hoffm. lich. t.* 32. *f.* 3

In Auſtriæ, Styriæ montanis ſyl terrram.

**** *Imbricati.*

ſaxicola. 32. L. ſubfoliaceus calcareus, foliolis adpr ſis anguſtis flavis, ſcutellis concoloribı *lich. t.* 17. *f.* 3. ſub Pſora.

Lichen candelaris. *Wulfen in Jacqu.* 3, *p.* 124. *t.* 6. *f.* 1.

Ad muros, ſaxa.

caſius. 33. L. cruſtaceus adpreſſus, albido-cinereus, ſcutellis concoloribus, tuberculis intermixtis cæſiis. *Hoffm. lich. t.* 8. *f.* 1. ſub Pſora.

Ad ſaxa, tecta.

pulveru-lentus. 34. L. adpreſſus, foliolis linearibus obtuſis lobatis crenatis, polline conſperſus, ſcutellis nigricantibus. *Hoffm. lich. t.* 8. *f.* 2. ſub Pſora.

Ad arborum corticem.

ſaxatilis. 35. L. imbricatus, foliolis ſinuatis ſcabris lacunoſis, ſcutellis badiis. *Syſt. veget. p.* 958. *Wulfen in Iacqu. collect. tom.* 4. *p.* 281. *t.* 20. *f.* 2.

Lichenoides vulgatiſſimum cinereo-glaucum lacunoſum & cirrhoſum. *Dill. muſc.* 188. *t.* 24 *f.* 83.

Ad ſaxa, arborum truncos.

ompha-lodes. 36. L. imbricatus, foliolis multifidis glabris obtuſis incanis, punctis vagis eminentibus. *Sp. pl. p.* 1609.

Lichenoides ſaxatile tinctorium, foliis piloſis purpureis. *Dill. muſc.* 185. *t.* 24. *f.* 80.

In Styriæ ſubalpinis ad rupes.

oliva-ceus. 37. L. imbricatus, foliolis lobatis nitidis lividis. *Sp. pl. p.* 1610.

Lichenoides olivaceum, ſcutellis amplioribus verrucoſis. *Dill. muſc.* 184. *t.* 24. *f.* 78.

Ad rupes, arborum truncos.

fahlu-nenſis. 38. L. imbricatus, foliolis linearibus dichotomis planiuſculis acutis nigris, ſcutellis atris. *Sp. pl. p.* 1610. *Web. ſpic. p.* 251. *Wulfen in Iacqu. miſcell. tom.* 2. *p.* 90. *t.* 10. *f.* 2.

Lichen fronde laciniata, undique nigerrima, ſcutellis concoloribus. *Hall. hiſt. n.* 2020.

Squamaria fahlunenſis. *Hoffm. lich. t.* 36. *f.* 2.

Lichenoides tinctorium atrum foliis minimis criſpis. *Dill. muſc.* 188. *t.* 24. *f.* 81.

In rupibus alpinis.

criſpus. 39. L. imbricatus, foliis lobatis truncati atroviridibus, ſcutellis concoloribus. *S p.* 959. *Wulfen in Iacqu. collect. tom. t.* 10. *f.* 1.

Lichenoides gelatinoſum atrovirens (rugoſum. *Dill. muſc.* 139. *t.* 19. *f.* 23

Ad ſaxa.

criſtatus. 40. L. foliis imbricatis dentato-ciliatis, ſ lio majoribus. *Sp. pl. p.* 1610. *Wulfer collect. tom.* 3. *p.* 139. *t.* 12. *f.* 1.

Lichenoides gelatinoſum, foliis imbric ſtatis. *Dill. muſc.* 140. *t.* 19. *f.* 26.

Ad ſaxa, muros vetuſtos.

decipiens. 41. L. tartareus aggregatus latericius, tube ginalibus nigris. *Hoffm. lich. t.* 43. *f.* Pſora. *Hedw. muſc.* 2. *t.* 1. *Dicks. fas*

Lichen elveloides. *Wulfen in Iacqu. co* 8. *p.* 108. *t.* 3. *f.* 3.

Ad terram, ſaxa.

lentigerus. 42. L. calcareus lacteus lobatus: lobis de tellisque albo-marginatis. *Hoffm. lich.* ſub. P.

Ad terram, rupes, muſcos.

parietinus. 43. L. imbricatus, foliis criſpis fulvis, pe loribus. *Sp. pl. p.* 1610. *Fl. dan. t.* 1

Lichenoides vulgare ſinuoſum, foliis luteis. *Dill. muſc.* 180. *t.* 24. *f.* 76.

Ad ſaxa, muros, arbores.

phyſodes. 44. L. imbricatus, laciniis obtuſis ſubinfla *p.* 1610.

Lichenoides ceratophyllum obtuſius & moſum. *Dill. muſc.* 154. *t.* 20. *f.* 49

Ad arborum truncos, aſſeres, ſa

ſtellaris. 45. L. imbricatus, foliolis oblongis laci ſtis cinereis, ſcutellis pullis. *Sp. pl.*

Lichenoides cinereum, ſegmentis argutis ſtellatis, ſcutellis nigris. *Dill. muſc.* 176. *t.* 24. *f.* 70.

Ad arborum truncos, ramos.

***** *Foliacei.*

ciliaris. 46. L. foliaceus erectiuſculus: laciniis linearibus ciliatis, ſcutellis pedunculatis crenatis. *Sp. pl. p.* 1611. *Fl. dan. t.* 711. *Scop. carn. tom.* 2. *p.* 382. *Wulfen in Iacqu. collect. tom.* 4. *p.* 244. *t.* 13. *f.* 1.

Lichen corniculatus, planus, ramulis capillaribus, ſcutellis petiolatis. *Hall. hiſt. n.* 1980.

Lichenoides ciliare. *Hoffm. lich. t.* 3. *f.* 4.

Lichenoides hiſpidum majus & rigidius, ſcutellis nigris. *Dill. muſc.* 150. *t.* 20. *f.* 45.

Ad arborum truncos.

hiſpidus. 47. L. foliaceus albus, laciniis ſubturbinatis ciliato-ſpinuloſis, ſcutellis ſubſeſſilibus. *Retzii ſcand. p.* 281. *Schreb. ſpic. p.* 126. *Wulfen in Iacqu. collect. tom.* 4. *p.* 246. *t.* 6.

Lichen fronde ſemipinnata, canaliculata, cinerea. *Hall. hiſt. n.* 2019.

Lichen tenellus. *Scop. carn. tom.* 2. *p.* 394. *Sibthorp oxon. p.* 329. *Web. ſpic. p.* 235.

Lichenoides hiſpidum. *Hoffm. lich. t.* 3. *f.* 3.

Lichenoides hiſpidum minus & tenerius, ſcutellis nigris. *Dill. muſc.* 152. *t.* 20. *f.* 46.

Ad ſaxa, arborum ramos, truncos.

islandicus. 48. L. foliaceus adſcendens laciniatus, marginibus elevatis ciliatis. *Sp. pl. p.* 1611. *Fl. dan. t.* 155. *Wulfen in Iacqu. collect. tom.* 4. *p.* 253. *t.* 8. *f.* 1.

Lichenoides islandicum. *Hoffm. lich. t.* 9. *f.* 1.

In alpinis, ſubalpinis ad terram.

nivalis. 49. L. ſoliaceus adſcendens laciniatus criſ|
lacunoſus albus, margine elevato.
1612. *Fl. dan. t.* 227.

Lichenoides lacunoſum candidum glab
diviæ criſpæ facie. *Dill. muſc.* 162. *t.*

In alpibus.

pulmonarius. 50. L. foliaceus laciniatus obtuſus glaber,
cunoſus, ſubtus tomentoſus, ſcutellis ı
bus. *Retzii ſcand. p.* 281. *Fl. dan. t.*

Pulmonaria reticulata. *Hoffm. lich. t.*

Lichenoides pulmoneum reticulatum
marginibus peltiferis. *Dill. muſc.* 2ı
f. 113.

Ad arborum truncos.

furfuraceus. 51. L. foliaceus decumbens furfuraceus: la
tis: ſubtus lacunoſis atris. *Sp. pl. p.*

Lichen cornubus planiuſculis ovatis, ſ
rinoſis, inferne atris & reticulatis.
n. 1976.

Lichenoides furfuraceum. *Hoffm. lich.*

In ſubalpinis ad arborum truncos

farinaceus. 52. L. foliaceus erectus compreſſus ramo
rucis marginalibus farinoſis. *Syſt. veg.*

Lichenoides ſegmentis anguſtioribus
nes verrucoſis & pulverulentis. *Dill.*
t. 23. *f.* 63.

Ad arborum truncos.

calicaris. 53. L. foliaceus erectus linearis ramoſus
convexus mucronatus. *Sp. pl. p.* 1613

Lichenoides coralliforme roſtratum
culatum. *Dill. muſc.* 170. *t.* 23. *f.* 6

Ad rupes, arborum truncos.

fraxineus. 54. L. foliaceus erectus oblongus lanceolatus ſublaciniatus lacunoſus glaber, ſcutellis ſubpedunculatis. *Sp. pl. p.* 1614.

Lichen utrinque lacunatus, fronde latiſſima, ſcutellis creberrimis. *Hall. hiſt. n.* 1985.

Platisma fraxineum. *Hoffm. lich. t.* 18. *f.* 1. 2.

Lichenoides longifolium rugoſum rigidum. *Dill. musc.* 165. *t.* 22. *f.* 59.

Ad arborum truncos.

prunastri. 55. L. foliaceus erectiusculus lacunoſus: ſubtus tomentoſus albus. *Sp. pl. p.* 1614.

Lichenoides cornutum, bronchiale molle ſubtus incanum. *Dill. musc.* 160. *t.* 21. *f.* 55.

Ad arborum truncos.

juniperinus. 56. L. foliaceus laciniatus crispus fulvus, peltis lividis. *Sp. pl. p.* 1614. *Fl. dan. t.* 1004.

Squamaria juniperina. *Hoffm. lich. t.* 7. *f.* 2.

Ad Pini, juniperi truncos.

caperatus. 57. L. pallide viridis rugoſus margine undulatus. *Sp. pl. p.* 1614. *Wulfen in Iacqu. collect. tom.* 4. *p.* 280. *t.* 20. *f.* 1.

Lichen fronde rotunde lobata, rugoſa, inferne atra & aſpera. *Hall. hiſt. n.* 2005.

Platisma caperatum. *Hoffm. lich. t.* 38. 39. 40. *f.* 1.

Lichenodes caperatum, roſacee expanſum, e ſulphureo virens. *Dill. musc.* 193. *t.* 25. *f.* 97.

Ad ſaxa, arborum truncos.

glaucus. 58. L. foliaceus depreſſus lobatus glaber: margine crispo farinaceo. *Sp. pl. p.* 1615.

Lichen fronde laciniata & crispa, inferne atra & aſpera. *Hall. hiſt. n.* 2007.

Lichenoides endiviæ foliis crispis ſplendentibus ſubtus nigricantibus. *Dill. musc.* 192. *t.* 25. *f.* 96.

In ſubalpinis ad arborum truncos.

fallax. 59. L. foliaceus diffuſus, laciniatus criſpus, ſ lis terminalibus. *Web. ſpic. p.* 244.

Lichen glaucus. *Wulfen in Iacqu. collect.* 4. *p.* 276. *t.* 19. *f.* 2.

Platisma fallax. *Hoffm. lich. t.* 46. *f.* 1.

Lichenoides membranaceum, tubæ fall æmulum. *Dill. muſc.* 165. *t.* 22. *f.* 58.

Lichen alpinus, membranaceus, elegans, plas lacinias diviſus, interne albus, ſup glauco ſubvireſcens, receptaculis florum a ribus, interna parte fuſcis. *Mich. gen. t.* 37.

In Carinthia, Carnia, Forojulio, I ad arborum truncos.

nigreſcens. 60. L. foliaceus gelatinoſus ſubrotundus loba goſus atrovirens, ſcutellis rufis. *Syſt. p.* 961.

Lichen papyraceus. *Wulfen in Iacqu. tom.* 3. *p.* 134. *t.* 10. *f.* 3.

Lichen Lactuca. *Web. ſpic. p.* 252.

Collema veſpertilio. *Hoffm. lich. t.* 37.

Lichenoides gelatinoſum membranaceum nigricans. *Dill. muſc.* 138. *t.* 19. *f.* 20.

Ad ſaxa, arborum truncos.

granoſus. 61. L. gelatinoſus ſinuato-plicatus, lobis culis rotundatis, tuberculis ſphæroidibu noſis aggregatis. *Retzii ſcand. p.* 284. *in Iacqu. collect. tom.* 3. *p.* 131. *t.* 1 *Fl. dan. t.* 462. *f.* 1. *Scop. carn. tom.* 2

Lichenoides gelatinoſum atro-virens tum, & granoſum. *Dill. muſc.* 140. *t.*

Ad ſaxa, terram, muſcos.

ſcrobiculatus. 62. L. coriaceus repens lobatus obtuſus ſup biculatus verrucoſus, inferne villoſus *Retzii ſcand. p.* 284. *Scop. carn. tom.*

Lichen fronde obtuſe lobata, inferne ſcabra, maculis rotundis varia. *Hall. hiſt.* n. 1989.

Lichen verrucoſus. *Huds. angl. ed.* 2. 545. *Wulfen in Iacqu. collect.* tom. 4. p. 276. t. 18. *f.* 2.

Pulmonaria verrucoſa. *Hoffm. lich.* t. 1. *f.* 1.

Lichenoides pulmoneum villoſum, ſuperficie ſcrobiculata & peltata. *Dill. muſc.* 216. t. 29. *f.* 114.

In montanis, ſubalpinis ad arborum truncos & radices, ad rupes.

Coriacei.

aquaticus. 63. L. ſubcoriaceus imbricatus lobatus: lobis obtuſis. *Weis crypt.* p. 77.

Lichen fluviatilis. *Web. ſpic.* p. 265. t. 4.

Platisma aquaticum. *Hoffm. lich.* t. 45. *f.* 1–5.

Lichenoides imbricatum, luridum. *Dill. muſc.* 224. t. 30. *f.* 128.

In alpibus judenburgenſibus ad rivulorum ſaxa.

reſupinatus. 64. L. coriaceus repens lobatus, peltis marginalibus poſticis. *Sp. pl.* p. 1615. *Wulfen in Iacqu. collect.* tom. 4. p. 257. t. 12. *f.* 1.

Lichenoides fuſcum, peltis poſticis ferrugineis. *Dill. muſc.* 206. t. 28. *f.* 105.

In ſubalpinis, alpinis ad terram.

venoſus. 65. L. coriaceus repens ovatus planus, ſubtus venoſus villoſus, peltis marginalibus horizontalibus. *Sp. pl.* p. 1615.

Lichen flabelliformis, ſubtus reticulatus, ſcutellis marginalibus planis. *Hall. hiſt.* n. 1993.

Peltigera venoſa. *Hoffm. lich.* t. 6. *f.* 2.

Lichenoides parvum virescens, peltis
bus planis. *Dill. muſc.* 208. *t.* 28. *f.*

In ſylvis ad terram.

aphtoſus. 66. L. foliaceus repens lobatus obtuſus pl
rucis ſparſis, pelta marginali adſcend
pl. p. 1616. *Fl. dan. t.* 762. *f.* 1.
Iacqu. collect. tom. 4. *p.* 266. *t.* 17.
tom. 2. *p.* 384.

Lichen fronde obtuſe lobata, ſuperne
inferne villoſa. *Hall. hiſt. n.* 1992.

Peltigera aphtoſa. *Hoffm. lich. t.* 6.

Lichenoides digitatum læte virens, v
gris notatum. *Dill. muſc.* 207. *t.* 28.

Cum præc.

caninus. 67. L. coriaceus repens lobatus obtuſ
ſubtus venoſus villoſus, pelta margin
dente. *Sp. pl. p.* 1616. *Fl. dan. t.* 76
in Iacqu. collect. tom. 4. *p.* 260. *t.*
Scop. carn. tom. 2. *p.* 383.

Lichenoides digitatum cinereum, lac
ſinuoſis. *Dill. muſc.* 200. *t.* 27. *f.* 1

Cum præc.

ſylvaticus. 68. L. coriaceus repens laciniatus lacun
marginalibus adſcendentibus. *Syſt. ve*
Wulfen in Iacqu. collect. tom. 4. *p.*
f. 2.

Lichen lacunatus exaſperatus, infe
& ſcutellifer. *Hall. hiſt. n.* 1987.

Peltigera ſylvatica. *Hoffm. lich. t.* 4

Lichenoides polyſchides villoſum
peltis parvis. *Dill. muſc.* 199. *t.* 9

Ad rupes, arborum truncos.

polydactylus. 69. L. coriaceus repens lobatus: lobis d
tis marginalibus adſcendentibus.
p. 285. *Wulfen in Iacqu. collect.*
262. *t.* 14. *f.* 2. *Leers herb. p.*
meth. p. 85.

Peltigera polydactylon. *Hoffm. lich. t. 4. f.* 1.

Lichenoides cynereum polydactylon. *Dill. muſc.* 207. *t.* 28. *f.* 107.

In ſylvaticis ad terram.

perlatus. 70. L. coriaceus repens lobatus glaber, ſubtus ater, ſcutellis pedunculatis integris. *Syſt. veget. p.* 962. *Wulfen in Iacqu. collect. tom.* 4. *p.* 273. *t.* 10.

Lichenoides glaucum perlatum ſubtus nigrum & cirrhoſum. *Dill. muſc.* 147. *t.* 20. *f.* 39.

Ad arborum truncos, radices, ad rupes.

horizontalis. 71. L. coriaceus repens planus ſubtus avenius, peltis marginalibus horizontalibus. *Syſt. veget. p.* 961. *Fl. dan. t.* 533. 765. *Wulfen in Iacqu. collect. tom.* 4. *p.* 265. *t.* 16.

Lichenoides ſubfuſcum, peltis horizontalibus. *Dill. muſc.* 205. *t.* 28. *f.* 104.

In ſylvis ad terram.

ſaccatus. 72. L. coriaceus repens ſubrotundus: peltis depreſſis ſubtus ſaccatis. *Sp. pl. p.* 1616. *Fl. dan. t.* 532. *f.* 3.

Lichenoides Lichenis facie, peltis acetabulis immerſis. *Dill. muſc.* 221. *t.* 30. *f.* 121.

In ſylvaticis ad terram.

croceus. 73. coriaceus repens ſubrotundus planus, ſubtus venoſus villoſus croceus, peltis ſparſis adnatis. *Sp. pl. p.* 1616. *Wulfen in Iacqu. collect. tom.* 4. *p.* 255. *t.* 11. *f.* 2. 3.

Lichen fronde rotunde lobata, plana, ſubtus venoſa, miniata. *Hall. hiſt. n.* 1994.

Peltigera crocea. *Hoffm. lich. t.* 41. *f.* 2. 4. *t.* 42. *f.* 4. 5.

Lichenoides ſubtus croceum, peltis adpreſſis. *Dill. muſc.* 221. *t.* 29. *f.* 120.

In ſummis Styriæ, Carinthiæ, Tyrolis alpium jugis ad terram.

ſarcoides. 74. L. coriaceus gelatinoſus carneus, ſcu
bus. *Syſt. veget. p.* 962. *Iacqu. misc
p.* 378. *t.* 22.

In ſylvis Auſtriæ ſubalpinis ad arl
cos. *Iacqu.*

******* *Umbilicati.*

miniatus. 75. L. umbilicatus gibbus punctatus: ſul
Sp. pl. p. 1617. *Fl. dan. t.* 532. *f.
in Iacqu. miscell. tom.* 2. *p.* 90. *t.*

Lichenoides coriaceum, nebuloſum,
punctatum, ſubtus fulvum. *Dill. mu*
30. *f.* 127.

In alpinis, ſubalpinis ad rupes.

velleus. 76. L. umbilicatus: ſubtus hirſutiſſimus
p. 1617.

Lichen fronde coriacea, inferne puſt
loſa, ſcutellis atris. *Hall. hiſt. n.* 19

Lichenoides coriaceum, latiſſimo foli
to & verrucoſo. *Dill. musc.* 545. *t.*

In rupibus alpium judenburgenſiu

puſtulatus. 77. L. umbilicatus: ſubtus lacunoſus, fur
adſperſus. *Syſt. veget. p.* 962. *Fl. da
f.* 2. *Wulfen in Iacqu. collect. tom.*

Lichen fronde cinerea, inferne lacı
busculis ſuperne nigris. *Hall. hiſt. n.*

Umbilicaria puſtulata. *Hoffm. lich. t.*

Lichenoides puſtulatum cinereum &
buſtum. *Dill. musc.* 226. *t.* 31. *f.* 13

In Styriæ, Carinthiæ, Tyrolis ru
pinis, ſubalpinis.

proboscideus. 78. L. umbilicatus laciniatus lævis, lacinii
ramoſo ſetaceo - ciliatis, peltis turbina
dunculatis. *Retzii ſcand. p.* 288. *Sp. p
Wulfen in Iacqu. collect. tom.* 2. *p.
f.* 2. *Fl. dan. t.* 471. *f.* 1. 2. *Hedw.
t.* 1.

Umbilicaria crinita. *Hoffm. lich.* t. 44. f. 1–9.

Lichenoides corneum, marginibus eleganter fimbriatis. *Dill. muſc.* 218. t. 29. f. 116.

In rupibus montanis, ſubalpinis, alpinis.

deuſtus. 79. L. umbilicatus undique lævis. *Sp. pl.* p. 1618.

Lichenoides coriaceum cinereum, peltis ater compreſſis. *Dill. muſc.* 219. t. 30. f. 117.

In rupibus alpinis.

polyphyllus. 80. L. umbilicatus polyphyllus undique lævis atrovirens crenatus. *Syſt. veget.* p. 962.

Lichenoides tenue pullum, foliis utrinque glabris, *Dill. muſc.* 225. t. 30. f. 129.

Cum præc.

complicatus. 81. L. umbilicatus polyphyllus imbricatus lobatus punctatus fuſco-viridis. *Retzii ſcand.* p. 288.

Lichen polyphyllus. *Wulfen in Iacqu. collect. tom.* 2. p. 190. t. 16. f. 1.

Lichenoides imbricatum luridum, *Dill. muſc.* 224. t. 30. f. 128.

In rupibus alpinis Styriæ, Carinthiæ, Tyrolis.

******** *Scyphiferi.*

cocciferus. 82. L. ſcyphifer ſimplex integerrimus, ſtipite cylindrico, tuberculis coccineis. *Sp. pl.* p. 1618.

Coralloides ſcyphiforme, tuberculis coccineis. *Dill. muſc.* 82. t. 14. f. 7.

In ſylvaticis ad terram, ſaxa.

pyxidatus. 83. L. ſcyphifer ſimplex crenulatus, tuberculis fuſcis. *Sp. pl.* p. 1619.

Coralloides ſcyphiforme, tuberculis fuſcis. *Dill. muſc.* 79. t. 14. f. 6.

In ſylvis ad terram.

fimbriatus. 84. L. ſcyphifer ſimplex denticulatus, ſt drico. *Sp. pl. p.* 1619.

Coralloides ſcyphiforme gracile, mai tatis. *Dill. muſc.* 84. *t.* 14. *f.* 8.

Cum præc.

gracilis. 85. L. ſcyphifer ramoſus denticulatus fili *pl. p.* 1619.

Coralloides ſcyphiforme ſerratum e libus gracilibus glabris. *Dill. muſc* *f.* 13.

Cum præc.

cornutus. 86. L. ſcyphifer ſimpliciuſculus ſubvent lycibus integris. *Sp. pl. p.* 1620.

Cladonia cornuta. *Hoffm. lich. t.* 2

Coralloides vix ramoſum, ſcyphis ol *muſc.* 90. *t.* 15. *f.* 14.

In ſylvaticis ad terram.

radiatus. 87. L. ſcyphifer ſubramoſus cylindraceu calycibus inæqualibus dentatis radiati *ſpic. p.* 122. *Retzii ſcand. p.* 290.

Coralloides ſcyphiforme cornutum. 92. *t.* 15. *f.* 16.

In ſylvis montanis, ſubalpinis

********* *Fruticuloſi.*

papillaria. 88. L. fruticuloſus perforatus albidus, obtuſis breviſſimis, tuberculis tern neis. *Ehrh. beyt. tom.* 2. *p.* 106. *p.* 291.

Lichen madreporæformis. *Wulfen lect. tom.* 3. *p.* 105. *t.* 3. *f.* 2.

Coralloides minimum fragile, mad naſcens. *Dill. muſc.* 107. *t.* 16. *f.*

In Carnioliæ, Carinthiæ ſylvi *Wulfen.*

Rr 2

rangiferinus. 89. L. fruticuloſus perforatus ramoſiſſimus, ramulis nutantibus, *Sp. pl. p.* 1620. *Fl. dan. t.* 180. 539.

Coralloides montanum, fruticuli ſpecie ubique candicans. *Dill. muſc.* 107. *t.* 16. *f.* 29.

Coralloides fruticuli ſpecie candicans, corniculis rufeſcentibus. *Dill. muſc.* 110. *t.* 16. *f.* 30.

In montanis, ſubalpinis, alpinis ad terram.

furcatus. 90. L. fruticuloſus tubuloſus ramoſus: ramis erectis furcatis. *Huds. angl. ed.* 1. 458. *Schreb. ſpic. p.* 117.

Lichen caule fruticoſo ſubfolioſo, ramis adſcendentibus repetito - bifurcatis punctiferis. *Hall. hiſt. n.* 1954.

Coralloides corniculis brevioribus & crebrioribus. *Dill. muſc.* 104. *t.* 16. *f.* 27.

Cum præc.

ſpinoſus. 91. L. fruticuloſus tubuloſus ramoſiſſimus ſpinoſus, ramulis tuberculatis ſubdigitatis. *Sibthorp oxon. p.* 334.

Coralloides ſparſum, caulibus tortuoſis & ſpinoſis. *Dill. muſc.* 101. *t.* 16. *f.* 25.

In aſcenſu alpium judenburgenſium ad terram.

ſubulatus. 92. L. fruticuloſus tubuloſus ſubdichotomus, ramis ſimplicibus ſubulatis. *Sp. pl. p.* 1621.

Coralloides corniculis longioribus & rarioribus. *Dill. muſc.* 102. *t.* 16. *f.* 26.

In alpinis ad terram.

vermicularis. 93. L. fruticuloſus lævis ſubramoſus: ramis diffuſis, tuberculis ſparſis. *Syſt. veget. p.* 963.

Lichen tauricus. *Wulfen in Jacqu. collect. tom.* 2. *p.* 177. *t.* 12. *f.* 2.

Cladonia taurica. *Hoffm. lich. t.* 34. *f.* 2.

Cum præc.

ſubuliformis. 94. L. fruticuloſus tubuloſus, filiformi ſimplex, ceſpitoſus, adſcendens, læ mus, ramulis pauciſſimis breviſſimis lis. *Ehrh. beytr. tom.* 3. *p.* 82.

Lichen vermicularis. *Dicks. faſc.* 6. *f.* 10.

Cladonia ſubuliformis. *Hoffm. lic* 1. 2. 3.

In alpibus judenburgenſibus.

paſchalis. 95. L. fruticuloſus ſolidus tectus folioli *Sp. pl. p.* 1621. *Fl. dan. t.* 151.

Lichen caule fruticoſo, fungifero, cis calcareis. *Hall. hiſt. n.* 1963.

Coralloides paſchale. *Hoffm. lich. t*

Coralloides criſpum & botryform *Dill. muſc.* 114. *t.* 17. *f.* 33.

In alpinis, ſubalpinis.

fragilis. 96. L. fruticuloſus ſolidus, ramulis ter obtuſis. *Sp. pl. p.* 1621. *Wulfen in cell. tom.* 2. *p.* 92. *t.* 9. *f.* 6.

Coralloides fragile. *Hoffm. lich. t.*

Coralloides alpinum, corallinæ m *Dill. muſc.* 116. *t.* 17. *f.* 34.

In alpibus Styriæ, Carinthiæ, rupes.

Botrytis. 97. L. fruticuloſus ſubramoſus, tubercu bus ſphæricis flavo - bruneis max *ſcand. p.* 292. *Hagen. lich. t.* 9. *f* *io Iacqu. collect. tom.* 4. *p.* 237. *t*

In Carinthia ad radices abietun

triſtis. 98. L. fruticuloſus ſolidus, ramulis c moſis, apicibus nigricantibus, ſcut libus. *Syſt. veget. p.* 963. *Web.* *t.* 5.

Lichen fruticoſus, durus, niger, chotomus. *Hall. hiſt. n.* 1966.

Lichen rigidus. *Syſt. veget. p.* 959. *Wulfen in Jacqu. miscell. tom.* 2. *p.* 86. *t.* 9. *f.* 6. *Collect. tom.* 2. *p.* 187. *t.* 13. *f.* 5.

Lichen fucoides. *Wulfen in Iacqu. collect. tom.* 3. *p.* 143. *t.* 12. *f.* 3.

Cornicularia triſtis. *Hoffm. lich. t.* 34. *f.* 1.

Coralloides corniculatum fuci tenuioris facie. *Dill. musc.* 118. *t.* 17. *f.* 37.

In rupibus alpium ſtyriacarum, carinthiacarum, tyrolenſium.

✿✿✿✿✿✿✿✿✿✿ *Filamentoſi.*

plicatus. 99. L. filamentoſus pendulus, ramis implexis, ſcutellis radiatis. *Sp. pl. p.* 1622.

Usnea vulgaris, loris longis implexis. *Dill. musc.* 56. *t.* 11. *f.* 1.

In arboribus.

divaricatus. 100. L. filamentoſus pendulus angulatus articulatus tomentoſus, ramis divaricatis, peltis orbiculatis ſeſſilibus. *Syſt. veget. p.* 964.

Usnea mollis, ramis longis compreſſis. *Dill. musc.* 62. *t.* 12. *f.* 5.

In ſubalpinis ad arborum truncos.

jubatus. 101. L. filamentoſus pendulus: axillis compreſſis. *Sp. pl. p.* 1622.

Usnea jubata nigricans. *Dill. musc.* 64. *t.* 12. *f.* 7.

Cum præc.

lanatus. 102. L. filamentoſus ramoſiſſimus decumbens implicatus opacus. *Sp. pl. p.* 1623. *Wulfen in Iacqu. miscell. tom.* 2. *p.* 93. *t.* 10. *f.* 5.

Lichen reticulatus. *Wulfen in Iacqu. collect. tom.* 2. *p.* 187. *t.* 13. *f.* 6.

In alpium ſtyriacarum, tyrolenſium, carinthiacarum rupibus.

pubeſcens. 103. L. filamentoſus ramoſiſſimus decum xus nitidus. *Sp. pl. p.* 1623. *Wulf miscell. tom.* 2. *p.* 87. *t.* 9. *f.* 7.

Usnea cæspitoſa exilis capillacea *musc.* 66. *t.* 13. *f.* 9.

Cum præc.

hirtus. 104. L. filamentoſus ramoſiſſimus erectu farinaceis ſparſis. *Sp. pl. p.* 1623.

Usnea hirta. *Hoffm. lich. t.* 30. *f.*

Usnea vulgatiſſima tenuior & brevi culis. *Dill. musc.* 67. *t.* 13. *f.* 12.

In arboribus, rupibus.

floridus. 105. L. filamentoſus ramoſus erectus, ſc tis. *Sp. pl. p.* 1624.

Usnea florida. *Hoffm. lich. t.* 30.

Usnea vulgatiſſima tenuior & brev biculis. *Dill. musc.* 69. *t.* 13. *f.* 1

Cum præc.

ochroleucus. 106. L. filamentoſus, dichotomo - ram erectiuſculus, inarticulatus inanis ramis divaricatis: apicibus furcatis *beytr. tom.* 3. *p.* 82.

Usnea ochroleuca. *Hoffm. lich. t.*

In alpibus.

vulpinus. 107. L. filamentoſus ramoſiſſimus erect inæquali - anguloſus. *Sp. pl. p.* 1623 *Iacqu. miscell. tom.* 2. *p.* 88. *t.* 1 *dan. t.* 226.

Usnea capillacea citrina, fruticuli *musc.* 73. *t.* 13. *f.* 16.

In Styriæ, Carinthiæ, Tyrolis alpinis ad arborum truncos.

radiciformis. 108. L. filamentoſus teres ramoſiſſimus formis. *Syſt. veget. p.* 294.

Sub cortice arborum.

FUCUS. Fructificatio veſiculæ cavæ.

natans. 1. F. caule filiformi ramoſo, foliis lanceolatis ſerratis, fructificationibus globoſis pedunculatis. *Syſt. veget. p.* 965.

In mare adriatico cum ſequentibus ſpeciebus.

volubilis. 2. F. fronde ſpirali: margine dilatato repando-dentato. *Wulfen in Iacqu. collect. tom.* 3. *p.* 149. *t.* 13. *f.* 2.

canaliculatus. 3. F. fronde plana dichotoma integerrima lineari-canaliculata, fructificationibus tuberculatis bipartitis obtuſis. *Syſt. veget. p.* 967. *Fl. dan. t.* 214.

nodoſus. 4. F. fronde compreſſa dichotoma, foliis integerrimis diſtichis, veſiculis innatis ſolitaris dilatatis. *Syſt. veget. p.* 967. *Fl. dan. t.* 146.

ſiliquoſus. 5. F. ſtirpe filiformi dichotoma, frondibus membranaceis enſiformibus ſolitariis ſerratis: terminalibus petiolo inflatis. *Retzii ſcand. p.* 297. *Sp. pl. p.* 1629. *Fl. dan. t.* 106.

granulatus. 6. F. fronde filiformi ramoſiſſima, ramulis acuminatis, veſiculis ſubrotundis cumulatis, ramis foliisque acutis omnibus innatis. *Syſt. veget. p.* 968. *Fl. dan. t.* 591.

Selaginoides. 7. F. fronde filiformi ramoſiſſima: ramis dichotomis, foliis ſubulatis alternis baſi veſiculoſis. *Syſt. veget. p.* 968. *Gmel. fuc. t.* 2. *A. f.* 1.

hirſutus. 8. F. fronde filiformi tereti dichotoma obtecta undique hirſutie. *Syſt. veget. p.* 969.

Filum. 9. F. fronde filiformi ſubfragili opaca. *Syſt. veget. p.* 969. *Fl. dan. t.* 821.

faſtigiatus. 10. F. fronde filiformi dichotoma ramoſiſſima faſtigiata obtuſa. *Syſt. veget. p.* 969. *Fl. dan. t.* 393.

furcellatus. 11. F. fronde filiformi dichotoma ramoſiſſima acuminata. *Syſt. veget. p.* 969. *Fl. dan. t.* 419.

ciliatus, 12. F. frondibus membranaceis lanceola ciliatis. *Syſt. veget. p.* 970. *Fl. dan*

Fucus caulescens. *Gmel. fuc. t.* 20.

criſpatus. 13. F. frondibus membranaceis ſublinea ſiſſimis crispis coloratis. *Syſt. veget. dan. t.* 826. *Wulfen in Iacqu. col p.* 162. *t.* 16. *f.* 2.

alatus. 14. F. frondibus membranaceis ſubdich tis, laciniis alternis decurrentibus *veget. p.* 970. *Fl. dan. t.* 352. *Gme*

plumoſus, 15. F. frondibus cartilagineis lanceolat plumoſis, caule filiformi compr *Syſt. veget. p.* 971, *Fl. dan. t.* 35

cartilagineus. 16. F. fronde cartilaginea compreſſa fu ſita: laciniis linearibus. *Syſt. veget.*

Fucus verſicolor, *Gmel. fuc. t.* 17.

gigartinus. 17. F. fronde cartilaginea filiformi con toma, fructificationibus globoſis pe minalibus ſubjacente ariſta. *Syſt. v*

ſpinoſus. 18. F. aphyllus cartilagineus ramoſus verticillato-ternis. *Syſt. veget. p. in Iacqu. collect. tom.* 3. *p.* 156.

conſervoides. 19. F. frondibus membranaceis linearib ramoſis, fructificationibus ſparſis ſe tundis. *Syſt. veget. p.* 971. *Wul collect. tom.* 3. *p.* 151. *t.* 14. *f.*

Fucus verrucoſus. *Gmel. fuc. t.* 14

ULVA. Fructificatio in membrana diaph

pavonia. 1. U. plana reniformis ſeſſilis decu *Syſt. veget. p.* 972.

Fucus pavonius. *Sp. pl. p.* 1630.

In mari adriatico cum ſequen

umbilicalis. 2. U. plana orbicularis monophylla peltata coriacea. *Retzii ſcand. p.* 302. *Sp. pl. p.* 1633.

Tremella marina umbilicata. *Dill. muſc.* 45. *t.* 8. *f.* 3.

inteſtinalis. 3. U. tubuloſa ſimplex. *Sp. pl. p.* 1632.

Tremella marina tubuloſa, inteſtinorum figura. *Dill. muſc.* 47. *t.* 9. *f.* 7.

compreſſa. 4. U. tubuloſa ramoſa compreſſa. *Sp. pl. p.* 1632.

Tremella marina tenuiſſima & compreſſa. *Dill. muſc.* 48. *t.* 9. *f.* 8.

latiſſima. 5. U. oblonga plana undulata membranacea viridis. *Sp. pl. p.* 1632.

Lactuca. 6. U. palmata prolifera membranacea, ramentis inferne anguſtatis. *Syſt. veget. p.* 972.

Tremella marina vulgaris, Lactucæ ſimilis. *Dill. muſc.* 42. *t.* 8. *f.* 1.

Linza. 7. U. fronde oblonga bullata. *Sp. pl. p.* 1633. *Fl. dan. t.* 889.

Tremella marina faſciata. *Dill. muſc.* 46. *t.* 9. *f.* 6.

CONFERVA. Tubercula inæqualia in fibris capillaribus longiſſimis.

rivularis. 1. C. filamentis ſimpliciſſimis æqualibus longiſſimis. *Sp. pl. p.* 1633. *Fl. dan. t.* 881.

Conferva fluviatilis ſericea, vulgaris & fluitans. *Dill. muſc.* 12. *t.* 2. *f.* 1.

In aquis.

fontinalis. 2. C. filamentis ſimpliciſſimis æqualibus digito brevioribus. *Sp. pl. p.* 1633. *Fl. dan. t.* 651. *f.* 3.

Conferva minima, Byſſi facie. *Dill. muſc.* 14. *t.* 2. *f.* 3.

In fontibus.

bulloſa. 3. C. filamentis æqualibus ramoſis bullas
cludentibus. *Sp. pl. p.* 1634.

Conferva paluſtris bombycina. *Dill. t.* 3. *f.* 11.

In aquis ſtagnantibus.

canalicularis. 4. C. filamentis æqualibus baſin verſus
bus. *Sp. pl. p.* 1634.

Conferva rivulorum capillacea, denſi
geſtis ramulis. *Dill. muſc.* 21. *t.* 4.

In rivulis, & in aquæductibus.

amphibia. 5. C. filamentis æqualibus ramoſis, exſicc
untibus in aculeos. *Sp. pl. p.* 1634.

Conferva amphibia fibrilloſa & ſpon
muſc. 22. *t.* 4. *f.* 17.

In foſſis.

reticulata. 6. C. filamentis reticulato-coadunatis.
1635. *Dill. muſc.* 20. *t.* 4. *f.* 14.

In aquis ſtagnantibus, fluviis.

fluviatilis. 7. C. filis ſimpliciſſimis ſetiformibus re
culis craſſioribus angulatis. *Sp. pl.*

Conferva fluviatilis lubrica ſetoſa,
cie. *Dill. muſc.* 39. *t.* 7. *f.* 47.

In fluviis.

capillaris. 8. C. filis geniculatis ſimplicibus: artic
tim compreſſis. *Sp. pl. p.* 1636.

Conferva filamentis longis genicula
bus. *Dill. muſc.* 25. *t.* 5. *f.* 25.

In aquis ſtagnantibus, vel lent

corallinoides. 9. C. filis geniculis dichotomis. *Sp. pl.*

Conferva marina gelatinoſa, coralli
niculata craſſior. *Dill. muſc.* 33. *t.*

In mari adriatico.

catenata. 10. C. filis geniculatis: articulis cylindricis, *Sp. pl. p.* 1636.

Conferva ramoſa, geniculis longioribus cateniformibus. *Dill. muſc.* 27. *t.* 5. *f.* 27.

Cum præc.

polymorpha. 11. C. filamentis geniculatis, ramis faſciculatis. *Sp. pl. p.* 1636.

Conferva marina geniculata nigra palmata. *Dill. muſc.* 32. *t.* 6. *f.* 35.

Cum præc.

glomerata. 12. C. filamentis geniculatis, ramulis brevioribus multifidis. *Sp. pl. p.* 1637.

Conferva fontinalis ramoſiſſima, glomeratim congeſta. *Dill. muſc.* 28. *t.* 5. *f.* 31.

In fontibus rivulis.

BYSSUS. Fibræ ſimplices lanuginoſæ.

phosphorea. 1. B. lanuginoſa violacea lignis adnaſcens. *Sp. pl. p.* 1638. *Dill. muſc.* 4. *t.* 1. *f.* 6.

In ligno putreſcente.

velutina. 2. B. capillacea viridis, filamentis ramoſis. *Syſt. veget. p.* 974.

Byſſus tenerrima viridis, velutum referens. *Dill. muſc.* 7. *t.* 1. *f.* 14.

Terræ adnaſcitur.

aurea. 3. B. lanuginoſa crocea, exſiccatione candida, filamentis brevibus ſimplicibus ramoſisque. *Sibthorp. oxon. p.* 328. *Sp. pl. p.* 1638.

Byſſus petræa crocea, glomerulis lanuginoſis. *Dill. muſc.* 8. *t.* 1. *f.* 16.

In rupibus humidis.

ſeptica. 4. B. filamentis longiſſimis tenuiſſimis ra plexis. *Huds. angl. ed.* 2. 607. *Sibth p.* 339.

Byſſus ſubterranea. *Scop. carn. tom.*

Byſſus evanida, floccoſa, nivea. *Dill. t.* 1. *f.* 9.

In locis ſubterraneis lignis put adnaſcitur.

F U N G I.

AGARICUS. Fungus horizontalis, ſubtus l

muſcari-us. 1. A. ſtipitatus, pileo ſanguineo verru rucis lamelliſque albis, ſtipite baſi bul *thorp oxon. p.* 339. *Sp. pl. p.* 1640. *Sc t.* 27. 28. *Curtis lond.*

Agaricus pſeudo - aurantiacus. *Bull t.* 122.

Duodecimi generis pernicioſorum fung ta ſpecies. *Cluſ. hiſt. p. CCLXXX.*

In ſylvis.

cæſareus. 2. A. ſtipitatus, volva maxima, pilei a phærici margine ſtriato, ſtipite craſſ *Retzii ſcand. p.* 311. *Scop. carn. tom.*

Agaricus aurantiacus. *Bulliard herb.*

Decimum ſeptimum genus eſculentor rum. *Cluſ. hiſt. p. CCLXXII.*

In ſylvis Pannoniæ, Croatiæ, Car toralis, Forijulii.

verruco-ſus. 3. A. ſtipitatus, pileo convexo cinereo verrucis lamelliſque albis, ſtipite ba *Huds. angl.* 2. 613. *Sibthorp oxon. p tis lond.*

In ſylvis.

procerus. 4. A. ſtipitatus, pileo campanulato ſubfuſco ſquamato, lamellis albidis, ſtipite longo ſquamato bulboſo annulato. *Sibthorp oxon. p.* 340. *Huds. angl.* 2. 612. *Curtis lond. Schæff. fung. t.* 22. 23. *Scop. carn. tom.* 2. *p.* 418.

Agaricus colubrinus. *Bulliard herb. t.* 78.

In ſylvaticis, campeſtribus.

campeſtris. 5. A. ſtipitatus, pileo convexo ſquamato albido, lamellis rufis. *Sp. pl. p.* 1641. *Schæff. fung. t.* 33.

Agaricus edulis. *Bulliard herb. t.* 134.

In campis, pratis, hortis.

aureus. 6. A. ſtipitatus fulvus, pileo leviter floccoſo, ſtipite lævi curvato annulato: annulo fugaci. *Sibthorp oxon. p.* 343. *Bulliard herb. t.* 92.

In ſylvis.

floccoſus. 7. A. ſtipitatus, faſciculoſus, pileo ſtipiteque piloſo-ſquamoſis flavo-fuſcis. *Curtis lond. Sibthorp oxon. p.* 343. *Schæff. fung. t.* 61.

Agaricus ſquamoſus. *Bulliard herb. t.* 266.

In arborum truncis.

eburneus. 8. A. ſtipitatus, niveus, lamellis bifidis decurrentibus, ſtipite brevi farcto. *Sibthorp oxon. p.* 344. *Bulliard herb. t.* 118.

In paſcuis.

integer. 9. A. ſtipitatus, pileo convexiuſculo viſcido purpureo, lamellis omnibus æqualibus albis. *Huds. angl.* 2. 610. *Sp. pl. p.* 1640. *Bolt. fung. t.* 1. *Schæff. fung. t.* 204. 214.

In ſylvis.

ſpadiceus. 10. A. ſtipitatus, pileo ſpadiceo pilis araneoſis marginato, lamellis concoloribus, ſtipite albido brevi baſi incraſſato. *Sibthorp oxon. p.* 346.

Agaricus araneoſus. *Bulliard herb.*

In ſylvis.

velatus. 11. A. ſtipitatus, pileo cervino glutino ſubalbidis, ſtipite craſſo ſubcylindri *oxon. p.* 347. *Schæff. fung. t.* 36.

In ſylvis.

terreus. 12. A. ſtipitatus, pileo convexo fragili nigro, lamellis albidis, ſtipite alb *Sibthorp oxon. p.* 347. *Schæff. fun*

In ſylvis.

cruſtuliniformis. 13. A. ſtipitatus, pileo ex albido rufes lis rufis, ſtipite furfuraceo albido. *S p.* 348. *Bulliard herb. t.* 308.

In pascuis, montanis ſylvaticis

craſſipes. 14. A. ſtipitatus, fuscus, pileo mutabil ciſo, lamellis fuscescentibus, ſtipit ſubulato, ſurſum craſſiori. *Dicks.* *Sibthorp oxon. p.* 350. *Schæff. fu*

In ſylvaticis.

lactifluus. 15. A. ſtipitatus, pileo plano carne lamellis rufis, ſtipite longo carne 1641. *Schæff. fung. t.* 5.

In ſylvis.

dulcis. 16. A. ſtipitatus, pileo convexo carn lamellis ſtipiteque concoloribus. *S p.* 351. *Bulliard herb. t.* 224.

In ſylvaticis.

delicioſus. 17. A. ſtipitatus, pileo teſtaceo, ſuc *Sp. pl. p.* 1641. *Schæff. fung. t.*

In ſylvis.

Liſteri. 18. A. ſtipitatus, pileo planiuſculo la gine deflexo, lamellis ramoſis pall *oxon. p.* 352.

Agaricus lactifluus acris. *Bulliard herb.* t. 200.

In paſcuis, ſylvis.

ſangui-neus. 19. A. ſtipitatus, ſanguineo-ruber unicolor, lamellis araneoſo fugaci aurantiaco. *Syſt. veget.* p. 975. *Wulfen in Iacqu. miscell.* tom. 2. p. 107. t. 15. f. 3.

In ſylvis.

viscidus. 20. A. ſtipitatus, pileo purpuraſcente-fusco viscido, lamellis fusco-purpuraſcentibus. *Sp. pl.* p. 1642.

In ſylvis.

virgine-us. 21. A. ſtipitatus niveus, pileo hemiſphærico, lamellis arcuatim per ſtipitem decurrentibus. *Syſt. veget.* p. 975. *Wulfen in Iacqu. miscell.* tom. 2. p. 104. t. 15. f. 1.

ceraceus. 22. A. ſtipitatus, pileo hemiſphærico ſtipiteque ſubfiſtuloſo flavis, lamellis aquoſe luteolis. *Syſt. veget.* p. 975. *Wulfen in Iacqu. miscell.* tom. 2. p. 105. t. 15. f. 2.

In paſcuis, ad margines ſylvarum.

ericeus. 23. A. ſtipitatus, pileo albo viscido margine tenui; lamellis decurrentibus ſtipiteque concoloribus. *Sibthorp oxon.* p. 353. *Bulliard. herb.* t. 188.

In paſcuis, ſylvaticis.

dentatus. 24. A. ſtipitatus, pileo convexo, lamellis baſi mucrone dentatis. *Sp. pl.* p. 1640. *Schæff. fung.* t. 302.

In paſcuis, montanis ſylvaticis.

pſittaci-nus. 25. A. pileo conico viscido viridi-flavo, lamellis luteis, ſtipite viridi. *Sibthorp oxon.* p. 356. *Schæff. fung.* t. 301.

In paſcuis.

velutipes. 26. A. ſtipitatus faſciculoſus, pileo plar vo, ſtipite nudo tenerrime villoſo *Curtis lond.*

Ad arborum truncos, ſtipites ram

tener. 27. A. pileo conico fulvo margine ſatu mellis atro-rufis infra marginem ext te concolori. *Sibthorp oxon. p.* 3(*fung.* t. 70.

In ſylvis, pratis.

ramealis. 28. A. ſtipitatus, pileo convexo candid non decurrentibus ſtipiteque concolo *thorp oxon. p.* 360. *Bulliard herb. t.*

In ramis arborum putreſcentibus.

clavus. 29. A. ſtipitatus, pileo luteo convexo, piteque albis. *Sp. pl. p.* 1644. *Bu t.* 148.

Inter muſcos, folia, & in ligno

alliaceus. 30. A. ſtipitatus, pileo convexo opaco co, ſubtus albo, ſtipite fiſtuloſo n *Syſt. veget. p.* 976. *Iacqu. auſtr. t.*

In ſylvis ſubalpinis.

eſculentus. 31. A. ſtipitatus, pileo convexo argillace albis, ſtipite gracili fiſtuloſo ex a flaveſcente. *Syſt. veget. p.* 976. *Wulf miſcell. tom.* 2. *p.* 103. *t.* 14. *f.* 4.

In ſylvaticis.

ebúrneus. 32. A. ſtipitatus, pileo campanulato ac livido nitido, lamellis albis, ſtipite nereo. *Huds. angl.* 2. 620. *Sibtho* 364. *Schæff. fung. t.* 52.

Ad truncos, radices arborum.

flavipes. 33. A. ſtipitatus, pileo campanulato pli fuſco, lamellis albidis, ſtipite fiſt *Sibthorp oxon. p.* 305. *Schæff. fung.*

In ſylvis.

minutulus. 34. A. ſtipitatus, pileo tenui ſtriato ſubfuſco, lamellis raris flaveſcentibus, ſtipite albido nudo. *Dicks. faſc.* 1. *p.* 16. *Schæff. fung. t.* 308.

In herbidis inſularum danubialium.

campanulatus. 35. A. ſtipitatus, pileo campanulato ſtriato pellucido, lamellis adſcendentibus, ſtipite nudo. *Sp. pl. p.* 1645. *Schæff. fung. t.* 211.

In ſylvaticis, paſcuis.

androſaceus. 36. A. ſtipitatus albus, pileo plicato membranaceo, ſtipite nigro. *Sp. pl. p.* 1644. *Bolt. fung. t.* 32. *Bulliard herb. t.* 64.

Ad arborum ramos deciduos.

fimiputris. 37. A. ſtipitatus, pileo pallide fuſco obtuſe conico, lamellis ſubnigris, ſtipite albido verſus apicem irregulariter annulato. *Sibthorp oxon. p.* 368. *Bulliard herb. t.* 66.

In paſcuis, hortis.

plicatilis. 38. A. ſtipitatus, pileo campanulato ſeu plano murino pellucido plicato. *Curtis lond. Sibthorp oxon. p.* 368.

In pratis, paſcuis.

ſtercorarius. 39. A. ſtipitatus, pileo ſubconico albo furfuraceo, lamellis cinereo-nigris, ſtipite albo fiſtuloſo. *Sibthorp oxon. p.* 369. *Bulliard herb. t.* 68.

In paſcuis, ad fimeta.

cylindricus. 40. A. ſtipitatus, pileo albo cylindrico flocceſo, lamellis purpuraſcenti-nigris uniformibus, ſtipite cylindrico albo. *Sibthorp. oxon. p.* 370.

Agaricus fimetarius. *Curtis lond.*

In pratis, paſcuis.

ovatus. 41. A. ſtipitatus, pileo ovato ſubplicato, ſtipite nudo ad baſin attenuato ſcabriuſculo: lamellis creberrimis ſubcoaleſcentibus. *Curtis lond. Sibthorp oxon. p.* 370.

In paſcuis, pratis, hortis.

appendiculatus. 42. A. ſtipitatus, pileo obtuſe conico ſtr dehiscente marginato, lamellis crebr fis, ſtipite fiſtuloſo albo. *Sibthorp ox Bulliard herb. t.* 317.

In pratis, hortis, pomariis.

micaceus. 43. A. ſtipitatus, pileo conico flavesce centro micaceo, lamellis creberrimis pite fiſtuloſo albo. *Sibthorp oxon. p. liard herb. t.* 246.

In cultis, ſylvaticis.

turbinatus. 44. A. ſtipitatus, pileo flavescente fusco baſi inæquali, lamellis ochroleucis b pite albo. *Sibthorp oxon. p.* 371. *Sc t.* 66.

Ad arborum truncos.

aquoſus. 45. A. ſtipitatus, pileo campanulato obt flavescente, lamellis atro-griſeis, fl loſo griſeo villoſo. *Huds. angl.* 2. 69 *fung. t.* 6. *Sibthorp oxon. p.* 371.

In pascuis, ſylvis, ad arborum t

niveus. 46. A. ſubſtipitatus totus niveus pileo din bro, ſtipite brevi ad baſin hirſuto. *p.* 976. *Iacqu. auſtr. t.* 288.

Agaricus *reſupinatus*, acaulis rever lamellis concoloribus, margine inflex *oxon. p.* 373.

In ſylvis ſubalpinis ad ligna put tum folia.

oſtreatus. 47. A. ſubſtipitatus fasciculato-imbricat nereo obovato margine involuto, la dis decurrentibus baſi ſubanaſtomiſan *tis lond. Syſt. veget. p.* 976. *Iacqu.* c

Ad arborum truncos.

quercinus. 48. A. acaulis, lamellis labyrinthiformi *p.* 1644. *Bolt. fung. t.* 73.

Ad truncos quercuum.

betulinus. 49. A. acaulis coriaceus villoſus, margine obtuſo, lamellis ramoſis anaſtomiſantibus. *Sp. pl. p.* 1645. *Bulliard herb. t.* 394. *Fl. dan. t.* 776. *f.* 1.

Ad arborum truncos.

alneus. 50. A. acaulis, lamellis bifidis pulverulentis. *Sp. pl. p.* 1645. *Schæff. fung. t.* 246.

Ad ligna putrida, arborum truncos.

ochraceus. 51. A. acaulis flabelliformis, ſupra pulvinatus ochraceus, lamellis albidis, margine cineraſcentibus. *Syſt. veget. p.* 977. *Wulfen in Iacqu. miſcell. tom.* 2. *p.* 110. *t.* 16. *f.* 2.

In Carinthia ad arborum emortuarum truncos. *Wulfen.*

MERULIUS. Fungus ſubtus venoſus.

cantharellus. 1. M. ſtipitatus, pileo irregulariter infundibuliformi aurantio, venis ramoſis anaſtomiſantibus, ſtipite ſubcompreſſo, concoloribus. *Sibthorp oxon. p.* 374.

Agaricus cantharellus. *Sp. pl. p.* 1639. *Bulliard herb. t.* 62.

In ſylvis.

umbonatus. 2. M. pileo umbonato cinereo, ſtipite concolori tereti, venis albis. *Syſt. nat. ed.* 13. *p.* 1430.

Agaricus muscoides. *Wulfen in Iacqu. miſcell. tom.* 2. *p.* 109. *t.* 16. *f.* 1.

In ſylvis, ad ſepes.

cornucopioides. 3. M. fuscus tenax, pileo lobato, venis ſubquadriplici ordine decurrentibus. *Syſt. nat. ed.* 13. *p.* 1429. *Schæff. fung. t.* 275.

Peziza cornucopioides. *Sp. pl. p.* 1650.

In ſylvis.

BOLETUS. Fungus horizontalis, ſubtus po

bovinus. 1. B. ſtipitatus, pileo olivaceo-fusco, albo fuscescentibus inæqualibus, ſtip albo. *Sibthorp oxon. p.* 375. *Sp. pl.*

Boletus edulis. *Bulliard herb. t.* 60.

In ſylvis.

luteus. 2. B. ſtipitatus, pileo ſubviscido, poris flaviſſimis, ſtipite albido, annulo *Sibthorp oxon. p.* 377. *Sp. pl. p.* 16 *fung. t.* 114.

In ſylvis.

olivaceus. 3. B. ſtipitatus, pileo olivaceo deflexo tabili, poris minutis flavis. *Sibtho* 376. *Schæff. fund. t.* 105.

Boletus tuberoſus. *Bulliard. herb. t.*

In ſylvaticis.

cinnamomeus. 4. B. ſtipitatus, pileo cinnamomeo, ſ nitido, poris angulatis, ſtipite tere loſo. *Iacqu. collect. tom.* 1. *p.* 116

In ſylvis montanis agri vindob

leptocephalus. 5. B. ſtipitatus, pileo plano tenui ſ ſubtus albo. *Syſt. veget. p.* 977. *I tom.* 1. *p.* 142. *t.* 12.

In ſylvis in putridis arborum

ramoſiſſimus. 6. B. ſtipitatus ramoſiſſimus pileis m ticeps. *Syſt. veget. p.* 978. *Iacqu. Schæff. fung. t.* 111. 265. 266. *Sc* 2. *p.* 470.

In quercubus, aliisque arbori

flabelliformis. 7. B. ſubſtipitatus, pileo coriaceo mu gine lobato plicato lacerove, ſup poris albidis. *Schæff. fung. t.* 113

In ſylvis Auſtriæ montanis.

ſquamoſus. 8. B. ſubſtipitatus planiuſculus lacero - ſquamoſus flaveſcens, poris tetragonis. *Huds. angl.* 2. 626. *Sibthorp oxon. p.* 378.

Boletus juglandis. *Schæff. fung. t.* 101. 102. *Bulliard. herb. t.* 19.

Ad arborum truncos.

rugoſus. 9. B. ſubſtipitatus perennis caſtaneus nitidus, pileo dimidiato rugoſo, poris albis. *Syſt. veget. p.* 977. *Iacqu. auſtr. t.* 169.

Boletus laccatus. *Timm. meg. p.* 269.

Boletus nitens. *Batſch fung. f.* 225. *Schæff. t.* 109. 110. &c.

Agaricus pſeudo - boletus. *Iacqu. auſtr. t.* 41.

Ad ſtipites truncatarum arborum, ramos deciduos.

verſicolor. 10. B. acaulis: faſciis diſcoloribus, poris albis. *Sp. pl. p.* 1645. *Bulliard. herb. t.* 86. *Schæff. fung. t.* 136.

Ad arborum, fruticum truncos, ſtipites.

ſuberoſus. 11. B. acaulis pulvinatus albus lævis, poris acutis difformibus. *Sp. pl. p.* 1645. *Schæff. fung. t.* 136. 263. *Bulliard herb. t.* 162.

Ad arborum truncos.

igniarius. 12. B. acaulis pulvinatus lævis, poris tenuiſſimis. *Sp. pl. p.* 1645. *Bulliard herb. t.* 82. *Schæff. fung. t.* 106.

Ad arborum truncos.

ſuaveolens. 13. B. acaulis, ſuperne lævis, ſalicinus. *Sp. pl. p.* 1646. *Bulliard herb. t.* 310. *Schæff. fung. t.* 124.

Ad ſalicum truncos.

officinalis. 14. B. acaulis pulvinatus, ſuperne eminentiis convexis diſcoloribus, ſubtus poris ochroleucis. *Batſch fung. p.* 111. *Iacqu. miſcell. tom.* 1. *t.* 20.

Polyporus ſeſſilis, convexo planus, coloribus fulvis, poris ochroleucis *n.* 2284.

Boletus Agaricum. *Allion. pedem.* *t*

Boletus Laricis. *Bulliard herb.* *t.* 29

Ad Laricis truncos.

cinnabarinus. 15. B. acaulis pulvinatus, totus intus & *Syſt. veget. p.* 977. *Iacqu. auſtr. t.*

In ſylvis ſubalpinis ad arborum

Medulla panis. 16. B. cruſtaceus albus effuſus difformis, *p.* 977. *Iacqu. miscell. tom.* 1. *p.* 14

Polyporus effuſus cruſtaceus albus, *n.* 2272.

Agaricum terreſtre, medullam pa *Mich. gen. p.* 121. *t.* 63. *f.* 2.

Ligno putrido & ipſi nudæ hum

HYDNUM. Fungus horizontalis, ſubtus

imbricatum. 1. H. ſtipitatum, pileo convexo imbri *p.* 1647.

Erinaceus esculentus albus craſſus. 132. *t.* 72. *f.* 2.

In ſylvis montanis, ſubalpinis.

repandum. 2. H. ſtipitatum, pileo convexo lævi *pl. p.* 1647. *Bulliard herb. t.* 172. *t.* 316.

Cum præc.

auriscalpium. 3. H. ſtipitatum, pileo dimidiato. *Sp.* *Curtis lond.*

Erinaceus parvus hirſutus ex fusco ſemiorbiculari, pediculo tenuiore. 132. *t.* 72. *f.* 8.

In ſylvis innascitur conis ra dis Pini.

gelatino-ſum. 4. H. ſtipitatum gelatinoſum, pileo dimidiato ſupra glaberrimo. *Syſt. veget. p.* 978. *Iacqu. auſtr. t.* 239.

Echinus cryſtallinus, gelatinoſus. *Hall. hiſt.* 2. 2319. *Scop. carn. tom.* 2. *p.* 472.

In ſylvis ſubalpinis innaſcitur ſemiputridis arborum ramis.

PHALLUS. Fungus ſubtus lævis, ſupra reticulatus.

eſculentus. 1. P. pileo ovato celluloſo, ſtipite nudo rugoſo. *Sp. pl. p.* 1648. *Bulliard herb. t.* 218.

Boletus eſculentus rugoſus fulvus. *Tourn. inſt.* 561. *Batt. fung. p.* 24. *t.* 2. *f. F.*

In ſylvis.

impudicus. 2. P. volvatus ſtipitatus, pileo celluloſo. *Sp. pl. p.* 1648. *Curtis lond. Bulliard herb. t.* 182.

Boletus Phalloides. *Tourn. inſt.* 562. *Batt. fung. p.* 23. *t.* 2. *f. A.*

In ſylvis.

CLATHRUS. Fungus ſubrotundus, cancellatus.

cancellatus. 1. C. acaulis ſubrotundus. *Sp. pl. p.* 1648.

Boletus cancellatus purpureus. *Tourn. inſt.* 561. *Batt. fung. p.* 23. *t.* 2. *f. E.*

In ſylvis Litoralis, Forijulii.

HELVELLA. Fungus turbinatus, ſuperne & inferne lævis.

Mitra. 1. H. pileo deflexo adnato lobato difformi. *Sp. pl. p.* 1649. *Bulliard herb. t.* 190.

In ſylvis montanis, ſubalpinis.

PEZIZA. Fungus concavus ex parte ſu[illegible]
na emittens.

punctata. 1. P. turbinata truncata disco puncta
1650. *Bulliard herb. t.* 252.

In ſtercore vaccino & equino.

epidendra. 2. P. ſubſtipitata campanulata, fragilis
intus coccinea. *Sibthorp oxon. p.* [illegible]

Peziza coccinea. *Bolt. fung. t.* 107

In umbroſis, humidis ad ran
ſtipites.

citrina. 3. P. ſubſtipitata citrina gregaria, cra
planiuscula. *Sibthorp oxon. p.* 386

Octoſpora citrina. *Hedw. musc.* 2.

Ad arborum truncos emortuos

coronaria. 4. P. cyathiformis utrinque glabra, l
erecto. *Syſt. veget. p.* 980 *Iacqu.*
1. *p.* 140. *t.* 10.

In ſylvis umbroſis.

ſcutellata. 5. P. plana: margine convexo pilo
1651.

Octoſpora ſcutellata. *Hedw. musc.*

In umbroſis, humidis ad terra

hirta. 6. P. hemisphaerica: extrinſecus piloſ
tus coccinea glabra. *Syſt. nat. ed.*

Octoſpora hirta. *Hedw. musc.* 2.

In ligno putrido.

leucoloma. 7. P. minuta depreſſa ruberrima: ma
briato. *Syſt. nat. ed.* 13. *p.* 1458.

Octoſpora leucoloma. *Hedw. mus*

In humidis, praecipue inter
innascitur.

coccinea. 8. P. multiformis : extus ex flavo pallens ; intus cinnabarina, margine aurantiaco crenulata. *Iacqu. auſtr. t.* 163. *Schæff. fung. t.* 148.

Peziza cava, albida, intus coccinea. *Hall. hiſt. n.* 2222.

In umbroſis humidis innaſcitur putridis ramulis per ſuperficiem terræ dejectis.

Auricula. 9. P. acaulis auriformis rugoſa membranacea fuſca. *Sibthorp oxon. p.* 387.

Tremella Auricula. *Sp. pl. p.* 1625.

Primum genus perniciosorum fungorum. *Cluſ. hiſt. p CCLXXVI.*

Ad arborum truncos.

cochlea- ta. 10. P. turbinata cochleata. *Sp. pl. p.* 1651. *Bulliard herb. t.* 154.

In hortorum umbroſis ad fimeta, in caldariis.

TREMELLA. Fungus gelatinoſus varie expanſus ex tota ſuperficie ſemina emittens.

juniperi- na. 1. T. ſeſſilis membranacea auriformis fulva. *Syſt. veget. p.* 965.

In junipero communi, Oxycedro.

Noſtoc. 2. T. plicata undulata fuſco-virescens. *Huds. angl.* 2. 564. *Sibthorp oxon. p.* 390.

Tremella terreſtris ſinuoſa, pinguis & fugax. *Dill. muſc.* 52. *t.* 10. *f.* 14.

In paſcuis, campis, alibique.

arborea. 3. T. ſeſſilis ſubrotunda undulata nigrescens. *Huds. angl.* 2. 563. *Sibthorp oxon. p.* 390.

Tremella arborea nigricans, minus pinguis & fugax. *Dill. muſc.* 54. *t.* 10. *f.* 15.

Ad arborum truncos.

criſpa. 4. T. terreſtris atro-virens tenera criſp: *oxon. p.* 391. *Dill. muſc.* 52. *t.* 10.

In umbroſis humidis.

granu-lata. 5. T. gelatinoſa ſphærica aggregata vir *angl.* 2. 566.

Ulva granulata. *Sp. pl. p.* 1633.

Tremella paluſtris, veſiculis ſphæric mibus. *Dill. muſc.* 55. *t.* 10. *f.* 17.

Ad foſſas, lacus, fluvios.

meſente-rica. 6. T. membranacea gelatinoſa congeſta rea. *Dicks. faſc.* 1. *p.* 14. *Sibthorp o: Syſt. veget. p.* 965. *Iacqu. miſcell.* 142. *t.* 13.

Ad arborum ramos emortuos, ſti

rufa. 7. T. longitudinaliter dimidiato-infund *Syſt. veget. p.* 965. *Iacqu. miſcell. ton t.* 14.

In ligno putrido.

hydnoi-dea. 8. T. compreſſo-plana difformiter digi *Syſt. veget. p.* 965. *Iacqu. miſcell.* 145. *t.* 16.

In ligno putrido.

purpurea. 9. T. ſubgloboſa ſeſſilis ſolitaria glab *Sp. pl. p.* 1626. *Bulliard herb. t.* 2

Lichenoides tuberculoſum amœne pur *muſc.* 127. *t.* 18. *f.* 6.

Ad arborum truncos, ramos mortuos.

NIDULARIA. Fungus coriaceus campanulatus in fundo capſulas lentiformes fovens.

campanulata. 1. N. campanulata livido-plumbea lævis: margine patente, capſulis glabris. *Sibthorp oxon. p.* 393.

Peziza lentifera. *Sp. pl. p.* 1649.

In campeſtribus, ad vias.

ſtriata. 2. N. conica fusca: externe villoſa; interne ſtriata, capſulis ſubtus tomentoſis. *Sibthorp oxon. p.* 393. *Bulliard herb. t.* 40. *f. A.*

Ad arborum ſtipites, lignum putridum.

lævis. 3. N. crucibuliformis ſordide lutea: intus lævis, margine erecto, capſulis glabris. *Sibthorp oxon. p.* 393. *Bulliard herb. t.* 40. *f. B. C.*

In ligno putrido.

CLAVARIA. Fungus lævis, oblongus, ſuperficie unica.

Ophiogloſſoides. 1. C. clavata integerrima compreſſa obtuſa. *Sp. pl. p.* 1652.

In ſylvis.

cæſpitoſa. 2. C. cæſpitoſa, caulibus ſimpliciſſimis, baſi nonnunquam unitis oblongo clavatis luteolis. *Syſt. veget. p.* 980. *Wulfen in Iacqu. miscell. tom.* 2. *p.* 98. *t.* 12. *f.* 2.

In ſylvis montanis, ſubalpinis.

alveloides. 3. C. cæſpitoſa, caulibus ſimpliciſſimis craſſiſſimis, baſi unitis, obverſe pyramidatis, ſtriatis. *Syſt. veget. p.* 980. *Wulfen in Iacqu. miscell. tom.* 2. *p.* 99. *t.* 12. *f.* 3. *Schæff. fung. t.* 164. *f.* 1.

In ſylvaticis.

digitata. 4. C. ramoſa lignea nigra. *Sp. pl. p.* 1652.

In ſylvis.

Hypoxylon. 5. C. ramoſo - cornuta compreſſa. *Sp.* |
In umbroſis ad lignum.

cornuta. 6. C. ramoſa, ramis ſimpliciſſimis, con no - truncatis. *Syſt. veget. p.* 980. *Iacqu. miscell. tom.* 2. *p.* 98. *t.* 14.
In ſylvis.

Corralloides. 7. C. ramis confertis ramoſiſſimis inæqu *pl. p.* 1652. *Bulliard herb. t.* 222.
In ſylvis.

faſtigiata. 8. C. ramis confertis ramoſiſſimis faſtigi luteis. *Sp. pl. p.* 1652. *Bolt. fung.*
In pascuis ſterilibus, montanis ſ

crispa. 9. C. foliacea ramoſiſſima, ramis pl per oras ſerratis. *Syſt. veget. p.* 98 *in Iacqu. miscell. tom.* 2. *p.* 110. *t.*
In ſylvis.

plebeja. 10. C. ramoſiſſima, ramis ſimpliciusculi apice denticulatis. *Syſt. veget. p.* 9 *in Iacqu. miscell. tom.* 2. *p.* 101. *t.*
In ſylvis.

muscoides. 11. C. ramis ramoſis acuminatis inæqua *Sp. pl. p.* 1652. *Fl. dan. t.* 775. *f. fung. t.* 173.
In ſylvis inter muscos.

TUBER. Fungus ſubterraneus carnoſus ſ ſemina fovens.

cibarium. 1. T. globoſum ſolidum verrucoſum tutum. *Sibthorp oxon. p.* 398.
Lycoperdon Tuber. *Sp. pl. p.* 16
In pascuis, montanis ſylvatic
In Auſtria, Moravia occurrit tice albo.

LYCOPERDON. Fungus ſubrotundus ſemina farinacea emittens.

* *Terreſtria.*

ſtellatum. 1. L. volva multifida patente, capitulo glabro breviter pedunculato, ore acuminato. *Sibthorp oxon p.* 398. *Sp. pl. p.* 1653. *Bolt. fung. t.* 179.

In ſylvis ſubalpinis.

Carpobolus. 2. L. volva multifida, fructu globoſo ex ſeminibus combinatis. *Sp. pl. p.* 1654. *Fl. dan. t.* 895.

Cum praec.

pedunculatum. 3. L. ſtipite longo, capitulo globoſo glabro: ore cylindrico integerrimo. *Sp. pl. p.* 1654. *Bulliard herb. t.* 294.

In campis, pratis, alibique.

verrucoſum. 4. L. ſubglobofum rufo-fuſceſcens, carne cæſio-rubeſcente, ſeminibus fuſcis, collo radicis plicato. *Sibthorp oxon. p.* 399. *Bulliard herb. t.* 24.

In ſylvaticis.

Boviſta. 5. L. ſubrotundum: lacerato-dehiſcens. *Sp. pl. p.* 1653.

In apricis herbidis.

** *Paraſitica.*

piſiforme. 6. L. globoſum ſcabrum: ore perforato. *Syſt. veget. p.* 982. *Iacqu. miſcell. tom.* 1. *p.* 137. *t.* 7.

In ligno putrido.

ramoſum. 7. L. pyriforme ſcabrum, radice filiformi ramoſiſſima prolifera. *Syſt. veget. p.* 982. *Iacqu. auſtr. t.* 224.

In ſubalpinis ligno putrido innaſcitur.

Epidendrum. 8. L. cortice farinaque purpurea. *Sp.* *Fl. dan. t.* 720. *Schæff. fung. t.* 1

In muſcoſis, & ligno putrido.

luteum. 9. L. ſubrotundum gregarium, glabru rato fuſco. *Syſt. veget. p.* 982. *I* *tom.* 1. *p.* 138. *t.* 8.

In ſylvis inter delapſa folia.

ÆCIDIUM. Capſula utrinque glabra, ſe dis non cohærentibus plena.

fuſcum. 1. Æ; ſeminibus fuſcis. *Syſt. nat. ed.*

In plantarum foliis.

pini. 2. Æ. oblongo - compreſſum pallidu aurantiis. *Syſt. nat. ed.* 13. *p.* 14

In foliis pini.

epiphyllum. 3. Æ. aggregatum, ore multifido l re fulvo.

Lycoperdon epiphyllum. *Sp. pl.*

Lycoperdon poculiforme. *Iacqu.* *p.* 122. *t.* 4. *f.* 1.

In dorſo foliorum Berberis, E rumque plantarum.

cancellatum. 4. Æ. cancellatum filis parallelis.

Lycoperdon cancellatum. *Syſt.* *Iacqu. auſtr. t.* 17. *Fl. dan. t.* 7

In averſa pyri communis fol

SPHÆRIA. Fungus variæ figuræ. Fruct rica, maſſa pulverulenta abs repleta.

maxima. 1. S. magna craſſa atra, ſuperne *Sibthorp oxon. p.* 402. *Dicks.* *Bolt. fung. t.* 181.

Ad lignum putreſcens.

fragifor-mis. 2. S. convexa obſcure rubra. *Hoffm. veg. crypt.* 1. *p.* 20. *t.* 5. *f.* 1.

Arborum cortici innaſcitur.

MUCOR. Fungus veſiculis ſtipitatis, in quibus ſemina numeroſa.

Mucedo. 1. M. ſtipitatus, capſula globoſa. *Sp. pl. p.* 1655.

In corporibus putridis.

miniatus. 2. M. ſtipitatus, capſula pyriformi. *Syſt. veget. p.* 982. *Jacqu. auſtr. t.* 299.

In ligno putrido, ſtercore cervino.

glaucus. 3. M. ſtipitatus, capitulo ſubrotundo aggregato. *Sp. pl. p.* 1656.

In Melonibus, pomis, aliisque fructibus corruptis.

cruſtaceus. 4. M. ſtipitatus, ſpicis digitatis. *Sp. pl. p.* 1656.

In cibis corruptis.

Eryſiphe. 5. M. albus, capitulis fuſcis ſeſſilibus. *Sp. pl. p.* 1656.

In foliis putridis.

ſepticus. 6. M. unctuoſus flavus. *Sp. pl. p.* 1656.

In ligno putrido.

INDEX GENERUM.

Acanthus. p. 350.
Acer. 546.
Achillea. 472.
Aconitum. 296.
Acorus. 196.
Acroſtichum. 553.
Actæa. 289.
Adiantum. 560.
Adonis. 307.
Adoxa. 218.
Aecidium. 655.
Aegilops. 543.
Aegopodium. 172.
Aesculus. 207.
Aethuſa. 160.
Agaricus. 637.
Agrimonia. 256.
Agroſtemma. 251.
Agroſtis. 40.
Aira. 42.
Ajuga. 316.
Alchemilla. 81.
Aliſma. 206.
Allium. 183.
Alopecurus. 34.
Alſine. 175.
Althæa. 383.
Alyſſum. 362.
Amaranthus. 516
Amygdalus. 271.
Anagallis. 102.
Anaſtatica. 354.
Anchuſa. 88.
Andromeda. 222
Andropogon. 54
Androſace. 94.
Anemone. 299.
Anethum. 69.
Angelica. 156.
Anthemis. 471.
Anthericum. 191
Anthoceros. 610
Anthoxanthum.
Anthyllis. 393.
Antirrhinum. 34
Apargia. 423.
Apium. 171.
Aquilegia. 298.

Arabis. 374.
Arbutus. 224.
Arctium. 438.
Arenaria. 244.
Aretia. 94.
Ariſtolochia. 495.
Arnica. 467.
Artemiſia. 449.
Arum. 497.
Arundo. 61.
Aſarum. 255.
Asclepias. 126.
Asparagus. 193.
Asperugo. 92.
Asperula. 71.
Asphodelus. 191.
Asplenium. 555.
Asprella. 35.
Aſter. 461.
Aſtragalus. 408.
Aſtrantia. 138.
Athamanta, 147.
Atragene. 303.
Atriplex. 545.
Atropa. 117.
Avena. 58.
Azalea. 103.
Ballota. 330.
Bartramia. 586.
Bartſia. 338.
Bellis. 469.
Berberis. 201.
Beta. 129.
Betonica. 327.
Betula. 513.
Bidens. 446.
Biscutella. 364.
Blaſia. 609.
Blitum. 2.
Boletus. 645.
Borago. 92.
Braſſica. 377.
Briza. 48.
Bromus. 54.
Bryonia. 525.
Bryum. 579.
Bunias. 352.
Buphthalmum. 474.
Bupleurum. 139.
Butomus. 219.
Buxbaumia. 600.
Buxus. 514.
Byſſus. 636.
Cacalia. 447.
Cactus. 270.
Calendula. 478.
Calla. 498.
Callitriche. 2.
Caltha. 315.
Campanula. 105.

Camphorosma. 81.
Cannabis. 533.
Capparis. 289.
Cardamine. 366.
Carduus. 440.
Carex. 502.
Carlina. 445.
Carpeſium. 454.
Carpinus. 522.
Carum. 169.
Caucalis. 143.
Celtis. 547.
Cenchrus. 543.
Centaurea. 475.
Ceraſtium. 252.
Ceratophyllum. 517.
Cerinthe. 91.
Chærophyllum. 164.
Chaiturus. 331.
Chara. 499.
Cheiranthus. 373.
Chelidonium. 289.
Chenopodium. 127.
Cherleria. 247.
Chlora. 210.
Chondrilla. 426.
Chryſanthemum. 469.
Chryſocoma. 448.
Chryſoſplenium. 225.
Cicer. 401.
Cichorium. 436.
Cicuta. 160.
Cineraria. 463.
Circæa. 4.
Ciſtus. 292.
Clathrus. 648.
Clavaria. 652.
Clematis. 304.
Clinopodium. 332.
Clypeola. 363.
Cnicus. 443.
Cochlearia. 360.
Colchicum. 206.
Colutea. 403.
Comarum. 288.
Conferva. 634.
Conium. 144.
Convallaria. 193.
Convolvulus. 103.
Conyza. 454.
Coris. 114.
Cornus. 79.
Coronilla. 404.
Corrigiola. 175.
Cortuſa. 99.
Corylus. 523.
Crambe. 353.
Cratægus. 274.
Crepis. 434.
Crithmum. 151.

coccinea. 8. P. multiformis : extus ex flavo pallens ; intus cinnabarina, margine aurantiaco crenulata. *Iacqu. auſtr. t.* 163. *Schæff. fung. t.* 148.

Peziza cava, albida, intus coccinea. *Hall. hiſt. n.* 2222.

In umbroſis humidis innaſcitur putridis ramulis per ſuperficiem terræ dejectis.

Auricula. 9. P. acaulis auriformis rugoſa membranacea fuſca. *Sibthorp oxon. p.* 387.

Tremella Auricula. *Sp. pl. p.* 1625.

Primum genus pernicioſorum fungorum. *Clus. hiſt. p.* CCLXXVI.

Ad arborum truncos.

cochleata. 10. P. turbinata cochleata. *Sp. pl. p.* 1651. *Bulliard herb. t.* 154.

In hortorum umbroſis ad fimeta, in caldariis.

TREMELLA. Fungus gelatinoſus varie expanſus ex tota ſuperficie ſemina emittens.

juniperina. 1. T. ſeſſilis membranacea auriformis fulva. *Syſt. veget. p.* 965.

In junipero communi, Oxycedro.

Noſtoc. 2. T. plicata undulata fuſco-virescens. *Huds. angl.* 2. 564. *Sibthorp oxon. p.* 390.

Tremella terreſtris ſinuoſa, pinguis & fugax. *Dill. musc.* 52. *t.* 10. *f.* 14.

In paſcuis, campis, alibique.

arborea. 3. T. ſeſſilis ſubrotunda undulata nigrescens. *Huds. angl.* 2. 563. *Sibthorp oxon. p.* 390.

Tremella arborea nigricans, minus pinguis & fugax. *Dill. musc.* 54. *t.* 10. *f.* 15.

Ad arborum truncos.

criſpa. 4. T. terreſtris atro-virens tenera criſp: *oxon. p.* 391. *Dill. muſc.* 52. *t.* 10.

In umbroſis humidis.

granulata. 5. T. gelatinoſa ſphærica aggregata vir: *angl.* 2. 566.

Ulva granulata. *Sp. pl. p.* 1633.

Tremella paluſtris, veſiculis ſphæric mibus. *Dill. muſc.* 55. *t.* 10. *f.* 17.

Ad foſſas, lacus, fluvios.

meſenterica. 6. T. membranacea gelatinoſa congeſta rea. *Dicks. faſc.* 1. *p.* 14. *Sibthorp o: Syſt. veget. p.* 965. *Iacqu. miſcell.* 142. *t.* 13.

Ad arborum ramos emortuos, ſti

rufa. 7. T. longitudinaliter dimidiato-infund: *Syſt. veget. p.* 965. *Iacqu. miſcell. ton t.* 14.

In ligno putrido.

hydnoidea. 8. T. compreſſo-plana difformiter digi *Syſt. veget. p.* 965. *Iacqu. miſcell.* 145. *t.* 16.

In ligno putrido.

purpurea. 9. T. ſubgloboſa ſeſſilis ſolitaria glabr *Sp. pl. p.* 1626. *Bulliard herb. t.* 2

Lichenoides tuberculoſum amœne purp *muſc.* 127. *t.* 18. *f.* 6.

Ad arborum truncos, ramos m mortuos.

NIDULARIA. Fungus coriaceus campanulatus in fundo capſulas lentiformes fovens.

campanulata. 1. N. campanulata livido-plumbea lævis: margine patente, capſulis glabris. *Sibthorp oxon. p.* 393.

Peziza lentifera. *Sp. pl. p.* 1649.

In campeſtribus, ad vias.

ſtriata. 2. N. conica fusca: externe villoſa; interne ſtriata, capſulis ſubtus tomentoſis. *Sibthorp oxon. p.* 393. *Bulliard herb. t.* 40. *f. A.*

Ad arborum ſtipites, lignum putridum.

lævis. 3. N. crucibuliformis ſordide lutea: intus lævis, margine erecto, capſulis glabris. *Sibthorp oxon. p.* 393. *Bulliard herb. t.* 40. *f. B. C.*

In ligno putrido.

CLAVARIA. Fungus lævis, oblongus, ſuperficie unica.

Ophioglosſoides. 1. C. clavata integerrima compreſſa obtuſa. *Sp. pl. p.* 1652.

In ſylvis.

cæſpitoſa. 2. C. cæſpitoſa, caulibus ſimpliciſſimis, baſi nonnunquam unitis oblongo clavatis luteolis. *Syſt. veget. p.* 980. *Wulfen in Iacqu. miscell. tom.* 2. *p.* 98. *t.* 12. *f.* 2.

In ſylvis montanis, ſubalpinis.

elveloides. 3. C. cæſpitoſa, caulibus ſimpliciſſimis craſſiſſimis, baſi unitis, obverſe pyramidatis, ſtriatis. *Syſt. veget. p.* 980. *Wulfen in Iacqu. miscell. tom.* 2. *p.* 99. *t.* 12. *f.* 3. *Schæff. fung. t.* 164. *f.* 1.

In ſylvaticis.

digitata. 4. C. ramoſa lignea nigra. *Sp. pl. p.* 1652.

In ſylvis.

Hypoxylon. 5. C. ramoſo-cornuta compreſſa. *Sp. pl.*

In umbroſis ad lignum.

cornuta. 6. C. ramoſa, ramis ſimpliciſſimis, cono no-truncatis. *Syſt. veget. p.* 980. I *Iacqu. miscell. tom.* 2. *p.* 98. *t.* 14. *j*

In ſylvis.

Corralloides. 7. C. ramis confertis ramoſiſſimis inæqua *pl. p.* 1652. *Bulliard herb. t.* 222.

In ſylvis.

faſtigiata. 8. C. ramis confertis ramoſiſſimis faſtigia luteis. *Sp. pl. p.* 1652. *Bolt. fung. t.*

In pascuis ſterilibus, montanis ſyl

crispa. 9. C. foliacea ramoſiſſima, ramis plan per oras ſerratis. *Syſt. veget. p.* 980. *in Iacqu. miscell. tom.* 2. *p.* 110. *t.* 1

In ſylvis.

plebeja. 10. C. ramoſiſſima, ramis ſimpliciusculis apice denticulatis. *Syſt. veget. p.* 98c *in Iacqu. miscell. tom.* 2. *p.* 101. *t.* 1

In ſylvis.

muscoides. 11. C. ramis ramoſis acuminatis inæqualit *Sp. pl. p.* 1652. *Fl. dan. t.* 775. *f.* *fung. t.* 173.

In ſylvis inter muscos.

TUBER. Fungus ſubterraneus carnoſus ſoli ſemina fovens.

cibarium. 1. T. globoſum ſolidum verrucoſum rac tutum. *Sibthorp oxon. p.* 398.

Lycoperdon Tuber. *Sp. pl. p.* 1653.

In pascuis, montanis ſylvaticis.

In Auſtria, Moravia occurrit va tice albo.

LYCOPERDON. Fungus ſubrotundus ſemina farinacea emittens.

** Terreſtria.*

ſtellatum. 1. L. volva multifida patente, capitulo glabro breviter pedunculato, ore acuminato. *Sibthorp oxon p.* 398. *Sp. pl. p.* 1653. *Bolt. fung. t.* 179.

In ſylvis ſubalpinis.

Carpobolus. 2. L. volva multifida, fructu globoſo ex ſeminibus combinatis. *Sp. pl. p.* 1654. *Fl. dan. t.* 895.

Cum præc.

pedunculatum. 3. L. ſtipite longo, capitulo globoſo glabro: ore cylindrico integerrimo. *Sp. pl. p.* 1654. *Bulliard herb. t.* 294.

In campis, pratis, alibique.

verrucoſum. 4. L. ſubgloboſum rufo-fuſcescens, carne cæſio-rubescente, ſeminibus fuscis, collo radicis plicato. *Sibthorp oxon. p.* 399. *Bulliard herb. t.* 24.

In ſylvaticis.

Boviſta. 5. L. ſubrotundum: lacerato-dehiscens. *Sp. pl. p.* 1653.

In apricis herbidis.

*** Paraſitica.*

piſiforme. 6. L. globoſum ſcabrum: ore perforato. *Syſt. veget. p.* 982. *Iacqu. miscell. tom.* 1. *p.* 137. *t.* 7.

In ligno putrido.

ramoſum. 7. L. pyriforme ſcabrum, radice filiformi ramoſiſſima prolifera. *Syſt. veget. p.* 982. *Iacqu. auſtr. t.* 224.

In ſubalpinis ligno putrido innascitur.

Epidendrum. 8. L. cortice farinaque *Fl. dan. t.* 720. *Sch*

In muſcoſis, &

luteum. 9. L. ſubrotundum greg rato fuſco. *Syſt. veg tom.* 1. *p.* 138. *t.* 8

In ſylvis inter

ÆCIDIUM. Capſula utrinqu dis non cohærenti

fuscum. 1. Æ; ſeminibus fuſcis.

In plantarum fc

pini. 2. Æ. oblongo - comp aurantiis. *Syſt. nat.*

In foliis pini.

epiphyllum. 3. Æ. aggregatum, or re fulvo.

Lycoperdon epiphyl

Lycoperdon poculif *p.* 122. *t.* 4. *f.* 1.

In dorſo foliorun rumque plantarum.

cancellatum. 4. Æ. cancellatum filis

Lycoperdon cancell *Iacqu. auſtr. t.* 17.

In averſa pyri

SPHÆRIA. Fungus variæ fi rica, maſſa pulv repleta.

maxima. 1. S. magna craſſa atr *Sibthorp oxon. p.* *Bolt. fung. t.* 181.

Ad lignum put

fragiformis. 2. S. convexa obſcure rubra. *Hoffm. veg. crypt.* 1. *p.* 20. *t.* 5. *f.* 1.

Arborum cortici innaſcitur.

MUCOR. Fungus veſiculis ſtipitatis, in quibus ſemina numeroſa.

Mucedo. 1. M. ſtipitatus, capſula globoſa. *Sp. pl. p.* 1655.

In corporibus putridis.

miniatus. 2. M. ſtipitatus, capſula pyriformi. *Syſt. veget. p.* 982. *Iacqu. auſtr. t.* 299.

In ligno putrido; ſtercore cervino.

glaucus. 3. M. ſtipitatus, capitulo ſubrotundo aggregato. *Sp. pl. p.* 1656.

In Melonibus, pomis, aliisque fructibus corruptis.

cruſtaceus. 4. M. ſtipitatus, ſpicis digitatis. *Sp. pl. p.* 1656.

In cibis corruptis.

Eryſiphe. 5. M. albus, capitulis fuscis ſeſſilibus. *Sp. pl. p.* 1656.

In foliis putridis.

ſepticus. 6. M. unctuoſus flavus. *Sp. pl. p.* 1656.

In ligno putrido.

Arabis. 374.
Arbutus. 224.
Arctium. 438.
Arenaria. 244.
Aretia. 94.
Ariſtolochia. 495.
Arnica. 467.
Artemiſia. 449.
Arum. 497.
Arundo. 61.
Aſarum. 255.
Asclepias. 126.
Asparagus. 193.
Asperugo. 92.
Asperula. 71.
Asphodelus. 191.
Asplenium. 555.
Asprella. 35.
Aſter. 461.
Aſtragalus. 408.
Aſtrantia. 138.
Athamanta. 147.
Atragene. 303.
Atriplex. 545.
Atropa. 117.
Avena. 58.
Azalea. 103.
Ballota. 330.
Bartramia. 586.
Bartſia. 338.
Bellis. 469.
Berberis. 201.
Beta. 129.
Betonica. 327.
Betula. 513.
Bidens. 446.
Biscutella. 364.
Blaſia. 609.
Blitum. 2.
Boletus. 645.
Borago. 92.
Braſſica. 377.
Briza. 48.
Bromus. 54.
Bryonia. 525.
Bryum. 579.
Bunias. 352.
Buphthalmum. 474.
Bupleurum. 139.
Butomus. 219.
Buxbaumia. 600.
Buxus. 514.
Byſſus. 636.
Cacalia. 447.
Cactus. 270.
Calendula. 478.
Calla. 498.
Callitriche. 2.
Caltha. 315.
Campanula. 105.

Camphoroſma. 81.
Cannabis. 533.
Capparis. 289.
Cardamine. 366.
Carduus. 440.
Carex. 502.
Carlina. 445.
Carpeſium. 454.
Carpinus. 522.
Carum. 169.
Caucalis. 143.
Celtis. 547.
Cenchrus. 543.
Centaurea. 475.
Ceraſtium. 252.
Ceratophyllum. 517.
Cerinthe. 91.
Chærophyllum. 164.
Chaiturus. 331.
Chara. 499.
Cheiranthus. 373.
Chelidonium. 289.
Chenopodium. 127.
Cherleria. 247.
Chlora. 210.
Chondrilla. 426.
Chryſanthemum. 469.
Chryſocoma. 448.
Chryſoſplenium. 225.
Cicer. 401.
Cichorium. 436.
Cicuta. 160.
Cineraria. 463.
Circæa. 4.
Ciſtus. 292.
Clathrus. 648.
Clavaria. 652.
Clematis. 304.
Clinopodium. 332.
Clypeola. 363.
Cnicus. 443.
Cochlearia. 360.
Colchicum. 206.
Colutea. 403.
Comarum. 288.
Conferva. 634.
Conium. 144.
Convallaria. 193.
Convolvulus. 103.
Conyza. 454.
Coris. 114.
Cornus. 79.
Coronilla. 404.
Corrigiola. 175.
Cortuſa. 99.
Corylus. 523.
Crambe. 353.
Cratægus. 274.
Crepis. 434.
Crithmum. 151.

Crocus. 24.
Cruciatella. 76.
Crypſis. 19.
Cucubalus. 238.
Cuscuta. 82.
Cyclamen. 100.
Cynogloſſum. 88.
Cynoſurus. 50.
Cyperus. 29.
Cytiſus. 402.
Cypripedium. 495.
Dactylis. 49.
Daphne. 213.
Datura. 116.
Daucus. 144.
Delphinium. 295.
Dentaria. 365.
Dianthus. 235.
Dicranum. 568.
Dictamnus. 220.
Didymodon. 574.
Digitalis. 348.
Dipſacus. 68.
Doronicum. 468.
Draba. 354.
Dracocephalum. 335.
Droſera. 180.
Dryas. 288.
Drypis. 175.
Echinophora. 141.
Echinops. 479.
Echium. 93.
Elaeagnus. 80.
Elatine. 218.
Elymus. 63.
Empetrum. 530.
Encalypta. 567.
Epilobium. 208.
Epimedium. 79.
Equiſetum. 549.
Erica. 212.
Erigeron. 455.
Eriophorum. 33.
Erodium. 379.
Ervum. 401.
Eryngium. 136.
Eryſimum. 371.
Erythronium. 188.
Eupatorium. 447.
Euphorbia. 259.
Euphraſia. 338.
Evonymus. 121.
Fagus. 521.
Ferula. 151.
Feſtuca. 51.
Ficus. 548.
Filago. 478.
Fontinalis. 586.
Fragaria. 282.
Fraxinus. 547.

Fritillaria. 187.
Fucus. 632.
Fumaria. 386.
Funaria. 579.
Galanthus. 182.
Galega. 407.
Galeobdolon. 327.
Galeopſis. 326.
Galium. 72.
Geniſta. 389.
Gentiana. 131.
Geranium. 380.
Geum. 286.
Gladiolus. 25.
Glaux. 124.
Glecoma. 324.
Globularia. 67.
Glycyrrhiza. 403.
Gnaphalium. 452.
Gratiola. 13.
Grimmia. 566.
Gymnoſtomum. 562.
Gypſophila. 233.
Hedera. 123.
Hedyſarum. 406.
Heliotropium. 86.
Helleborus. 314.
Helvella. 648.
Hemerocallis. 196.
Heracleum. 154.
Herniaria. 127.
Hesperis. 374.
Hibiscus. 385.
Hieracium. 428.
Hippocrepis. 405.
Hippophaë. 532.
Hippuris. 1.
Holcus. 542.
Holoſteum. 66.
Hordeum. 64.
Horminum. 335.
Hottonia. 100.
Humulus. 533.
Hyacinthus. 195.
Hydnum. 647.
Hydrocharis. 536.
Hyoscyamus. 116.
Hypnum. 588.
Hyoſeris. 435.
Hypericum. 418.
Hypochæris. 437.
Hyſſopus. 320.
Iaſione. 480.
Iberis. 360.
Ilex. 82.
Illecebrum. 124.
Impatiens. 482.
Imperatoria. 167.
Inula. 464.
Iris. 25.

Iſatis. 353.
Iſnardia. 80.
Iſopyrum. 314.
Juglans. 521.
Juncus. 197.
Jungermannia. 604.
Juniperus. 537.
Ixia. 24.
Lactuca. 425.
Lamium. 325.
Lapſana. 436.
Laſerpitium. 151.
Lathræa. 341.
Lathyrus. 395.
Laurus. 219.
Lavatera. 385.
Ledum. 221.
Lemna. 500.
Leontodon. 428.
Leonurus. 331.
Lepidium. 356.
Leucojum. 182.
Leskea. 597.
Lichen. 610.
Liguſticum. 155.
Liguſtrum. 3.
Lilium. 186.
Limoſella. 349.
Lindernia. 349.
Linum. 177.
Lithoſpermum. 87.
Lolium. 62.
Lonicera. 113.
Loranthus. 201.
Lotus. 415.
Lunaria. 365.
Lychnis. 251.
Lycoperdon. 654.
Lycopodium. 550.
Lycopſis. 92.
Lycopus. 14.
Lyſimachia. 101.
Lythrum. 255.
Malva. 384.
Marchantia. 603.
Marrubium. 330.
Matricaria. 470.
Medicago. 416.
Meeſia. 585.
Melampyrum. 339.
Melica. 43.
Meliſſa. 334.
Melittis. 336.
Mentha. 323.
Menyanthes. 100.
Mercurialis. 535.
Merulius. 644.
Mespilus. 275.
Milium. 39.
Mœhringia. 215.

Momordica. 525.
Monotropa. 221.
Montia. 66.
Morus. 515.
Mucor. 656.
Myagrum. 351.
Myoſotis. 86.
Myoſurus. 181.
Myriophyllum. 517.
Myrtus. 270.
Narciſſus. 183.
Nardus. 34.
Neckera. 599.
Nepeta. 321.
Nidularia. 652.
Nigella. 299.
Nymphæa. 291.
Oenanthe. 159.
Oenothera. 208.
Olea. 3.
Onoclea. 554.
Ononis. 391.
Onopordum. 444.
Onosma. 91.
Ophiogloſſum. 552.
Ophrys. 491.
Orchis. 483.
Origanum. 332.
Ornithogalum. 188.
Ornithopus. 405.
Orobus. 394.
Orobanche. 349.
Osmunda. 553.
Oſyris. 531.
Oxalis. 250.
Pæderota. 12.
Pæonia. 294.
Panicum. 37.
Papaver. 290.
Parietaria. 544.
Paris. 218.
Parnaſſia. 176.
Paſtinaca. 168.
Pedicularis. 341.
Peltaria. 364.
Peplis. 201.
Peucedanum. 149.
Peziza. 649.
Phaca. 407.
Phalaris. 36.
Phallus. 648.
Phascum. 561.
Phellandrium. 159.
Phillyrea. 3.
Phleum. 38.
Phlomis. 331.
Phyſalis. 117.
Phyteuma. 111.
Pimpinella. 170.
Pinguicula. 14.

Pinus. 523.
Piſtacia. 532.
Plantago. 76.
Poa. 44.
Pohlia. 578.
Polemonium. 104.
Polycarpon. 66.
Polycnemum. 23.
Polygala. 387.
Polygonum. 215.
Polypodium. 557.
Polytrichum. 601.
Populus. 534.
Portulaca. 255.
Potamogeton. 83.
Potentilla. 282.
Poterium, 519.
Prenanthes. 427.
Primula. 96.
Prunella. 337.
Prunus. 272.
Pterigynandrum. 587.
Pteris. 554.
Pulmonaria. 89.
Punica. 271.
Pyrola. 223.
Pyrus. 277.
Quercus. 519.
Ranunculus. 308.
Raphanus. 378.
Reſeda. 257.
Rhamnus. 118.
Rhinanthus. 338.
Rhodiola. 535.
Rhododendron. 222.
Rhus. 172.
Ribes. 122.
Riccia. 609.
Roſa. 279.
Rosmarinus. 15.
Rubia. 76.
Rubus. 281.
Rumex. 202.
Ruppia. 85.
Ruscus. 538.
Ruta. 220.
Sagina. 85.
Sagittaria. 518.
Salicornia. 1.
Salix. 526.
Salſola. 129.
Salvia. 15.
Salvinia. 552.
Sambucus. 173.
Samolus. 112.
Sanguiſorba. 78.
Sanicula. 137.
Santolina. 448.
Saponaria. 234.
Satureja. 319.

Satyrium. 489.
Saxifraga. 225.
Scabiola. 68.
Scandix. 161.
Schoenus. 27.
Scilla. 190.
Scirpus. 30.
Scleranthus. 232.
Scolymus. 438.
Scorpiurus. 406.
Scorzonera. 420.
Scrophularia. 347.
Scutellaria. 336.
Sedum. 247.
Selinum. 145.
Sempervivum. 268.
Senecio. 458.
Serapias. 493.
Serratula. 439.
Seſeli. 167.
Sherardia. 70.
Sibbaldia. 180.
Sideritis. 322.
Silene. 239.
Sinapis. 377.
Siſymbrium. 369.
Sium. 157.
Smilax. 534.
Smyrnium. 169.
Solanum. 118.
Soldanella. 99.
Solidago. 463.
Sonchus. 424.
Sorbus. 275.
Sparganium. 501.
Spartium. 388.
Spergula. 254.
Sphæria. 655.
Sphagnum. 561.
Spiræa. 278.
Splachnum. 565.
Stachys. 328.
Stæhelina. 447.
Staphylea. 174.
Statice. 176.
Stellaria. 242.
Stellera. 215.
Stipa. 58.
Stratiotes. 536.
Swertia. 130.
Symphytum. 90.
Syringa. 4.
Tamarix. 174.
Tamus. 533.
Tanacetum. 448.
Taxus. 538.
Tetraphis. 564.
Teucrium. 317.
Thalictrum. 306.
Theſium. 125.

Thlaspi. 357.
Thymus. 332.
Tilia. 292.
Tordylium. 141.
Tormentilla. 286.
Tortula. 575.
Tozzia. 341.
Tragopogon. 420.
Trapa. 80.
Tremella. 650.
Tribulus. 220.
Trichomanes. 560.
Trichoſtomum. 573.
Trientalis. 207.
Trifolium. 411.
Triglochin. 205.
Triticum. 64.
Trollius. 313.
Tuber. 658.
Tulipa. 188.
Turritis. 376.
Tuſſilago. 456.
Typha. 501.
Ulmus. 130.
Ulva. 633.
Urtica. 515.
Utricularia. 14.
Uvularia. 187.
Vaccinium. 211.
Vaillantia. 544.
Valeriana. 20.
Veratrum. 540.
Verbascum. 115.
Verbena. 322.
Veronica. 4.
Viburnum. 172.
Vicia. 397.
Vinca. 125.
Viola. 480.
Viscum. 531.
Vitex. 350.
Vitis. 124.
Weiſia. 577.
Wulfenia. 13.
Xanthium. 516.
Xeranthemum. 453.
Zannichellia. 499.
Zoſtera. 498.

Errata.

p. 25 lin. 20 ieterjacent lege interjacent.
— 33 — 14 Eriphorum — Eriophorum
— 116 — 22 & 28. Hyosciamus — Hyo
— 152 — 21 Arcangelica — Archangelic
— 201 — 25 ſtigmatæ — ſtigmate.
— 274 — 25 ſuſtantia — ſubſtantia.
— 441 — 10 deſtoratus — defloratus.
— 468 — 12 Pardelianches — Pardalianc
— 482 — 9 arivis — arvis.

5